



The International Lily Register and Checklist 2007

First Supplement



© 2009 The Royal Horticultural Society
80 Vincent Square, London SW1P 2PE, United Kingdom
www.rhs.org.uk

International Lily Registrar:
Kate Donald

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright holder

ISBN 9781902896977

Printed and bound in the UK by Page Bros, Norwich

Cover: *Lilium* 'Willcrovidii', drawing of Award of Merit plant by Winifred Walker, 1932.
Image courtesy of RHS Herbarium

The International Lily Register and Checklist 2007 *First Supplement*

| | |
|---|---------|
| Introduction | page 1 |
| Notes on the entries | page 1 |
| Horticultural classification | page 2 |
| Register and Checklist, Oct 2005–Aug 2007 | page 4 |
| Hybridizers, registrants and others | page 62 |

The lily epithets listed here were registered between 1 October 2005 and 31 August 2007. Details of lilies with unregistered names are published also.

Although registration is a voluntary procedure and does not confer any legal protection on the plant, as the International Cultivar Registration Authority for *Lilium*, The Royal Horticultural Society urges all hybridizers, raisers and introducers to register their lily names to reduce the potential confusion caused by new epithets which are the same as, or very similar to, existing names. Epithets which conform to the Articles (and, ideally, the Recommendations) of the latest edition of the *International Code of Nomenclature for Cultivated Plants* (ICNCP) are acceptable for registration. Registration, for which a certificate is issued on request, is free of charge, and should take place before a plant is released or described in a catalogue.

Registration forms are available as a download from RHS Online: www.rhs.org.uk; from Mrs Kate Donald, International Lily Registrar, 16 Midtown of Inverasdale, Poolewe, Ross-shire IV22 2LW, UK; e-mail: lily@rhs.org.uk; tel: +44 (0)1445 781717; and from the Regional Representatives listed below.

The Royal Horticultural Society is indebted to the following Regional Representatives for their continued efforts to encourage the registration of lily names and for contributing amendments to *The International Lily Register and Checklist* and its *Supplements*:

• **Australia:**

Mr Neil Jordan, 27 Scarfe Street, Camdale, Tasmania 7320
E-mail: neil.jordan@gotalk.net.au

• **Czech Republic:**

Ing. Břetislav Mičulka, 687 07 Velehrad 225
E-mail: miculka.b@tiscali.cz

• **The Netherlands:**

Mrs Saskia Bodegom, Koninklijke Algemeene Vereeniging voor Bloembollencultuur, Postbus 175, 2180 AD Hillegom
E-mail: bodegom@kavb.nl

• **New Zealand:**

Mrs Merle I. Lepper, P.O. Box 1394, Christchurch
E-mail: merle.lepper@xtra.co.nz

• **North America:**

Mrs Kathy Andersen, 2565 Charlestown Road, Phoenixville, Pennsylvania 19460, USA
E-mail: ksa2006@verizon.net

Notes on the entries

The format of entries is similar to the layout adopted for *The International Lily Register and Checklist 2007*.

Parentage

The seed parentage is given first, preceding the larger multiplication sign.

Roles

The sequence and explanation of the abbreviations is as follows:

S: Selected by,

H: Hybridizer (reinstated in favour of the rather ambiguous term, Raiser),

G: Grown to first flowering by,

N: Named by,

I: Introduced by,

REG: Registrant.

Colour

Numbers and letters in parentheses refer to the *RHS Colour Chart* (1966 (reprinted 1986), 1995, 2001 and 2007). A colour name preceding a *Colour Chart* reference is taken from *A Contribution toward Standardization of Color Names in Horticulture: Application of the Universal Color Language to the Colors of the Royal Horticultural Society's Colour Chart* by R.D. Huse and K.L. Kelly, edited by D.H. Voss, The American Rhododendron Society, 1984.

Nomenclatural Standard

Sharp colour images, clearly showing the characteristics of the plant(s), and/or, if practicable, pressed specimens, are very helpful and greatly appreciated. These, with the completed form, constitute a permanent, definitive record (nomenclatural standard) of the cultivar or Group. Those held in the RHS Herbarium at RHS Garden Wisley are under a designated WSY reference number.

The Registrar would like to thank her predecessor, Victoria Matthews, and colleagues in the Botany Department and Library at Wisley for their help and advice – particularly Dr Alan Leslie – and most especially, Duncan Donald, the International Clematis Registrar.

• **Further information to augment existing lily records is always welcome.**

Horticultural classification

Division I Asiatic hybrids

Hybrids derived from the following species and interspecific hybrids: *amabile*, *bulbiferum*, *callosum*, *cernuum*, *concolor*, *dauricum*, *dauidii*, *L. × hollandicum*, *lancifolium* (syn. *tigrinum*), *lankongense*, *leichtlinii*, *L. × maculatum*, *pumilum*, *L. × scottiae*, *wardii* and *wilsonii*.

Flowers usually small to medium-sized. Flower colour often uniform, or with contrasting tepal tips and/or throat. Spots absent, or when present, well-defined and often rounded. Conspicuous brush-marks sometimes present. Tepal margins usually smooth or slightly ruffled. Flowers usually with little or no scent.

Examples: 'Ariadne' I(c/d), Citronella Group I(c/d), 'Mont Blanc' I(a/b-c).

Division II Martagon hybrids

Hybrids of martagon type derived from the following species and interspecific hybrids: *L. × dalhansonii*, *hansonii*, *martagon*, *medeoloides* and *tsingtauense*.

Flowers usually small, often numerous, mostly down-facing, typically with rather thick, recurved tepals and often of Turk's cap form. Spots usually present, numerous, often on at least three-quarters of each tepal. Flowers with little or an unpleasant scent. Buds often hairy. Bulbs often mauve or orange-yellow. Early flowering.

Examples: 'Cadense' II(b/d), 'Claude Shride' II(c/d), *L. × dalhansonii* 'Marhan' II(c/d).

Division III Euro-Caucasian hybrids

Hybrids derived from the following species and interspecific hybrids: *candidum*, *chalcedonicum*, *kesselringianum*, *monadelphum*, *pomponium*, *pyrenaicum* and *L. × testaceum*.

Flowers mostly small to medium-sized, often bell-shaped to Turk's cap-shaped, often down-facing. Flower colour often in rather pale muted shades. Spots absent to numerous. Tepal-margins smooth, often gently reflexed. Flowers often scented. Many are lime tolerant.

Examples: Caucasian Group III(c/d), 'Moonlight Madonna' III(-/a), 'Zeus' III(c/d).

Division IV American hybrids

Hybrids derived from the following American species and interspecific hybrids: *bolanderi*, *L. × burbankii*, *canadense*, *columbianum*, *grayi*, *bumboldtii*, *kellyanum*, *kelloggii*, *maritimum*, *michauxii*, *michiganense*, *occidentale*, *L. × pardaboldtii*, *pardalinum*, *parryi*, *parvum*, *philadelphicum*, *pitkinense*, *superbum*, *vollmeri*, *washingtonianum* and *wigginsii*.

Flowers small to medium-sized, mostly down-facing. Flower colour often strong yellow to orange or orange-red, often with contrasting centre and tepal tips. Spots often very conspicuous, over at least half of each tepal, often surrounded by a paler halo. Tepals rather narrow, margins smooth, usually gently to strongly reflexed. Flowers with little scent.

Examples: Bellingham Group IV(c/d), 'Lake Tulare' IV(c-c/d), 'Shuksan' IV(c/d).

Division V Longiflorum lilies

Hybrids or selections derived exclusively from *formosanum*, *longiflorum*, *philippinense* and *wallichianum*.

Flowers medium-sized to large, often few, trumpet-shaped. Flower colour typically uniform (white). Spots, papillae and brushmarks absent. Tepal margins smooth. Flowers usually scented.

Examples: Formolongi Group V(-/a), *longiflorum* 'Slocum's Ace' V(b/a).

Division VI Trumpet and Aurelian hybrids

Hybrids derived from the following species and interspecific hybrids: *L. × aurelianense*, *brownii*, *L. × centigale*, *henryi*, *L. × imperiale*, *L. × kewense*, *leucanthum*, *regale*, *rosthornii*, *sargentiae*, *sulphureum* and *L. × sulphurgale* (but excluding hybrids of *henryi* with all species listed in Division VII). Aurelian hybrids are derived from a combination of *henryi* and trumpet lilies.

Flowers medium-sized to large, in all flower forms. Flower colour white, cream, yellow to orange or pink, often with a contrasting star-shaped throat and/or strong bands of colour outside. Trumpets usually scented, without markings; other types often with spotting as small streaks in two bands at the base. Tepal-margins smooth or twisted with irregular ruffling, tips often reflexed.

Examples: Herald Angel Group VI(a/a), *L. × kewense* 'White Henry' VI(b-c/c).

Division VII Oriental hybrids

Hybrids derived from the following species and interspecific hybrids: *auratum*, *japonicum*, *nobilissimum*, *L. × parkmanii*, *rubellum* and *speciosum* (but excluding all hybrids of these with *henryi*).

Flowers usually medium-sized to very large. Inner tepals often very broad and overlapping at the base to give a 'closed' centre. Flower colour mostly white to pink to purplish red, some golden yellow; ground colour often white, with a contrasting central ray. Spots absent to numerous, sometimes over half of each tepal. Papillae and nectaries often conspicuous. Flowers usually scented. Often late-flowering.

Examples: *L. × parkmanii* Imperial Silver Group VII(b/c), 'Woodriff's Memory' VII(a-b/b).

Division VIII Other hybrids

Hybrids not covered by any of the previous Divisions, I-VII. Includes all interdivisional hybrids, such as Asiatic/Trumpet (Asiapets or AT) hybrids, *longiflorum*/Asiatic (LA) hybrids, *longiflorum*/Oriental (LO) hybrids, Oriental/Asiatic (OA) hybrids and Oriental/Trumpet (Oriepets or OT) hybrids.

Hybrids of *henryi* with *auratum*, *japonicum*, *nobilissimum*, *L. × parkmanii*, *rubellum* and *speciosum* (excluded from Divisions VI and VII) are placed here.

Examples: 'Oberhof' VIII(b-c/-) AT, 'Salmon Classic' VIII(a-b/b) LA, 'Sky Treasure' VIII(b/a) LO, 'Sunny Crown' VIII(a/b) OA, 'Yelloween' VIII(a/b) OT.

Division IX Species and cultivars of species

Includes all species and their subspecies, varieties and forms, and cultivars selected therefrom (excluding those derived exclusively from *formosanum*, *longiflorum*, *philippinense* and *wallichianum*, which are placed in Div. V).

Examples: *auratum* var. *rubrovittatum* 'Crimson Beauty' IX(b/c); *canadense* var. *editorum* 'Chocolate Chips' IX(c/a); *lancifolium* 'Splendens' IX(c/d); *nepalense* IX(c/a).

Sub-divisions

If desired, it is possible to further classify lilies of all Divisions according to flower aspect and form. Flower aspect is given first and the two are separated by a forward slash (/).

Flower aspect:

- a** up-facing;
- b** out-facing;
- c** down-facing

Flower form:

- a** trumpet-shaped;
- b** bowl-shaped;
- c** flat (or with tepal tips recurved);
- d** tepals strongly recurved (with the Turk's cap form as the ultimate state).

Examples: 'Madison' I(a/b) indicates an Asiatic hybrid with up-facing, bowl-shaped flowers; 'Lightning Bug' II(b/d) a martagon hybrid with out-facing flowers with strongly recurved tepals; 'Snowdon' VII(a-b/-) an Oriental hybrid with flowers varying from up- to out-facing, but flower form unknown; *speciosum* 'Crimson Glory' IX(b-c/d) a selection from *speciosum* with out- to down-facing flowers with strongly recurved tepals.

Register and Checklist, Oct 2005–Aug 2007

‘Adriana’ I(a-b/b)

Unknown × (unknown × ‘Blizzard’)

H: B. Mićulka, pre-2006 **N:** B. Mićulka, pre-2006

I: B. Mićulka, 2006 **REG:** B. Mićulka, 2006

Inside of tepals pink, central portion pale apricot, darker over nectaries; throat deep pink at base.

Outside of tepals pale pink, midribs greenish, hairy. Spots small, oblong, dark brown, on a third of each tepal, fairly numerous, but fewer on outer tepals; nectaries pink-granulated; pollen orange-brown; stigma dark old rose. Fls 165 mm wide; tepals 95 × 43 mm, tips slightly recurved; pedicels 70 mm, green, stout. Lvs 160 × 14 mm, dull deep green, dense. Stems 0.85 m, green, stout, with fls in a whorled infl. of medium length. Mid-June.

‘After Eight’ VII(a/b)

(*L. × parkmanii* ‘Allegra’ × ‘Star Gazer’) × un-named seedling) × un-named seedling

H: Bischoff Tulleken Lelies BV, pre-2001

N: Bischoff Tulleken Lelies BV, 2001

REG: Mak Breeding BV, 2006

Inside of tepals strong purplish red (60C/60D), shading to yellowish white (155D) towards margins and pale greenish yellow (1D) along edge of nectaries; midveins deep red (60A); throat yellowish white (155D). Outside of outer tepals moderate purplish red (186B); outside of inner tepals dark purplish pink (186C); base and margins of each tepal yellowish white (155D); in outer tepals in upper two-thirds only; midrib of each tepal strong yellow-green (144C) at base, shading to yellowish white (155D); midribs of outer tepals strong yellow-green (144A) at top. Spots and most papillae dark red (59A); a few papillae pale yellow-green (155A); nectaries strong yellow-green (144C); pollen moderate orange (172D); stigma pinkish grey (201D). Fls 140–220 mm wide, scented; tepals 116 × 68 mm, margins slightly ruffled, tips recurved. Lvs 140 × 47 mm. Stems 0.45–0.7 m, green with darker markings, with c.8 fls.

‘Agneta’ VII(a/b)

‘Melbourne’ × white-flowered TL seedling

H: De Jong Beheer BV, pre-1999 **N:** De Jong Lelies Holland BV, 1999 **REG:** De Jong Lelies Holland BV, 2007

Inside of tepals yellowish white (155D), flushed pale greenish yellow (2D) at base; midveins light greenish yellow (1C) from above nectaries to up to half the length of each tepal; throat yellowish white (155D). Outside of outer tepals yellowish white (155D); midribs strong yellow-green (144C) at base and in upper half, shading to pale greenish yellow (2D) between. Outside of inner tepals yellowish white (155D), flushed brilliant greenish yellow (2B) at base; midribs light yellow-green (144D) at base and top, shading to pale greenish yellow (2D) between. Spots absent; papillae few, yellowish white (155D); nectaries light yellow-green (144D); pollen vivid orange-yellow (21A); stigma pale yellow-green (193D). Fls 250 mm wide, scented;

tepals 135 × 65 mm, margins ruffled, tips recurved. Lvs 185 × 48 mm. Stems 1 m, green, with up to 3 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Akemi’ VII(b-a/b-c)

‘Towny’ × ‘Sorbonne’

H: De Jong Lelies Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2006

Inside of inner tepals deep purplish pink (N66D), strong purplish red (64C) towards margins; inside of outer tepals deep purplish pink (N66C); midveins deep yellowish pink (47C), with a little yellowish white (155D) at top; throat yellowish white (155D). Outside of outer tepals strong purplish pink (63C), a little yellowish white (155D) beside midribs; midribs light yellow-green (150D) shading to strong yellow-green at base and top (143C and 143A respectively). Outside of inner tepals light purplish pink (63D) shading to deep purplish pink (64D) towards margins and yellowish white (155D) at base; midribs strong yellow-green (144C) at base, shading to pale yellow-green (155A), and light yellow-green (145B) at top. Spots strong purplish red (58B); papillae yellowish white (155D) and strong purplish red (58B); nectaries strong yellow-green (144C); pollen strong orange (N25A); stigma moderate yellow-green (138C). Fls 175 mm wide, becoming paler with age; tepals 116 × 60 mm, margins ruffled, tips recurved. Lvs 150 × 52 mm. Stems 1 m, green, with c.6 fls.

‘Alanno’ VIII(a/b) LA

Seedling BG 95-60 × seedling AJ 94-15

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals brilliant yellow (21C), strong orange (24A) towards margins; inside of outer tepals brilliant orange-yellow (21B), strong orange (25A) towards margins; throat pale yellow (8D). Outside of outer tepals light orange-yellow (22B), strong orange (24A) towards margins and a broad stripe of brilliant yellow-green (142B) along midribs. Outside of inner tepals light orange-yellow (22C) shading to 23C towards margins; midribs strong yellow-green (144C). Spots very few, dark red (59A); papillae strong orange (24A); nectaries strong yellow-green (143C); pollen dark red (187A); stigma moderate reddish orange (178C). Fls 190 mm wide; tepals 118 × 59 mm, margins smooth, tips slightly recurved. Lvs 162 × 18 mm, dark green. Stems 1–1.5 m, green with darker markings, with c.6 fls.

‘Alaska’ I(a-b/b)

Syn. of ‘Atoll’.

In 1993, ‘Atoll’ was granted a US Plant Patent under the cultivar epithet ‘Alaska’, which is the accepted name in countries in which ‘Alaska’ has legal protection. However, ‘Atoll’ was granted Dutch Plant Breeders’ Rights in 1990 and is, therefore, the prior name accepted by the ICRA for registration. In *ILL&C 2007* ‘Alaska’ appears as a trade designation.

'Albisola' VII(a-b/b)

Seedling 93-47 × seedling RM 94-01

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals deep purplish pink (N57C), shading to strong purplish pink (68B) towards margins; midveins vivid purplish red (N57B) at top. Inside of outer tepals deep purplish pink (N57D); midveins vivid purplish red (N57B); each tepal pale yellow (11D) beside nectaries, and above nectaries of inner tepals; throat yellowish white (155D). Outside of tepals deep purplish pink (N57D), shading to N57C towards margins of outer tepals only; all tepals yellowish white (155D) at base; midribs of outer tepals light yellow-green (145B) at base shading to yellowish white (155D); midribs of inner tepals strong yellow-green (144C) at base shading to pale yellow-green (155A); all midribs strong yellow-green (144B) at top. Spots strong purplish red (60D); papillae yellowish white (155D) and strong purplish red (60D); nectaries light yellow-green (145B); pollen dark orange (N163A); stigma light grey (188D). Fls 200 mm wide, scented, becoming more bluish with age; tepals 125 × 76 mm, margins ruffled, tips recurved. Lvs 185 × 43 mm. Stems 1.1 m, green with darker markings at leaf nodes, with c.6 fls.

'Aleš' I(a/c)

('Olga' (Jošt) × 'Windahloo') × ('Rusalka' × unknown)

H: B. Mičulka, pre-2005 **N:** B. Mičulka, pre-2005**I:** B. Mičulka, 2006 **REG:** B. Mičulka, 2005

Inside of tepals glowing yellow; outside of tepals paler yellow; midribs greenish. Brushmarks deep mahogany, fairly large; spots dark mahogany, fairly large, lightly distributed over a quarter of each tepal; nectaries white-felted; pollen brown; stigma brownish and pinkish. Fls 150 mm wide; tepals 83 × 35 mm, margins slightly ruffled, tips only very slightly recurved; pedicels 80 mm. Lvs 110 × 16 mm, dull dark green. Stems 0.95 m, green, stout and rigid. Late June.

'Alfa' (De Jong) I(a/b)

Tetraploid Asiatic × Asiatic

H: De Jong Lelies Beheer BV, pre-2005 **N:** De JongLelies Holland BV, 2005 **REG:** De Jong Lelies

Holland BV, 2006

Inside of tepals vivid yellow (12A); throat brilliant greenish yellow (2B). Outside of outer tepals brilliant greenish yellow (151D) shading to vivid yellow (12A) towards margins; midribs strong yellow-green (144C) at base, flushed dark red (59A) and wholly 59A at top; outside of inner tepals vivid yellow (12A), brilliant greenish yellow (1B) at base; midribs strong yellow-green (144C). Spots absent; papillae vivid yellow (12A); nectaries moderate olive-green (137A); pollen strong orange (169B) and dark reddish orange (172B); stigma brilliant yellow (13B). Fls 180 mm; tepals 102 × 54 mm, margins smooth, tips slightly recurved. Lvs 130 × 15 mm. Stems 0.85–0.95 m, dark with a few green markings, with c.8 fls.

'Alka' VI(b-c/c-d)

(('Lady Alice' × unknown) × (unknown × 'Toni')) × unknown

H: B. Mičulka, pre-2006 **N:** B. Mičulka, pre-2006**I:** B. Mičulka, 2006 **REG:** B. Mičulka, 2007

Tepals pale beige-apricot; central portion darker; tips whitish; midribs pale green. Spots absent; papillae apricot, over half of each tepal; nectaries green, broad; pollen apricot-ochre; stigma green. Fls 135 mm wide; tepals 95 × 37 mm, stiff, tips strongly recurved; pedicels slender, sparsely spotted brown; with secondary buds. Lvs 135 × 19 mm, dull dark green. Stems 1.3 m, stout, green, infl. of average height. Mid-July.

'All is All' VIII(a-b/-) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Inside of inner tepals vivid yellow (9A/B), inside of outer tepals 9A. Outside of inner tepals brilliant yellow (9C) to pale greenish yellow (9D). Spots absent; papillae present; nectaries green; pollen red-brown; stigma dark purple, spotted. Tepals of short to medium length and of narrow to medium width, margins ruffled, tips slightly recurved to recurved. Lvs of medium length and width to broad. Stems tall to very tall, green with darker stripes and spots, with few to average no. of fls. Late to very late.

'Allysee' VIII(a-b/b) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007

Tepals pale greenish yellow (2D); midveins green at base, shading to yellow. Outside of outer tepals tinged purple at base; young fls more whitish. Spots absent; papillae present; nectaries green; pollen red-brown; stigma grey-green. Tepals of medium length and width, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and of medium width to broad. Stems of average height to tall, green, with up to 6 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Alois' I(b/b)

'Dawn Star' × 'Feuer und Rauch'

H: J. Dudek, pre-2006 **G:** J. Dudek, pre-2006**N:** J. Dudek, pre-2006 **I:** J. Dudek, 2006**REG:** J. Dudek, 2006

Inside of tepals white, with greenish centre and throat. Outside of tepals pale yellow-green; midribs pale. Spots purple, numerous, fairly large, on half of each tepal; papillae absent; nectaries green; pollen brown; stigma yellowish. Fls 160 mm wide; tepals 80 × 30 mm, tips slightly recurved; pedicels 80 mm; with secondary buds. Lvs 130 × 12 mm, scattered, glossy. Stems 1.2 m, pale green; infl. of average height. Mid-July.

'Altesino' VII(a/b)

Seedling 88-88 × seedling 91-62

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of tepals deep purplish red (61A) shading to moderate purplish red (70A) towards margins; margins yellowish white (155D); midveins deep purplish red (59B); throat pale greenish yellow (2D). Outside of tepals moderate purplish red (64A) shading to yellowish white (155D) towards margins and base; midribs of outer tepals strong yellow-green (144B) at base, with a small stripe of moderate purplish red (64A), shading to 64A with yellowish white (155D), and strong yellow-green (144A) at top. Midribs of inner tepals yellowish white (155D) from base to up to half the length of the tepal, and moderate yellow-green (147D) at top. Spots dark red (59A); papillae moderate purplish red (64A); nectaries strong yellow-green (144B); pollen strong brown to moderate reddish orange (172A/N172A); stigma greyish reddish purple (N77C). Fls 220 mm wide, slightly scented; tepals 135 × 70 mm, margins slightly ruffled, tips recurved. Lvs 190 × 40 mm. Stems 1–1.2 m, light green with darker markings at leaf nodes, with c.3 fls.

'Alusta' VIII(a-b/-) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Inside of inner tepals vivid yellow (9A/B); inside of outer tepals and outside of inner tepals vivid to brilliant yellow (9B/C); midveins yellow-green at base. Spots absent; papillae present; nectaries green; pollen red-brown; stigma grey-green. Tepals of medium length to long and of medium width, margins slightly ruffled, tips recurved. Lvs of medium length to long and of medium width. Stems tall, green with faint darker stripes and spots, with few fls. Late to very late.

'Amaril' V(b/a)

Parentage unknown

H: J.W.A. van der Wereld BV, pre-2005
REG: J.W.A. van der Wereld BV, 2005

Inside of tepals and outside of inner tepals yellowish white (155B); midveins green at base. Spots and papillae absent; nectaries green; pollen orange-yellow; stigma white-grey. Tepals long to very long, broad, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length to long and of medium width to broad. Stems tall, green, with few fls. Late.

'Antares' VII(a-b/b-c)

Seedling 88-99 × seedling RM 95-34

H: Vletter en Den Haan Beheer BV, pre-2005
N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2007
Inside of tepals strong purplish pink (73B), shading to moderate purplish pink (68C) towards margins; margins of upper half of each tepal, midveins above nectaries, and throat, yellowish white (155D). Outside of outer tepals strong purplish pink (73B), shading to deep purplish pink (73A) towards margins and yellowish white (155D) beside midribs; midribs strong yellow-green (N144C at base and top; N144D between). Outside of inner tepals moderate purplish pink (186D); margins towards tips, and base beside midribs, yellowish white (155D); midribs strong yellow-green (N144C) at base and top, shading to

yellowish white (155D) between. Spots vivid purplish red (N57B); papillae yellowish white (155D); nectaries strong yellow-green (N144C); pollen orange-brown; stigma pale green (189C). Fls 250 mm wide, scented; tepals 135 × 75 mm, margins slightly ruffled, tips recurved. Lvs 225 × 64 mm. Stems 1–1.4 m, green, with up to 4 fls.

Syn. 'Alcazar'

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Antequera' I(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2007

Inside of inner tepals strong orange-yellow (17A/N25D); midveins strong orange (N25C); throat vivid yellow (14B). Inside of outer tepals strong orange (N25C) to strong orange-yellow (N25D); tips 17A. Outside of outer tepals brilliant yellow (13B); margins strong orange-yellow (17A); midribs flushed brilliant yellow-green (154C); strong yellow-green (144C) at top. Outside of inner tepals brilliant yellow (14C); margins strong orange-yellow (17A); midribs vivid yellow (12A). Spots and papillae absent; nectaries strong yellow-green (144A); pollen dark reddish orange to brownish orange (172B to 172C); stigma strong yellow-green (144C) with moderate red (179B) stripes. Fls 190 mm wide; tepals 105 × 52 mm, margins smooth, tips recurved. Lvs 180 × 14 mm, dark green. Stems 1.1–1.3 m, green, with up to 10 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Anza' VII(a/b)

Parentage unknown

H: World Breeding BV, pre-2001 **N:** World Breeding BV, 2001 **REG:** World Breeding BV, 2007

Inside of tepals strong purplish red (63A), paler towards tips; becoming more blue with age. Outside of inner tepals strong purplish red (64B/C). Spots and papillae present; nectaries yellow-green; pollen orange-brown; stigma light purple. Tepals of short to medium length and of medium width, margins ruffled, tips slightly recurved to recurved. Lvs of medium length and broad. Stems of average height to tall, green with darker spots and stripes, with few to average no. of fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Apricot Pixels' I(a/b)

'Cha Cha Cha' × 'Bright Beauty'

H: Laan Flora Facilities BV, pre-2005 **N:** Laan Flora Facilities BV, 2005 **REG:** Laan Flora Facilities BV, 2006

Inside of tepals vivid yellow (13A) shading to strong orange (24A) towards tips of inner tepals and margins of outer tepals; margins of inner tepals vivid yellow (15A); throat vivid yellow (16A). Outside of outer tepals brilliant yellow (13C) shading to strong orange-yellow (17A) towards margins; midribs strong yellow-green (143C) at base shading to 144C on

yellow ground, and strong yellow-green (143A) at top, with spot of greyish purple (N77A). Outside of inner tepals light yellow (14D) shading to vivid yellow (17B) towards margins; midribs moderate yellow-green (139D) at base shading to strong yellow-green (144C), and dark red (59A) at top. Spots dark red (59A), numerous, mostly distributed above throat and along inside margins of tepals, fewer along outside margins; papillae absent; nectaries moderate yellow-green (148A); pollen strong reddish orange (169A); stigma deep red (60A). Fls 115 mm wide, becoming paler with age; tepals 70 mm × 38 mm, margins smooth, tips straight. Lvs 160 × 15 mm. Stems 0.8–1.1 m, green, with *c.* 6 fls.

'Aprilia' VII(b-a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of inner tepals deep purplish red (59B); midveins dark red (59A); throat yellowish white (155D). Inside of outer tepals moderate purplish red (59C), tinged yellowish white (155D) at tips; midveins deep purplish red (59B). Outside of outer tepals deep purplish red (59B), yellowish white (155D) at base beside midribs; midribs strong yellow-green (N144D) at base and tinged moderate olive-green (146A) at top. Outside of inner tepals moderate purplish red (59C), shading to strong purplish red (60B) towards margins, and yellowish white (155D) at base beside midribs; midribs strong yellow-green (N144C) at base and tinged yellowish white (155D) at top. Spots and papillae dark red (187A and 59A respectively); nectaries strong yellow-green (N144C); pollen strong orange (N163B); stigma dark greyish red (N186C). Fls 205 mm wide, scented; tepals 115 × 60 mm, margins slightly ruffled, tips recurved. Lvs 215 × 47 mm. Stems 1.15–1.2 m, green with darker markings at base, with up to 6 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Arcachon' VIII(a/b) LA

Seedling BW 97-45 × seedling W 98-11

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals greenish white (157D); inside of outer tepals yellowish white (155D) flushed light yellow-green (145D), margins yellowish white (155D); throat light yellow-green (145C). Outside of outer tepals yellowish white (155D) flushed light yellow-green (145D); midribs light yellow-green (142C) shading to strong yellow-green (143B) at top and base; outside of inner tepals yellowish white (155D); midribs light yellow-green (144D) at base shading to 145D. Spots moderate purplish red (58A), few; papillae yellowish white (155D); nectaries strong yellow-green (144A); pollen strong reddish orange (169A), scarce; stigma brilliant greenish yellow (1B). Fls 190 mm wide; tepals 115 × 50 mm, margins smooth, tips recurved. Lvs 140 × 23 mm. Stems 1.1–1.4 m, green, with *c.* 6 fls.

'Argeles' VII(b-a/c)

Seedling 93-25 × seedling RM 96-26

H: Vletter en Den Haan Beheer BV, pre-1993

N: Vletter en Den Haan Beheer BV, 1993

REG: Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals vivid purplish red (61C) shading to strong purplish red (64C) towards margins; a small patch of light yellow-green (150D) above nectaries. Inside of outer tepals strong purplish red (67A) shading to deep purplish red (71A) towards margins; very pale purple (69D) at base; throat yellowish white (155D). Outside of outer tepals strong purplish red (64B) shading to moderate purplish red (64A) towards margins; midribs dark greenish yellow (152C) shading to strong yellow-green at base and top (143C and 143A respectively). Outside of inner tepals deep purplish pink (64D) to strong purplish red (64C); yellowish white (155D) at base; midribs strong yellow-green (143C) at base shading to strong yellow (153D). Spots strong purplish red (58B); papillae yellowish white (155D) and light purplish pink (65B); nectaries strong yellow-green (143C); pollen dark orange (N163A); stigma bluish white (112D). Fls 200 mm wide; tepals 115 × 60 mm, margins ruffled, tips recurved. Lvs 190 × 47 mm. Stems 1.1–1.3 m, green, with *c.* 3 fls.

'Armanda' VIII(b-c/a-b) LO

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2001

N: Marklily CV, 2001

REG: Testcentrum voor Siergewassen BV, 2007

Tepals greenish white (155C); deep purplish pink (N57D) beside base of midveins; midveins yellow-green at base. Spots and papillae absent; nectaries green; pollen orange-brown; stigma light grey. Tepals long to very long and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs long and of medium width to broad. Stems very tall, green with darker markings, with few fls. Late to very late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Arrow' I(a-b/b)

Parentage unknown

H: De Vries Veredeling BV, pre-2003

N: De Vries Veredeling BV, 2003

REG: De Vries Veredeling BV, 2007

Tepals vivid reddish orange (inside of tepals N30C, outside N30B); base of midveins strong orange (N25B). Spots few; papillae present; nectaries orange; pollen orange-brown; stigma orange. Tepals short to very short and narrow, margins (very) slightly ruffled, tips slightly recurved to recurved. Lvs of short to medium length, and of narrow to medium width. Stems 1–1.2 m, green with darker markings, with up to 10 fls. Mid-season.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Atlantico' VII(a-b/b-c)

Parentage unknown

H: World Breeding BV, pre-2003

N: Liliande BV, 2003

REG: World Breeding BV, 2007

Tepals deep purplish pink (N66C); base of midveins brilliant greenish yellow (4A). Spots numerous; papillae present; nectaries green; pollen orange-brown; stigma grey-green. Tepals of short to medium length and of medium width, margins ruffled, tips recurved. Lvs of medium length to long, and of medium width to broad. Stems 1.3 m, green with darker speckles and stripes, with up to 7 small to medium-sized fls. Mid-season to late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Atvasara'

Unregistered pollen parent of 'Priedula'. No further details known.

'Avanti' I(a/-)

Parentage unknown

H: Lybelmex Hillegom BV, pre-1993

REG: Lybelmex Hillegom BV, 2007

Inside of tepals strong yellowish pink (37B), flushed yellow at base and red-pink towards tips. Outside of tepals moderate pink (49B). Spots and papillae present; nectaries green; pollen and stigma orange.

Tepals short to very short and of narrow to medium width, margins (very) slightly ruffled, tips very slightly recurved. Lvs of short to medium length, and of narrow to medium width. Stems of average height, green, with few to average no. of fls.

'Avocado' VIII(a/b) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2006

Inside of inner tepals brilliant orange-yellow (23B) shading to light orange-yellow (23C) towards margins; tips light yellow (11B); midveins vivid orange-yellow (23A). Inside of outer tepals vivid yellow (13A) shading to light orange-yellow (16B) towards margins; midveins brilliant orange-yellow (23B); throat yellowish white (155D). Outside of outer tepals pale yellow (11D); margins light yellow (14D); midribs strong yellow-green (143C) at base shading to yellowish white (N155D) and light yellow (14D) at top. Outside of inner tepals brilliant yellow (15C) on pale yellow ground (11C); margins light orange-yellow (23C); midribs strong yellow-green (143C) at base shading to pinkish white (N155C). Spots and papillae absent; nectaries strong yellow-green (143B); pollen strong brown (172A); stigma very pale purple (85D) with a little very light purple (85C). Buds flushed strong reddish orange (N34B). Fls 195 mm wide; tepals 135 × 48 mm, margins slightly ruffled, tips recurved. Lvs 220 × 35 mm. Stems 1.2–1.55 m, dark green, with darker markings at base of lvs, with c.7 fls.

'Badessa' VIII(a/b) LA

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals strong orange-yellow (24B); margins and above throat strong orange (25A); throat pale

greenish yellow (2D). Outside of outer tepals light orange-yellow (19A) shading to strong orange (25B) towards margins; midribs greyish red (182B) at base shading to yellowish grey (194C), and strong yellow-green (143A) at top. Outside of inner tepals light orange-yellow (24C) shading to brilliant orange (25C) towards margins; midribs greyish red (182B) at base shading to greyish yellow-green (194A), and strong yellow-green (144B) at top. Spots and papillae absent; nectaries strong yellow-green (144A); pollen moderate reddish brown (175A); stigma deep purplish red (59B). Fls 180 mm wide; tepals 110 × 56 mm, margins smooth, tips very slightly recurved. Lvs 170 × 28 mm. Stems 1.2–1.6 m, green with darker markings, with c.5 fls.

'Bagnara' VIII(a/b) LA

Seedling BW 96-21 × seedling W 92-11

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Tepals yellowish white (155D); throat light yellow-green (145C); midribs of outer tepals strong yellow-green at base and top (143C and 144C respectively); midribs of inner tepals strong yellow-green (143C) at base flushed dark greenish yellow (152D). Spots and papillae absent; nectaries strong yellow-green (143A); pollen strong orange (N25B); stigma light greenish yellow (3C). Fls 175 mm wide; tepals 100 × 50–60 mm, margins smooth, tips recurved. Lvs 170 × 20 mm. Stems 1–1.4 m, dark with green markings, with c.4 fls. Resembles an Asiatic (Div. 1) hybrid.

'Bankir' I(b/d)

('Rondo' × Sutter's Gold Group) × unknown

H: K. Braun, pre-2006 **N:** K. Braun, pre-2006

I: K. Braun, 2006 **REG:** K. Braun, 2006

Tepals yellow, throat and midribs yellow-green. Spots dark brown, large, oval, over three-quarters of each tepal; papillae absent; nectaries green; pollen dark brown; stigma yellow-green. Fls 100 mm wide; with secondary buds; tepals 82 × 40 mm, strongly recurved; pedicels 80 mm, green. Lvs 70 × 16 mm, dull mid-green. Stems 0.8 m, rigid, green, felted; infl. short. Early July.

'Bantam' VII(a/b)

Seedling 92-59 × seedling RM 95-73

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals deep purplish pink (inner tepals N66D, outer N66C); margins and midveins strong purplish red (64C), margins of inner tepals closely spotted; throat yellowish white (155D). Outside of outer tepals strong purplish red (64C); margins 64B; midribs strong yellow-green (144C) at base, shading to moderate olive-green (146A) at top. Outside of inner tepals dark purplish pink (186C), shading to deep purplish pink (68A) towards margins and yellowish white (155D) at base beside midribs; midribs light yellow-green (144D) at base and top, flushed deep purplish pink (68A) towards top. Spots and papillae strong purplish red (63A); nectaries strong yellow-green

(144C); pollen vivid reddish orange (34B); stigma very pale green (189D). Fls 235 mm wide, scented, colour intensifying with age; tepals 140 × 75 mm, margins slightly ruffled, tips recurved. Lvs 170 × 55 mm. Stems 1.3 m, green, with up to 3 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Barbarella' VII(a)

Parentage unknown

H: Mak-Leek Inc., pre-2006 **REG:** Mak-Leek Inc., 2006

Inside of tepals strong purplish red (60C); midveins purple-red at base. Outside of inner tepals strong purplish red (71B). Spots and papillae present; nectaries green; pollen orange-brown; stigma grey-green. Tepals of medium length and of medium width to broad, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length and width to broad. Stems of average height, green with darker spots and stripes, with few fls. Early to mid-season.

'Barbina' I(a/b)

Published in *ILR 1982 Twenty-first Supplement* and *ILR&C 2007* as an LA hybrid (Div. VIII (a/b)) with the parentage 'Snowcap' × 'Longicanda'. Reclassified as an Asiatic hybrid (Div. I (a/b)) with the parentage 'Snowcap' × unknown, in July, 2008 (*in litt.*, Ing. B. Mićulka).

'Barbizon' VIII(a/b) LA

Seedling BW 96-03 × seedling AW 94-09

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals yellowish white (155D), margins of inner tepals closely spotted vivid purplish red (61C); throat light yellow-green (145D). Outside of outer tepals yellowish white (155D), with a stripe of light yellow-green (142D) beside midribs and smaller stripes of vivid purplish red (61C); midribs strong yellow-green at base and top (143B and 144C respectively), base flushed vivid purplish red (61C) shading to 61C towards top. Outside of inner tepals yellowish white (155D), margins closely spotted vivid purplish red (61C); midribs light yellow-green (144D). Spots and papillae absent; nectaries strong yellow-green (143A); pollen brownish orange (171B); stigma strong greenish yellow (151C). Fls 180 mm wide; tepals 114 × 56 mm, margins smooth, tips recurved. Lvs 114 × 15 mm. Stems 1–1.2 m, green with darker markings, with c.4 fls.

'Beau Soleil' VIII(a/b) LA

(*longiflorum* 'White Fox' × 'Brandaris') × yellow-flowered Asiatic

H: Testcentrum voor Siergewassen BV, pre-2000

N: Marklily CV, 2000 **REG:** Testcentrum voor Siergewassen BV, 2006

Inside of tepals brilliant yellow (13B), tips and margins of inner tepals spotted strong orange-yellow (17A); outer tepals shading to 17A towards margins; throat light greenish yellow (1C). Outside of tepals vivid yellow (12A); outer tepals shading to vivid

yellow (15A) towards margins; flushed moderate red (181C) at base; midribs vivid yellow (12A), strong yellow-green at base and top (143C and 143B respectively). Inner tepals shading to brilliant yellow (15C) towards margins; midribs vivid yellow (12A), strong yellow-green (143C) at base with a small stripe of moderate red (181B). Spots and papillae absent; nectaries strong yellow-green (143A); pollen dark orange (N163A); stigma brilliant greenish yellow (5A). Fls 215 mm wide, slightly scented; tepals 126 × 60 mm, margins smooth, tips recurved. Lvs 170 × 24 mm. Stems 1.2 m, green, with c.8 fls.

'Belforte' VII(a/b)

Seedling 92-35 × seedling RM 98-19

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals moderate purplish red to dark purplish pink (186B/C), with narrow margin of pale purplish pink (65D); brilliant greenish yellow (3A) beside and above nectaries; midveins flushed moderate reddish orange (35B); throat greenish white (155C). Outside of outer tepals moderate purplish red (186B), basal third and margins of each tepal moderate purplish pink (186D); yellowish white (155D) at base. Outside of inner tepals dark purplish pink (186C) in upper half; basal half moderate purplish pink (186D); yellowish white (155D) at base; midribs strong yellow-green (144B) at base; upper half of midribs of inner tepals, and smaller upper portion of midribs of outer tepals, yellowish white (155D) flushed purplish pink; strong yellow-green at top (144A in outer tepals, 144C in inner tepals). Spots deep purplish red (59B); papillae mostly 59B, but a few brilliant greenish yellow (3A); nectaries strong yellow-green (144B); pollen dark reddish orange (175B/C); stigma light brownish grey (201B). Fls 195 mm wide, becoming paler with age; tepals 120 × 72 mm, margins slightly ruffled, tips recurved. Lvs 215 × 57 mm, dark green. Stems 1.05–1.4 m, light green, with c.3 fls.

'Belladonna' VIII(b-a/b) OT

Seedling PG 95-43 × seedling RH 97-09

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals light greenish yellow: inner 4C, outer 5C; throat pale yellow-green (4D). Outside of outer tepals pale yellow-green (4D); midribs strong yellow-green (144C) at base shading to light yellow-green (154D), and 145C with 145B at top. Outside of inner tepals pale yellow-green (4D) shading to light greenish yellow (4C) towards margins; midribs pale yellow-green (149D) at base shading to brilliant yellow-green (150C). Spots absent; papillae light greenish yellow (4C); nectaries strong yellow-green (144C); pollen dark red (187A); stigma dark pink (182C). Fls 180 mm wide, scented; tepals 100 × 52 mm, margins slightly ruffled, tips recurved. Lvs 190 × 30 mm. Stems 1–1.3 m, green with a little darker marking, with c.3 fls.

'Belle Amour' VII(a-b/b-c)

Parentage unknown

H: World Breeding BV, pre-2005 **N:** World Breeding BV, 2005 **REG:** World Breeding BV, 2007

Inside of inner tepals moderate purplish pink (68C) to light purplish pink (68D), midveins yellow at base, shading to purple-red above. Inside of outer tepals deep purplish pink (N66C/D); outside of inner tepals N66D. Spots and papillae present; nectaries green; pollen orange; stigma grey-green with purple spots.

Tepals of medium length and width, margins ruffled to strongly ruffled, tips recurved. Lvs of medium length, and of medium width to broad. Stems 1.2 m, green, with up to 6 fls. Early to mid-season.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Bellini' VIII(a-b/b) LO

Parentage unknown

H: World Breeding BV, pre-2005 **N:** World Breeding BV, 2005 **REG:** World Breeding BV, 2007

Inside of all tepals and outside of inner tepals pale yellow-green (4D); midveins white at base. Spots absent; papillae inconspicuous; nectaries green; pollen red-brown; stigma purple. Tepals of medium length to long and of narrow to medium width, margins slightly ruffled to ruffled, tips recurved. Lvs long and of medium width to broad. Stems 1.5 m, green with darker spots and stripes, with few fls. Very late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Belo Horizonte' I(a/b)

(Seedling × 'Tango Nocturno') × ('Harmony' seedling × 'Connecticut King')

H: Bischoff Tulleken Lelies BV, pre-2001**N:** Bischoff Tulleken Lelies BV, 2001 **REG:** Mak Breeding BV, 2006

Inside of tepals brilliant yellow (7A); throat light greenish yellow (1C). Outside of outer tepals light greenish yellow (4B); margins brilliant greenish yellow (6A) with dark red (59A) spots; midribs flushed moderate reddish orange (178D), strong yellow-green (143C) at base and 143A at top, sometimes strong greenish yellow (151C). Outside of inner tepals light greenish yellow (4B) shading to brilliant greenish yellow (5B) towards margins, margins with dark red (59A) spots; midribs dark reddish orange (178B) at base shading to strong yellow-green (143A) at top. Spots numerous, large and small, dark red (59A); papillae brilliant yellow (7A); nectaries strong yellow-green (143A); pollen greyish red (178A); stigma moderate purplish red (58A). Fls 195 mm wide; tepals 104 × 56 mm, margins smooth, tips slightly recurved. Lvs 150 × 23 mm. Stems 0.84 m, green with darker markings, with c.5 fls.

'Beluga' VII(a/b)

Seedling RM 96-26 × seedling RM 99-18

H: Vletter en Den Haan Beheer BV, pre-2004**N:** Vletter en Den Haan Beheer BV, 2004**REG:** Vletter en Den Haan Beheer BV, 2007

Inside of tepals strong purplish red (64B), narrow margins and throat yellowish white (155D); midveins

moderate purplish red (64A). Outside of tepals moderate purplish red (outer tepals 64A, inner 70A), narrow margins and base beside midribs yellowish white (155D); all midribs strong yellow-green (144C) at base, midribs of outer tepals shading to 144B towards top, those of inner tepals shading to light yellow-green (150D) and 145C at top. Spots deep red (60A); papillae yellowish white (155D) with deep red (60A); nectaries light yellow-green (144D); pollen moderate reddish orange (171A); stigma light grey (188D). Fls 230 mm wide, slightly scented; tepals 153 × 83 mm, margins ruffled, tips slightly recurved. Lvs 195 × 40 mm. Stems 1.1–1.2 m, light green, with up to 4 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Biaritz' VIII(a-b/b) OT

Seedling PG 95-43 × seedling RH 97-09

H: Vletter en Den Haan Beheer BV, pre-2004**N:** Vletter en Den Haan Beheer BV, 2004**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals brilliant yellow (7A), pale greenish yellow (9D) beside midveins towards tips.

Inside of outer tepals brilliant yellow (9C), midveins and towards margins shading to vivid yellow (9B), a small patch of yellowish white (155D) at base beside nectaries; throat pale yellow-green (4D). Outside of outer tepals pale yellow-green (4D) shading to light greenish yellow (5C) towards margins, yellowish white (155D) at base. Outside of inner tepals light greenish yellow (8C) shading to 8B towards margins, pale yellow-green (4D) beside midribs; all midribs strong yellow-green (144C) at base shading to light yellow-green (154D); midribs of outer tepals strong yellow-green (144C) at top with 144A; midribs of inner tepals brilliant yellow-green (150C) at top with dark greenish yellow (152D). Spots absent; papillae vivid yellow (9A); nectaries strong yellow-green (144B); pollen greyish brown (166A); stigma greyish purple (N77A). Fls 245 mm wide, scented; tepals 145 × 75 mm, margins ruffled, tips recurved. Lvs 210 × 30 mm. Stems 1.2–1.35 m, with darker markings at leaf nodes, with c.4 fls.

'Biscarrosse' VII(a/b-c)

Seedling 92-82 × seedling 93-61

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of tepals moderate purplish pink (62B); above and beside midveins, margins towards tips, and throat, all yellowish white (155D). Outside of outer tepals strong purplish pink (62A), up to half of each tepal shading to pale purplish pink (65C) towards margins; narrow margins of yellowish white (155D) towards tips and beside midribs; midribs strong yellow-green (144C) at base, flushed strong purplish pink (62A), shading to strong greenish yellow (151B), moderate yellow-green (138B) at top. Outside of inner tepals light purplish pink (62C), narrow margins of yellowish white (155D) towards tips and beside midribs; midribs strong yellow-green (144C) at base shading to 144B flushed light purplish pink

(62C). Spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144B); pollen moderate reddish orange (171A); stigma light violet (92C). Fls 220 mm wide; tepals 125 × 70 mm, margins slightly ruffled, tips recurved. Lvs 165 × 36 mm. Stems 1–1.4 m, light green, with c.3 fls.

'Black Label' I(a/b)

(Seedling × Sundrop Group) × ('Red Carpet' × 'Pineapple')

H: Bischoff Tulleken Lelies BV, pre-2001

N: Bischoff Tulleken Lelies BV, 2001

REG: Mak Breeding BV, 2006

Inside of tepals strong red (46A), margins of inner tepals deep red (53A); throat strong orange (169B). Outside of outer tepals moderate red (180A) shading to 47A towards margins. Outside of inner tepals moderate red (180B) shading to strong red (53B) towards margins. Spots greyish purple (N77A); papillae strong orange (169B); nectaries strong yellow-green (143B); pollen dark reddish orange (178B); stigma dark red (59A). Fls 120–155 mm wide; tepals 80 × 45 mm, margins slightly ruffled, tips recurved. Lvs 105 × 15 mm. Stems 0.4–0.6 m, green, with c.9 fls.

'Black Out' I(b/b)

This lily was registered as 'Black-out' in 2000. In 2001 Dutch Plant Breeders' Rights were granted under the cultivar epithet 'Black Out', which becomes the spelling accepted by the ICRA (ICNCP, 2004: Art. 28.2).

BOGOTA VIII(a/b) LA

Trade designation of 'Zanlagota'.

'Bolsena' VII(a/c)

Seedling 93-07 × seedling RM 99-59

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals deep purplish pink (N57D); midveins vivid purplish red (N57B) for up to half their length; throat yellowish white (155D). Outside of outer tepals dark purplish pink (186C), yellowish white (155D) at base beside midribs; midribs strong yellow-green (143C) at base and 143B at top, shading to pale yellow-green (149D) between. Outside of inner tepals light purple (75B), yellowish white (155D) at base beside midribs; midribs moderate yellow-green (143D) at base, shading to light yellow-green (150D), and 145C at top. Spots and papillae strong purplish red (63A); nectaries strong yellow-green (144C); pollen moderate reddish brown (175A); stigma light greenish grey (190D). Fls 225 mm wide, scented; tepals 130 × 60 mm, margins slightly ruffled, tips recurved. Lvs 230 × 50 mm. Stems 1.2 m, green with darker markings, with up to 6 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Bonita' VII(a-b/b)

A temporary name for 'Una Bonita' prior to registration.

'Bonsoir' VIII(a-b/b) LA

Seedling BJ 96-21 × OR 91-15

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals strong orange (25A), midveins and towards margins shading to vivid orange (28B); throat strong orange-yellow (24B). Outside of outer tepals brilliant orange-yellow (23B) shading to strong orange (25A) towards margins; midribs dark greenish yellow (152D) at base. Outside of inner tepals strong orange (25B) shading to 25A towards margins; midribs dark greenish yellow (152C) at base. Spots and papillae absent; nectaries moderate olive-green (137A); pollen strong reddish orange (169A); stigma strong orange-yellow (24B). Fls 190 mm wide, slightly scented, becoming paler with age; tepals 115 × 46 mm, margins smooth, tips recurved. Lvs 136 × 20 mm. Stems 1.2–1.4 m, green with darker markings, with c.6 fls.

'Briancon' VII(a/b)

Seedling 92-50 × seedling RM 96-26

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of inner tepals deep purplish pink (N66D); base and narrow margins yellowish white (155D); midveins deep pink (50B); throat yellowish white (155D). Inside of outer tepals deep purplish pink (N66D); margins, from base to up to half the length of each tepal, deep purplish pink (N66C), shading to yellowish white (155D); midveins strong red (50A). Outside of outer tepals moderate purplish red (186B), yellowish white (155D) beside midribs; midribs strong yellow-green (144C) at base, and 144A at top, shading to light yellow-green (144D) between. Outside of inner tepals moderate purplish pink (186D), yellowish white (155D) towards base beside midribs; midribs strong yellow-green (N144D) at base, shading to light yellow-green (144D). Outside of all tepals with narrow margins of yellowish white (155D) in upper half. Spots deep purplish red (61A); papillae yellowish white (155D); nectaries strong yellow-green (144C) edged brilliant greenish yellow (1B); pollen strong brown (172A); stigma light greenish grey (190D). Fls 220 mm wide, scented, colour becoming darker with age; tepals 135 × 65 mm, margins ruffled, tips recurved. Lvs 170 × 58 mm. Stems 1.2–1.4 m, green, with 1 or 2 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Brňanka' VI(b-c/b-d)

('Lady Alice' × (unknown × 'Toni')) × unknown

H: B. Mičulka, pre-2007 **N:** B. Mičulka, pre-2007

I: B. Mičulka, 2007 **REG:** B. Mičulka, 2007

Inside of tepals creamy apricot, central portion and throat orange-apricot; tips whitish. Midribs yellow-green. Spots absent; papillae small, raised, resembling rusty brown dashes over two-thirds of each tepal; nectaries dark green; pollen brown; stigma green. Fls 125 mm wide; tepals 95 × 38 mm, stiff, margins slightly ruffled, tips recurved; pedicels 60 mm, sparsely spotted brown; with secondary buds. Lvs 120 × 19 mm, pale green. Stems 1.15 m, with short infl. Mid-July.

'Broken Heart' VII(b-a/c)

Seedling 9-13-SP × 'Star Gazer'

N: Rooyakkers Lelie BV, 2007 **REG:** Rooyakkers Lelie BV, 2007

Inside of inner tepals deep purplish pink (64D) on a paler ground; inside of outer tepals strong purplish red (64C), one margin of each outer tepal tinged greenish white (155C); all midveins with a strong red (53B) stripe; throat greenish white (155C). Outside of outer tepals strong purplish red (64C) on a paler ground; midribs moderate reddish brown (166B) at base, shading to brilliant yellow-green (154B), and strong yellow-green (144A) at top. Outside of inner tepals greenish white (155C) flushed strong purplish red (64C). Spots deep red (60A); papillae absent; nectaries strong yellow-green (143C); pollen and stigma absent. Fls 160–200 mm wide, double; tepals 90 × 30 mm, margins slightly ruffled, tips recurved. Lvs 185 × 42 mm, dark green. Stems 0.8–1.2 m, pale green with some darker markings, with up to 5 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Bronte' I(a/b)

Seedling RS 91-23 × seedling AM 95-07

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of tepals strong purplish pink (62A) shading to deep purplish pink (64D) towards margins and tips; outer tepals with a larger patch of deep purplish pink (64D) at tips; throat pale purplish pink (62D). Outside of outer tepals moderate purplish pink (62B) shading to deep purplish pink (64D) towards margins; yellowish white (155D) at base; midribs strong yellow-green (143C) at base and top. Outside of inner tepals strong purplish pink (63C) shading to strong purplish red (63B) towards margins; a small patch of yellowish white (155D) at base; midribs strong yellow-green (143C) at base. Spots strong purplish red (60B); papillae pale purplish pink (62D); nectaries strong yellow-green (143C); pollen brownish orange (N167B); stigma light yellow-green (142D) with a small flush of strong purplish red (59D) below. Fls 170 mm wide, becoming dark purplish pink (186C) with age; tepals 100 × 45 mm, margins smooth, tips recurved. Lvs 124 × 15 mm. Stems 1–1.1 m, dark with green markings, with c.4 fls.

'Bruno' I(a/b)

'Centurion' × 'Bold Knight' seedling

H: T.D. Willoughby, 1991 **G:** T.D. Willoughby,1993 **N:** T.D. Willoughby, 2000 **REG:** T.D. and

S.K. Willoughby, 2006

Inside of tepals strong red (46A) shading to 41B towards margins; tips vivid red (44A); throat strong red (41B). Outside of tepals vivid reddish orange (44C). Spots 1 mm across, oblong, black, raised, 45–50 evenly distributed in lower half of each tepal; papillae slightly frosted; nectaries strong red (41B); pollen moderate reddish orange (171A); stigma dark red-purple. Fls 135 mm wide; tepals 75 × 42 mm, margins smooth, tips straight. Lvs 70 × 25 mm, moderate olive-green (137A). Stems 1.4 m, moderate olive-green (137A), with up to 12 fls in a racemose

infl. Mid-July (USDA Zone 3, Alberta, Canada).

Named after a late friend of the raiser, Herbert

"Bruno" Sunley, who bred 'Bold Knight'.

'Brushmania' I(a/c)

('Echo' × ('Mirabel' × unknown)) × ('Koral' × ('Mirabel' × unknown))

H: B. Mićulka, pre-2006 **N:** B. Mićulka, pre-2006**I:** B. Mićulka, 2006 **REG:** B. Mićulka, 2006

Inside of tepals yellow; throat paler yellow. Outside of tepals paler yellow; midribs greenish. Brushmarks small, faint, mahogany brown; spots sparse, dark brown, fairly large, over a quarter of each tepal; papillae absent; nectaries green, feathered white; pollen dark brown; stigma dark red-brown. Fls 110 mm wide; tepals 65 × 30 mm, tips slightly recurved; pedicels 75 mm, spotted brown. Lvs 80 × 10 mm, dull mid-green. Stems 0.75 m, green, stiff, infl. short. Late June.

'Burlesca' VII(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of tepals dark purplish pink (186C), with a patch of yellowish white (155D) above and beside nectaries, and at margins towards tips of outer tepals; throat yellowish white (155D). Outside of outer tepals pale purplish pink (65C), yellowish white (155D) at base; midribs strong yellow-green (143B) at base shading to light yellow-green (145D), and strong yellow-green (144B) at top. Outside of inner tepals moderate purplish pink (186D) shading to dark purplish pink (186C) towards margins; up to half the length of each inner tepal yellowish white (155D) beside midribs; midribs strong yellow-green (143B) at base shading to 144C and pale greenish yellow (2D), and strong yellow-green (144C) at top. Spots absent; papillae yellowish white (155D); nectaries strong yellow-green (143C); pollen moderate reddish orange to brownish orange (171A/B); stigma greenish grey (N189C). Fls 240 mm wide, slightly scented; tepals 150 × 83 mm, margins ruffled, tips recurved. Lvs 185 × 55 mm. Stems 1.12–1.4 m, green with darker markings at base, with c.3 fls.

'Bursa' I(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2007

Inside of tepals vivid reddish orange (N30C); midveins of inner tepals, above nectaries of outer tepals, and all margins vivid reddish orange (N30B); throat strong orange (30D). Outside of outer tepals strong orange-yellow (24B) flushed light yellow-green (144D), shading to vivid reddish orange (N30B) towards margins. Outside of inner tepals vivid reddish orange (30B), strong yellowish pink (32C) beside midribs; midribs light yellow-green (144D). Spots and papillae absent; nectaries dark green (136A); pollen greyish brown (166A); stigma vivid reddish orange (33B). Fls 185 mm wide; tepals 110 × 55 mm, margins smooth, tips slightly recurved. Lvs

153 × 18 mm. Stems 1.1–1.4 m, green, with up to 8 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Bussetto' VIII(a/b) LA

Seedling BG 95-07 × seedling AG 95-14
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006
Inside of tepals brilliant greenish yellow (6A), margins brilliant yellow (7A); throat light yellow-green (2C). Outside of outer tepals light greenish yellow (7D) shading to vivid yellow (9A) towards margins; midribs light yellow-green (145B) at base. Outside of inner tepals vivid yellow (9B) shading to 9A towards margins, pale greenish yellow (2D) at base. Spots dark red (59A); papillae absent; nectaries strong yellow-green (143B); pollen brownish orange (166C); stigma dark red (59A). Fls 195 mm wide; tepals 125 × 67 mm, margins smooth, tips slightly recurved. Lvs 164 × 21 mm. Stems 1–1.35 m, green with darker markings, with c. 5 fls.

'Cadisha' VIII(a/b) LA

'Akina' × seedling LA 93007-4T
H: De Jong Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2007
Inside of tepals strong orange (25A), shading to N25A towards margins; midveins of inner tepals strong orange (N25B); throat light orange-yellow (24C). Outside of outer tepals strong orange (24A), shading to 25A towards margins; strong yellow-green (143C) beside midribs and dark red (183B) at base; midribs strong yellow-green (143C). Outside of inner tepals strong orange (25B), shading to 25A towards margins; midribs dark red (183B) tinged strong yellow-green (143C). Spots few, dark red (59A); papillae absent; nectaries light olive (152A); pollen dark red (183B); stigma strong reddish orange (34C). Fls 180 mm wide, slightly scented; tepals 133 × 58 mm, margins smooth, tips slightly recurved. Lvs 170 × 30 mm. Stems 1.2–1.25 m, dark, with up to 3 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Caitlin's Sunrise' I(a/b)

'Lily Simonet' × 'Golden Summer'
H: T.D. Willoughby, 1991 **G:** T.D. Willoughby, 1993 **N:** T.D. Willoughby, 1996 **I:** Honeywood Lilies, 2007 **REG:** T.D. and S.K. Willoughby, 2006
Inside of tepals vivid reddish orange (42A), with broad vivid yellow (15A) central flare with brush-marked edges; throat vivid reddish orange (42A). Outside of tepals strong orange (25B); margins strong red (46A). Scattered, bold spots on or near flare; papillae absent; nectaries strong reddish orange (42B); pollen deep orange-yellow (163A); stigma vivid red (45A). Fls 130 mm wide; tepals 75 × 35 mm, margins smooth or slightly ruffled, tips slightly recurved. Lvs 100 × 15 mm, moderate yellowish green (138A). Stems 1 m, moderate yellowish green (138A), with up to 15 fls in a racemose inf. Mid-July (USDA Zone 3, Alberta, Canada). Named after the hybridizer's eldest daughter.

'Calvados' VII(a)

Parentage unknown
H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005
Inside of tepals strong to vivid purplish red (61B/C), lighter towards margins and tips; base of midveins deep red (53A). Outside of inner tepals deep purplish pink (N66D). Spots numerous; papillae present; nectaries green changing to yellow; pollen orange; stigma dark purple. Tepals of medium length to long and of medium width, margins slightly ruffled, tips recurved. Lvs of medium length, broad. Stems of average height, green with darker stripes and spots, with few fls. Mid-season to late.

'Candle Glow' VIII(a-b/-) OT

Parentage unknown
H: World Breeding BV, pre-2005 **REG:** World Breeding BV, 2005
Inside of inner tepals light greenish yellow (8B/C), pale yellow towards tips (8D). Inside of outer tepals light greenish yellow (8C); base of midveins brilliant yellow (8A). Outside of inner tepals light greenish yellow (8C). Spots absent, papillae present, nectaries green; pollen orange-brown; stigma purple. Tepals of medium length to long and of medium width, margins slightly ruffled, tips recurved. Lvs of medium length to long and of medium width to broad. Stems of average height to tall, green, with few to average no. of fls. Mid-season to late.

'Cantarino' VII(a-b/c-b)

Parentage unknown
H: Testcentrum voor Siergewassen BV, pre-2004
N: Marklily CV, 2004 **REG:** Testcentrum voor Siergewassen BV, 2007
Tepals greenish white (155C); midveins pale yellow (8D) at base. Spots absent; papillae present; nectaries green; pollen orange-brown; stigma purple. Tepals of medium length and width, margins ruffled to strongly ruffled, tips slightly recurved to recurved. Lvs of medium length and broad to very broad. Stems of average height to tall, green with faint darker speckles and stripes, with few to average no. of fls. Mid-season to late.
Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Caravelle' VII(a/b)

Seedling RM 95-50 × seedling RM 99-59
H: Vletter en Den Haan Beheer BV, pre-2004
N: Vletter en Den Haan Beheer BV, 2004
REG: Vletter en Den Haan Beheer BV, 2007
Inside of tepals deep purplish pink (70C), shading to light purple (84C) towards margins, tinged pale greenish yellow (2D) above nectaries; midveins strong purplish pink (68B) towards top; throat yellowish white (155D). Outside of outer tepals dark purplish pink (186C), shading to very pale purple (69D) towards margins and yellowish white (155D) at base beside midribs. Outside of inner tepals moderate purplish pink (186D) and yellowish white (155D) beside midribs for up to half the length of each tepal. All midribs strong yellow-green (144B) at base and

top, shading to pale yellow-green (149D) between. Spots strong purplish red (58B); papillae yellowish white (155D); nectaries strong yellow-green (144C); stigma pale green (189C). Fls 210 mm wide, scented; tepals 115 × 70 mm, margins ruffled, tips slightly recurved. Lvs 190 × 55 mm. Stems 1–1.1 m, green with darker markings at base, with up to 4 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Cardiff’ I(a)

Parentage unknown

H: World Breeding BV, pre-2001 **N:** World Breeding BV, 2001 **REG:** World Breeding BV, 2005
Inside of inner tepals strong orange yellow to moderate yellow (163B/C), paler towards base; midveins vivid yellow (13A) at base. Outside of inner tepals strong orange-yellow (N163C/D). Spots few, papillae present, nectaries green changing to red; pollen red-brown; stigma purple. Young fls more reddish, becoming more yellow and paler with age; tepals very short to short and of narrow width, margins slightly ruffled, tips slightly recurved. Lvs of medium length and of narrow to medium width. Stems tall, green with darker stripes and spots, with few to average no. of fls. Early to mid-season.

‘Carre’ VII(a/b-c)

Seedling 88-99 × seedling RM 94-23

H: Vletter en Den Haan Beheer BV, pre-1997

N: Vletter en Den Haan Beheer BV, 1997

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals and margins moderate purplish red (64A and 70A respectively); midveins deep red to strong purplish red (60A/B); throat greenish white (155C). Outside of outer tepals moderate purplish red (64A); margins strong purplish red (60B); midribs strong yellow-green (144B) from base to up to half their length, shading through pale purplish pink (62D) to strong purplish red (64B) towards top; top strong yellow-green (144A). Outside of inner tepals strong purplish red (64B), shading through moderate purplish red (186B) towards strong purple (77B) at margins; midribs moderate purplish pink (70C), strong yellow-green (144B) at base. Spots deep red (53A); papillae absent; nectaries strong yellow-green (144B); pollen moderate reddish orange (N172A); stigma purplish grey (N187C). Fls 210 mm wide; tepals 130 × 64 mm, margins slightly ruffled, tips recurved. Lvs 175 × 54 mm, mid-green. Stems 1.25 m, dark, with *c.* 4 fls.

‘Caruso’ VII(b-c/c)

Seedling 93-25 × seedling RM 96-26

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals deep purplish pink (N66D) shading to deep purplish pink (70C) towards margins; midveins deep purplish pink (67C) at immediate top; throat yellowish white (155D). Inside of outer tepals very light purple (75C); deep purplish pink (N66C) towards tips; narrow margins and above and beside nectaries, yellowish white (155D). Outside of outer

tepals moderate purplish pink (186D); strong purplish red (63B) at tips and yellowish white (155D) at base; midribs strong yellow-green (144C) at base, shading to yellowish white (155D), and moderate to strong yellow-green (138B/143A) at top. Outside of inner tepals moderate purplish pink (186D); midribs strong yellow-green (143C) at base shading to yellowish white (155D) for up to half their length; upper half deep purplish pink (N66D), and light yellow-green (138D) at top. Spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen dark reddish orange (172B); stigma pale yellow-green (193D). Fls 260 mm wide, scented; tepals 135 × 67 mm, margins ruffled, tips recurved. Lvs 165 × 38 mm. Stems 1–1.3 m, green, with up to 6 fls.

‘Casanova Revue’ I(a/b)

‘Czardas’ × seedling St. 18/88

H: G. Steinbrück, 1991 **G:** G. Steinbrück, 1996

N: G. Steinbrück, 2006 **I:** G. Steinbrück, 2006

REG: G. Steinbrück, 2007

Inside of tepals strong red (53B), with yellow blotches towards tips. Outside of outer tepals green-yellow. Outside of inner tepals strong red (53B). Spots small, faint, brown, over three-quarters of each tepal; papillae absent; nectaries white shading to pale violet; pollen brownish orange (165B); stigma yellowish white (155D). Fls 135 mm wide; tepals 70 × 35 mm, margins smooth, tips slightly recurved. Lvs 100 × 20 mm, scattered. Stems 1.2 m, dark green, with up to 10 fls. Mid-July. Named after a Revue staged at the Friedrichstadtpalast, Berlin.

‘Cervo’ VII(a/b)

Seedling 90-19 × seedling RF 94-01

H: Vletter en Den Haan Beheer BV, pre-2007

N: Vletter en Den Haan Beheer BV, 2007

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals strong purplish red (60B), shading to moderate purplish red (64A) towards margins; midveins of outer tepals tinged yellowish white (155D) towards top; throat yellowish white (155D). Outside of tepals strong purplish red: outer tepals 60B, shading to 60C towards margins; inner tepals 64B, shading to 64C towards margins; yellowish white (155D) at base beside midribs; all midribs strong yellow-green (144C) at base shading to light olive (152A) towards top; midribs of inner tepals strong purplish red (64B) between. Spots and papillae dark red (59A); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma light yellow-green (150D). Fls 200 mm wide, slightly scented; tepals 115 × 70 mm, margins smooth, tips slightly recurved. Lvs 200 × 56 mm. Stems 1.1–1.2 m, dark with green markings, with up to 4 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Cevennes’ VIII(a/b) LA

Seedling BG 95-53 × seedling G 90-05

H: Vletter en Den Haan Beheer BV, pre-1994

N: Vletter en Den Haan Beheer BV, 1994

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals vivid yellow (13A) shading to 17B

towards margins; throat light greenish yellow (1C). Outside of tepals vivid yellow (outer 12A, inner 13A), all shading to vivid yellow (15A) towards margins; midribs of outer tepals light yellow-green (144D), strong yellow-green (144A) at top; midribs of inner tepals strong yellow-green (144C) at base shading to vivid yellow (13A). Spots and papillae absent; nectaries strong yellow-green (143A); pollen strong brown (172A); stigma brilliant greenish yellow (1B). Fls 195 mm wide; tepals 117 × 53 mm, margins smooth, tips slightly recurved. Lvs 177 × 25 mm. Stems 1.2–1.4 m, green with darker markings, with *c.* 7 fls.

‘Chiara’ VIII(b/a-b) LO

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2003

N: Marklily CV, 2003 **REG:** Testcentrum voor Siergewassen BV, 2007

Inside of tepals light purplish pink (63D) on a white ground; midveins green at base. Outside of inner tepals pale purplish pink (65D). Spots and papillae absent; nectaries green; pollen pale brown; stigma pale grey. Tepals long and broad, margins (very) slightly ruffled, tips slightly recurved to recurved. Lvs long and of medium width to broad. Stems tall, green, with few fls. Late to very late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Christine’ VI(c/a)

Unknown Trumpets

G: D. Fielding, 2000 **N:** D. Fielding, 2000

REG: D. Fielding, 2007

Inside of tepals light purplish pink (62C); margins mostly white; throat brilliant greenish yellow (2B). Outside of tepals very light purple (75C). Spots and papillae absent; nectaries strong yellow-green (142A); pollen deep orange-yellow (163A). Fls 135 mm wide, slightly scented; tepals 120 × 50 mm, margins slightly ruffled, tips slightly recurved. Lvs 145 × 13 mm, scattered, dark green, glossy. Stems 0.9 m, dark green, robust, with 25 or more fls in a pyramidal infl.

January (Australia).

Named after the grower’s wife.

‘Cilesta’ VIII(a/b) LA

Seedling BG 96-15 × seedling XA 01-15

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals strong orange (25A), shading to N25A towards margins; midveins N25A too; throat pale orange-yellow (23D). Outside of outer tepals strong orange-yellow (24B), shading to strong orange (N25A) towards margins; midribs strong yellow-green (143C) at base and 144A at top), shading to light yellow-green (144D) between. Outside of inner tepals strong orange (25B), shading to vivid yellowish pink (28A) towards margins and light orange-yellow (16C) in basal third beside midribs; midribs strong yellow-green (144C) from base to up to half the length of each tepal, shading to strong orange (25B). Spots and papillae absent; nectaries strong yellow-green (143A); pollen greyish reddish brown (200B); stigma dark red

(59A). Fls 228 mm wide; tepals 120 × 55 mm, margins smooth, tips slightly recurved. Lvs 155 × 26 mm.

Stems 1.4–1.5 m, green, with up to 5 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Cocktail Twins’ I(a/b)

‘Fata Morgana’ × ‘Fata Morgana’ F₂

H: Laan Flora Facilities BV, pre-2005 **N:** Laan Flora Facilities BV, 2005 **REG:** Laan Flora Facilities BV, 2006

Inside of tepals strong red (46A) above throat shading to moderate red (179B); a small flush of vivid yellow (15A) at tips; throat yellowish white (155D). Outside of outer tepals brownish orange (172C) shading to vivid reddish orange (34B) towards margins, with a broad stripe of light yellow-green (145C) towards deep purplish red (59B) tips. Outside of inner tepals moderate red (179B) shading to 179A towards margins; midribs moderate yellowish pink (N170D). Spots greyish purple (N77A), densely distributed over lower portion of inside of inner tepals; papillae absent; nectaries yellowish white (155D); stamens absent; stigma dark red (59A). Buds slightly hairy. Fls 185 mm wide, double, pedicels hairy; tepals 95 × 37 mm, margins smooth, tips slightly recurved. Lvs 152 × 24 mm. Stems 0.9–1.3 m, dark with green spots, with *c.* 5 fls.

‘Colgate’ VIII(a/-) LA

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Inside of tepals yellowish white (155D), light yellow on opening. Outside of inner tepals yellowish white (155B). Spots few; papillae present; nectaries green; pollen orange-brown; stigma yellow-green. Tepals of short to medium length and of medium width, margins slightly ruffled, tips slightly ruffled to ruffled. Lvs long and of narrow to medium width. Stems tall, green, with few to average no. of fls. Early to mid-season.

‘Columbia’ VII(a/b)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2006

N: Marklily CV, 2006 **REG:** Testcentrum voor Siergewassen BV, 2007

Inside of tepals greenish white (155C); brilliant yellow (8A) in basal half and along midveins. Outside of inner tepals greenish white (155C). Spots yellow; papillae present; nectaries pale green; pollen reddish brown; stigma purple. Tepals of medium length to long and of medium width to broad, margins ruffled, tips slightly recurved to recurved. Lvs of medium length to long and of medium width to broad. Stems of average height to tall, green with darker speckles and stripes, with few to average no. of fls. Mid-season to late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Compassion' VIII(b-c/b) LO

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2003**N:** Marklily CV, 2003 **REG:** Testcentrum voor Siergewassen BV, 2007

Inside of tepals deep purplish pink (N66C/D); midveins pale purple-red at base; throat green. Outside of inner tepals deep purplish pink (N66D). Spots and papillae absent; pollen brown; stigma pale grey-green. Tepals long and of medium width to broad, margins (very) slightly ruffled, tips recurved to strongly recurved. Lvs long to very long and of medium width to broad. Stems up to 1.9 m, green with some darker spots and stripes, with up to 6 fls. Very late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Conception' I(a/b)

Seedling AF 99-11 × seedling AF 99-07

H: Vletter en Den Haan Beheer BV, pre-2004**N:** Vletter en Den Haan Beheer BV, 2004**REG:** Vletter en Den Haan Beheer BV, 2007

Inside of tepals dark red (187C), shading to 187B towards margins; throat strong red (53C). Outside of tepals moderate red (185B), shading to dark red (59A) towards margins; midribs deep red (185A). Spots and papillae dark red (187A); nectaries strong yellow-green (143B); pollen dark reddish orange (175C); stigma dark red (187A). Fls 180 mm wide; tepals 100 × 50 mm, margins smooth, tips recurved. Lvs 160 × 15 mm. Stems 1.2–1.4 m, dark with a few green markings, with up to 9 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).**'Concordia' VII(a/b-c)**

(Seedling × 'Star Gazer') × red-flowered seedling

H: Bischoff Tulleken Lelies BV, pre-2003**N:** Bischoff Tulleken Lelies BV, 2003**REG:** Mak Breeding BV, 2006

Inside of tepals deep red (60A/53A) shading through strong purplish red (60C), and strong reddish purple (70B) in inner tepals, towards yellowish white (155D) margins; margins of outer tepals narrower; brilliant greenish yellow (6C) along edge of nectaries; throat yellowish white (155D). Outside of outer tepals moderate purplish red (186B) shading to yellowish white (155D) towards margins and base; midribs moderate purplish red (186B) at base shading to pale yellowish pink (159D), and light yellow-green (144D) with strong yellow-green (144A) at top. Outside of inner tepals dark purplish pink (186C), with yellowish white (155D) veins and base; moderate purplish pink (186B) beside midribs; midribs light yellow-green (145C) at base with moderate purplish pink (186B) stripe, shading to yellowish white (155D), and strong yellow-green (144B) at top. Spots and papillae dark red to deep purplish red (59A/B); nectaries strong yellow-green (144C); pollen moderate reddish orange (171A); stigma dark purplish red (N79B). Fls 230 mm wide, slightly scented; tepals 134 × 63 mm, margins smooth, tips recurved. Lvs 170 × 34 mm. Stems 0.85–0.95 m, green with darker markings at leaf nodes, with c.6 fls.

'Conquistador' I(a/-)

Syn. of 'Conquestador'.

This lily was registered as 'Conquistador' in 1997 and a description under this spelling was published in *ILR 1982 Sixteenth Supplement*. However, in 1993, Dutch Plant Breeders' Rights had been granted to this lily under the cultivar epithet 'Conquestador', which becomes the spelling accepted by the ICRA (ICNCP, 2004: Principles 3, 5 and Art. 24.4). In 2000, a US Plant Patent was granted to 'Conquistador', which remains the accepted spelling in countries in which 'Conquistador' has legal protection (ICNCP, 2004: Art. 28.2).

'Cornas' VII(a-b/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-1996**N:** Vletter en Den Haan Beheer BV, 1996**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of tepals and throat yellowish white (155D); midveins of inner tepals pale yellowish green (155A), those of outer tepals pale yellow-green (4D). Outside of tepals yellowish white (155D); pale yellow-green (4D) beside midribs of outer tepals; midribs of outer tepals strong yellow-green at base and top (143A and 143B respectively), shading to light greenish yellow (4C) between; midribs of inner tepals strong yellow-green (143C) at base, shading to light greenish yellow (4B) above. Spots and papillae yellowish white (155D); nectaries strong yellow-green (145A); pollen strong to moderate reddish orange (169A/171A); stigma very pale green (130D). Fls 220 mm wide, slightly scented; tepals 115 × 65 mm, margins ruffled, tips recurved. Lvs 215 × 53 mm. Stems 1.2 m, green, with c.7 fls.

CYRANO VIII(a/b) LO

Trade designation of 'Zanlorano'.

'Decennia' VII(a-b/b)

Seedling RW 98-21 × seedling RW 98-06

H: Vletter en Den Haan Beheer BV, pre-1999**N:** Vletter en Den Haan Beheer BV, 1999**REG:** Vletter en Den Haan Beheer BV, 2006

Tepals and throat yellowish white (155D); midveins flushed pale yellow-green (4D); midribs strong yellow-green (143C) at base shading to brilliant greenish yellow (151D); midribs of outer tepals strong yellow-green (143B) at top. Spots absent; papillae present; nectaries light yellow-green (144D); pollen strong to dark reddish orange (169A/173A); stigma bluish white (112D). Fls 195 mm wide, scented; tepals 125 × 80 mm, margins slightly ruffled, tips slightly recurved. Lvs 195 × 55 mm. Stems 1 m, green with darker markings at leaf nodes, with c.5 fls.

'Deshima' (Bischoff Tulleken) VII(a-b/c-b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2007**N:** Bischoff Tulleken Lelies BV, pre-2007

Tepals yellowish white (155D). Spots absent; papillae present; nectaries pale green; pollen pale brown; stigma grey-green spotted purple. Tepals of medium length to long and of medium width, margins (very) strongly ruffled, tips recurved. Lvs of medium length

to long and broad to very broad. Stems of average height to tall, green, with few fls. Mid-season. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007). An application for registration was turned down in 2007, as this cultivar epithet was registered by Van der Salm in 1988 for another lily (ICNCP, 2004: Art. 27.1). However, as Dutch Plant Breeders' Rights were granted to this lily in 2002, the epithet remains the accepted name (ICNCP, 2004: Art. 28.2).

'Desiderio' VIII(a-b/b) LA

(*longiflorum* 'White Fox' × 'Brandaris') × orange-flowered Asiatic mixture

H: Testcentrum voor Siergewassen BV, pre-2004
N: Marklily CV, 2004 **REG:** Testcentrum voor Siergewassen BV, 2006

Inside of tepals strong orange (25B); margins and midveins 25A; throat brilliant orange (25C). Outside of outer tepals brilliant orange (25C); margins strong orange (25B); base and top of midribs strong yellow-green (143C), with some 143A at top. Outside of inner tepals strong orange (25B); midribs light olive (152B) at base. Spots dark red (59A); papillae absent; nectaries deep yellowish green (141A); pollen strong reddish orange (169A); stigma strong orange (30D). Fls 200 mm wide; tepals 115 × 56 mm, margins slightly ruffled, tips recurved. Lvs 166 × 16 mm. Stems 1.2 m, green, with c.8 fls.

'Dexter' VII(a-b/-)

Parentage unknown

H: Bottema, pre-2007 **REG:** Bottema Bloembollen BV, 2007

Inside of tepals deep purplish pink (68A) to strong purplish pink (73B); midveins orange-yellow at base, shading to purple-red. Outside of inner tepals strong purplish pink (73B). Spots and papillae present; nectaries green; pollen orange-brown; stigma purple. Tepals of medium length and of narrow to medium width, margins ruffled to strongly ruffled, tips slightly recurved to recurved. Lvs of medium length, broad. Stems of short to average height, green, with few to average no. of fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Diabolo' VIII(a/b) LA

Seedling BM 96-24 × seedling RS 91-16

H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006
Inside of tepals moderate purplish pink (65A); margins and midveins strong purplish pink (67D); above and beside throat pale purplish pink (65D); throat yellowish white (155D) with a narrow, strong yellow-green (144C) edge. Outside of tepals dark purplish pink (186C); margins of outer tepals strong purplish red (63B); margins of inner tepals strong purplish pink (63C); midribs of outer tepals strong yellow-green (144C) at base. Spots deep purplish red (61A); papillae absent; nectaries moderate olive-green (137A); pollen dark orange (N163A); stigma light yellow-green (145C). Fls 170 mm wide; tepals 103 ×

54 mm, margins smooth, tips recurved. Lvs 148 × 18 mm. Stems 0.9–1.3 m, dark, with c.5 fls.

'Diamond Willough' I(a/b)

'Lily Simonet' × pink-flowered seedling bred by Fred Fellner

H: T.D. Willoughby, 1991 **G:** T.D. Willoughby, 1993
N: T.D. and S.K. Willoughby, 2005 **I:** L. Westfall, 2007 **REG:** T.D. and S.K. Willoughby, 2006

Inside of tepals strong red (46A), tips 50A; throat strong red (51A) with a brilliant orange-yellow (23B) blaze in nectary furrow. Outside of tepals deep purplish pink (54B) overlaid with dark red (187A); midribs light yellow (20B). Brilliant orange-yellow (23B) 15-mm-long blaze in centre of each tepal. Spots dark red (187A) scattered over lower three-quarters of each tepal; papillae absent; nectaries pale yellowish pink (36D), frosted at base; pollen strong reddish orange (169A); stigma brilliant orange (29A). Fls 140 mm wide; tepals 75 × 30 mm, margins smooth, tips slightly recurved. Lvs 75 × 8 mm, scattered, moderate olive-green (137A). Stems 1.1 m, moderate olive-green (137A), with up to 30 fls in a racemose infl. Mid-July (USDA Zone 3, Alberta, Canada). Published in *Valley K Greenhouses Cat.*, Spring 2006. Name reflects the diamond-shaped blaze in the nectary furrows and celebrates the North American Lily Society's 60th Anniversary Show held in Edmonton, Canada, in July 2007.

'Digital Globe' VIII(a/b) LA

Seedling BG 95-60 × seedling AJ 94-08

H: Vletter en Den Haan Beheer BV, pre-2006
N: Vletter en Den Haan Beheer BV, pre-2006

REG: Vletter en Den Haan Beheer BV, 2007
Inside of tepals light orange-yellow (24C); margins and midveins brilliant orange (25C); base of inner tepals closely spotted dark red (59A); throat pale yellow (11D). Outside of tepals pale orange-yellow (23D), shading to brilliant orange (25C) towards margins; midribs strong yellow-green: 143C at base, shading to N144D, with a small stripe of 144B at top. Spots very few, if any, dark red (59A); papillae few, brilliant orange (25C); nectaries moderate olive-green (137A); pollen greyish brown (166A); stigma strong orange (30D). Fls 190 mm wide; tepals 120 × 65 mm, margins smooth, tips recurved. Lvs 160 × 24 mm. Stems 1.1–1.5 m, green with darker markings, with up to 5 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Distant Drum' VII(b-a/c)

Seedling 9-13-SP × 'Broadway'

N: Rooyakkers Lelie BV, 2007 **REG:** Rooyakkers Lelie BV, 2007

Inside of tepals strong purplish red (60B), shading to 60D on paler ground towards margins; margins and throat greenish white (155C). Outside of tepals deep purplish pink on paler ground: outer tepals N66C, inner N66D; margins greenish white (155C); midribs of outer tepals tinged pale greenish yellow (1D). Spots dark red (59A); papillae absent; nectaries strong yellow-green (143B); pollen and stigma absent. Fls 195 mm wide, double; tepals 96 × 26–42 mm,

margins ruffled, tips recurved. Lvs 160 × 53 mm. Stems 0.8–1 m, pale green, with up to 6 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Divine’ VIII(a-b/a) LA

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Tepals deep red to strong purplish red (60A/B). Spots present; papillae absent; nectaries purple-red; pollen dark brown; stigma speckled purple. Fls 170 mm wide; tepals 158–165 × 50–60 mm, margins slightly ruffled, tips slightly recurved. Lvs 195–262 × 17–39 mm. Stems 0.9–1.05 m, green with darker stripes and spots, with 1–6 fls. Late.

‘Dolcetto’ VIII(b/b) LO

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2004 **N:** Marklily CV, 2004 **REG:** Testcentrum voor Siergewassen BV, 2007

Inside of tepals strong purplish pink (73B) to light purplish pink (73C), paler towards base, margins and tips; midveins pale purple-red at base. Outside of inner tepals light purple (75B) to very light purple (75C). Spots and papillae present; nectaries green; pollen red-brown; stigma grey-green. Tepals long and of medium width to broad, margins slightly ruffled, tips recurved. Lvs of medium length to long and of medium width to broad. Stems up to 1.45 m, green, with few fls. Very late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Doris Hall’ VIII(a/b) OA

White-flowered Asiatic × pink-flowered Oriental

H: R. Hall, 2003 **G:** R. Hall, 2006 **N:** R. Hall, 2006 **REG:** R. Hall, 2006

Inside of tepals golden orange-yellow; throat golden yellow. Outside of tepals pink. Brushmarks orange-red; nectaries orange-yellow; pollen and stigma orange-brown. Fls 112 mm wide; tepals 90 × 40 mm, margins smooth, tips slightly recurved. Lvs scattered, 160 × 20 mm. Stems to 1 m; green, mottled brownish green, with c.8 fls; bulbils present. Seed pods short and stout. Named after hybridizer’s mother.

‘Dublin’ I(a/-)

Parentage unknown

H: Fa Jaap Boon, pre-2002 **N:** Fa Jaap Boon, pre-2002 **REG:** Fa Jaap Boon, 2007

Inside of tepals strong orange (24A); midveins orange at base. Outside of tepals strong orange (25B). Spots and papillae present; nectaries green; pollen orange-brown; stigma purple-red. Tepals short to very short and narrow, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length and of narrow to medium width. Stems of average height to tall, green with darker stripes and spots, with few to average no. of fls. Early.

‘Du Chateau’ VII(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-1997

N: Vletter en Den Haan Beheer BV, 1997

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals deep to dark purplish pink (68A/186C) on pale ground; throat and above nectaries greenish white (155C). Outside of outer tepals dark to moderate purplish pink (186C/D); midribs very pale purple (73D), strong yellow-green (144B) at base and shading to brilliant yellow-green (154C) towards strong yellow-green (144A) top. Outside of inner tepals moderate purplish pink (186D); midribs very pale purple (76D), strong yellow-green (144B) at base and shading to brilliant yellow-green (150C) towards strong yellowish green (N144A) top. Spots absent; papillae moderate purplish pink (186D) and yellowish white (155D); nectaries strong yellow-green (144B); stigma light olive-grey (197B) with small, greyish purple (N77A) stripes. Fls 260 mm wide; tepals 150 × 85 mm, margins slightly ruffled, tips recurved. Lvs 192 × 54 mm, dark green. Stems 0.85–1.1 m, light green, with c.3 fls.

‘Dzeguzēni’ I(b/b-c)

‘Rotaļa’ × ‘Saulstarīte’

H: Ā. Zorgevics, 1984 **G:** Ā. Zorgevics, 1988

N: Ā. Zorgevics, 2006 **I:** University of Latvia Botanical Garden, 2006 **REG:** University of Latvia Botanical Garden, 2006

Tepals slender, light yellowish pink (26D); throat greenish. Brushmarks and evenly scattered small- to medium-sized spots, claret-red, mostly over basal third of each tepal; filaments pale, very long and widely exerted, pollen dark yellow; style long, incurving, stigma pale violet. Fls 120 mm wide; tepals 70 × 20 mm, margins slightly ruffled, tips twisted, recurved. Lvs scattered, 90 × 10 mm, glossy. Stems 0.9 m, purple-green, infl. long, with c.7 fls. Bulbs white. Early to mid-July. Published in *Lilium Balticum. Latvijas liliju selekcionāru izaudzētās lilijas*, No. 2 (1994). The name means Cuckoo Nestlings in Latvian.

‘Eagle’ VIII(a/b) LO

‘Tetra Sea Treasure’ × ‘Tetra Dame Blanche’

H: De Jong Lelies Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2006

Tepals and throat yellowish white (155D); inside of tepals, particularly midveins of outer tepals, flushed light purplish pink (65B); midribs of outer tepals strong yellow-green (144C) at base and top, shading to brilliant greenish yellow (151D) between; midribs of inner tepals strong yellow-green (143C) at base. Spots numerous, vivid purplish red (61C); papillae present; nectaries light yellow-green (144D); pollen strong reddish orange (169A); stigma very pale green (123D). Fls 210 mm wide, scented; tepals 140 × 77 mm, margins slightly ruffled, tips recurved. Lvs 170 × 52 mm, dark green. Stems 0.9–1 m, green, usually twin-flwd.

'Eberswalde' VI(c/d)

henryi × a mixture of pollen from 'Oneida', 'Lady Jane' and *L. × kewense* 'White Henryi'

H: G. Steinbrück, 1990 **G:** G. Steinbrück, 1994

N: G. Steinbrück, 2007 **I:** G. Steinbrück, 2007

REG: G. Steinbrück, 2007

Inside of tepals yellowish white (155D) with light yellowish pink (159A) central flare along and beside midvein in basal third. Outside of tepals white with mid-green midribs. Spots small, raised, of irregular shape, a few of the uppermost markings dashes, not raised, all reddish crimson, closely and evenly distributed on lower third of each tepal, mostly between the central flare and margins; papillae brilliant greenish yellow (6A); nectaries large, dark green, deep in throat; filaments yellowish white, very long; stamens large; pollen moderate reddish brown (175A); style almost as long as stamens; style and stigma pale yellowish green. Fls 80 mm wide, slightly scented; tepals 75 × 22 mm, margins smooth, tips strongly recurved. Lvs 130 × 30 mm, scattered. Stems 1.25 m, mid-green with reddish light marbling, with up to 15 fls. Mid- to late July (Germany).

Named after the city which hosted an international lily exhibition in 2007.

'Edessa' VII(b-a/b)

Seedling RW 96-17 × seedling 93-14

H: Vletter en Den Haan Beheer BV, pre-1994

N: Vletter en Den Haan Beheer BV, 1994

REG: Vletter en Den Haan Beheer BV, 2006

Tepals and throat yellowish white (155D); midveins yellowish white (155B); midribs of outer tepals strong yellow-green (144C) at base shading to pale greenish yellow (2D), and light yellow-green (144D) with strong yellow-green (143B) at top; midribs of inner tepals light yellow-green (144D) at base shading to yellowish white (155D). Spots moderate purple (79D) over one side of each inner tepal; papillae yellowish white (155D); nectaries strong to light yellow-green (144C/D); pollen dark reddish orange (173A); stigma dark purplish red (N79B). Fls 230 mm wide, scented; tepals 140 × 74 mm, margins of outer tepals strongly ruffled, margins of inner tepals smooth to slightly ruffled, all tips recurved. Lvs 165 × 37 mm. Stems 1.05 m, light green with darker markings at base, with c.3 fls.

'Egg Nog' II(c/c)

('Charlene' × 'Frontier') × 'Vernon White'

H: E.E. Fox, 1993 **G:** E.E. Fox, 1999 **N:** E.E. Fox, 2000 **I:** Fox Lily Ranch, 2006 **REG:** E.E. Fox, 2006

Inside of tepals light yellow (11B), with white stripe beside midveins; throat pale yellow (11C). Outside of tepals white; margins narrow, light yellow (11B); midribs deep pink (47D). Spots sparse, deep purplish red (187D), in basal half of each tepal; papillae absent; nectaries strong orange-yellow (22A); pollen strong reddish orange (31A); stigma light yellow (17D). Fls 78 mm wide, slightly scented; tepals 39 × 18 mm, margins smooth, tips slightly recurved. Lvs whorled, 175 × 40 mm. Stems 1.25 m, strong yellow-green (143B), mottled strong reddish purple (72A) near top, with up to 18 fls. Late June to early July.

Nomenclatural standard: colour print supplied by registrant (WSY 0100436). Fl. colours reminiscent of the ingredients of an eggnog.

'Ela' I(a-b/b)

'Korál' × 'Brushstroke'

H: K. Braun, pre-2006 **N:** K. Braun, pre-2006

I: K. Braun, 2006 **REG:** K. Braun, 2006

Tepals orange. Brushmarks dark brown, fairly large; spots and papillae absent; nectaries pale orange; pollen brown; stigma pale orange. Fls 110 mm wide; tepals 72 × 36 mm, tips recurved; pedicels 85 mm, dark brown; with secondary buds. Lvs 94 × 96 mm, pale green, glossy. Stems 0.9 m, stiff, spotted brown, infl. long. Mid-July.

'El Condor' VIII(b/b) LO

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007

REG: Testcentrum voor Siergewassen BV, 2007

Inside of tepals strong purplish red (60B to C), paler towards margins, tips and base; base of midveins green. Outside of inner tepals strong purplish red (71C) to strong reddish purple (72C). Spots and papillae absent; nectaries green; pollen orange-brown; stigma pale grey. Tepals long and broad, margins slightly ruffled, tips recurved. Lvs long to very long and of medium width to broad. Stems tall, green with darker markings, with few fls. Late to very late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'El Dorado' I(a/b)

In 1984, the lily registered in 1983 by Fa P. Hoff & Zn as 'Eldorado', was granted Dutch Plant Breeders' Rights under the cultivar epithet 'El Dorado', which becomes the spelling accepted by the ICRA (ICNCP, 2004: Art.24.4).

'El Santo' VII(a/b)

Red-flowered seedling × (seedling × ('Rosalie' × 'Star Gazer'))

H: Bischoff Tulleken Lelies BV, pre-2003

N: Bischoff Tulleken Lelies BV, 2003 **REG:** Mak Breeding BV, 2006

Inside of tepals dark red (59A) shading to strong purplish red (59C) towards margins; margins of inner tepals narrow, very pale purple (69D); throat and narrow margins of outer tepals yellowish white (155D). Outside of outer tepals moderate purplish red (59C); margins narrow, yellowish white (155D); midribs strong yellow-green at base and top (143B and 144C respectively). Outside of inner tepals strong purplish red (59D) shading to moderate purplish red (186A) towards margins; base yellowish white (155D); midribs strong yellow-green (144C) at base shading to moderate purplish red (186A), and light yellow-green (144D) at top. Spots and sparse papillae greyish purple (N77A); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma brilliant violet (86D). Fls 240 mm wide; tepals 125 × 57 mm, margins strongly ruffled, tips recurved. Lvs 190 × 50 mm. Stems 1 m, dark with green markings, with c.4 fls.

'Elysée' VII(a-b/b-c)

'Elysée' was registered by Royal Van Zanten Flowerbulbs BV in 2007. Details and a description of 'Elysée' were published in *ILR&C* 2007.

'Elzo' VIII(a-b/-) LA

Seedling BG 95-32 × seedling G 90-05

H: Vletter en Den Haan Beheer BV, pre-2006

N: Vletter en Den Haan Beheer BV, pre-2006

REG: Vletter en Den Haan Beheer BV, 2006

Inside and outside of inner tepals vivid yellow (12A); base of midveins yellow-green. Spots absent; papillae very sparse or absent; nectaries green; pollen orange-brown; stigma yellow-green. Tepals of medium length and of medium width to broad, margins slightly ruffled, tips slightly recurved. Lvs of medium length to long and of medium width. Stems 1.4 m, green, with up to 10 fls. Mid- to late season.

The registrant advises that this cv is no longer in cultivation.

'Emanuella' VII(a-b/b)

Seedling RM 96-65 × seedling RM 98-16

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals deep purplish pink (70C); inside of outer tepals light purplish pink (65B); inside of all tepals shading to strong purplish pink (68B) towards small yellowish white (155D) tip; above and beside nectaries brilliant yellow (11A); throat yellowish white (155D). Outside of tepals light purplish pink (63D); outer tepals yellowish white (155D) at base; inner tepals yellowish white (158C) at base shading to strong purplish pink (63C) at tips; all midribs strong yellow-green (144C) at base shading to yellowish white (155D); midribs of outer tepals strong yellow-green (144B) at top. Spots and papillae vivid purplish red (61C); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma very pale green (128D). Fls 240 mm wide, scented; tepals 135 × 76 mm, margins ruffled, tips recurved. Lvs 175 × 45 mm. Stems 1.2 m, dark at base shading to green with darker markings towards top, with c.4 fls.

'Embrun' I(a/c)

Seedling OR 93-21 × seedling OR 88-02

H: Vletter en Den Haan Beheer BV, pre-2006

N: Vletter en Den Haan Beheer BV, 2006

REG: Vletter en Den Haan Beheer BV, 2007

Inside of inner tepals deep red (53A), shading to dark red (187C) towards margins; margins dark red (187B), shading to deep red (185A) to strong red (53B) at tips; throat brilliant greenish yellow (1B). Inside of outer tepals deep red (53A), shading to strong red (53B) towards tips; margins dark red (187C). Outside of outer tepals moderate red (181B) to dark reddish orange (173A), shading to moderate red (181A) towards margins; margins deep red (53A); base with some strong yellow-green (144A). Outside of inner tepals moderate red (182A), shading to greyish red (184A) to moderate red (184B) towards margins; midribs moderate red (181A), with some strong yellow-green (144A) at base. Spots absent; papillae

deep red (53A); nectaries strong yellow-green (143B); pollen dark reddish orange (175B); stigma dark reddish orange (173A). Fls 177 mm wide; tepals 96 × 55 mm, margins smooth, tips strongly recurved. Lvs 176 × 13 mm, dark green. Stems 1.4 m, dark, with up to 8 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Enigma' VII(a/-)

Parentage unknown

H: J.A. and J. Moerman, pre-2007 **N:** J.A. and J.

Moerman, pre-2007 **REG:** J.A. and J. Moerman, 2007

Inside of tepals vivid purplish red (67B) to deep purplish pink (67C); midveins white at base, shading to purple-red. Outside of tepals strong purplish pink (68B). Spots numerous; papillae present; nectaries green; pollen orange-brown; stigma grey-green.

Tepals of medium length and width, margins strongly ruffled, tips (slightly) recurved. Lvs of medium length and of medium width to broad. Stems of average height to tall, green with faint darker markings, with few fls. Mid-season.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Enjoy' VII(a-b/-)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2006

N: Marklily CV, 2006 **REG:** Testcentrum voor Siergewassen BV, 2007

Inside of tepals deep purplish pink (68A) to strong purplish pink (68B) on paler ground, becoming paler with age and shading to nearly white towards base; midveins light yellow at base. Outside of inner tepals strong purplish pink (68B). Spots white on a pink background; papillae present; nectaries green; pollen orange-brown; stigma purple. Tepals of medium length and width, margins ruffled to strongly ruffled, tips recurved to strongly recurved. Lvs of medium length to long, and broad. Stems up to 1.1 m, green with darker markings, with up to 4 fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Enterprise NWT' I(c/d)

Parentage: 'Marlene' × orange-flowered I(b/c) seedling

H: D. Crook, 2001 **G:** D. Crook, 2003

N: D. Crook, 2007 **I:** Valley K Greenhouses, post-2008 **REG:** D. Crook, 2007

Inside of tepals and throat deep pink (48A). Outside of tepals open with rusty orange overlay which burns back to margins within a couple of days, becoming moderate red (N34C); midribs light yellowish pink (27A). Spots few, small, round, evenly distributed between throat and lowermost third of each tepal, greyish red (178A); papillae absent; nectaries strong yellowish pink (34D); pollen vivid reddish orange (33A); stigma moderate pink (49B). Fls 128 mm wide; tepals 60 × 25 mm, margins smooth, tips strongly recurved; pedicels 100 mm. Lvs 90 × 5 mm, scattered. Stems 1 m, strong yellow-green (143C), with up to 14 fls in a very open, racemose infl. Late July (NW Territories, Canada). The name is a community in Northwest Territories, Canada.

'Epirus' VIII(a/b-c) LA

Seedling BG 96-110 × seedling AG 95-11
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006
 Inside of tepals brilliant greenish yellow (5A); inner tepals strong yellow-green (144B) above nectaries; outer tepals strong yellow-green (144C) above and beside nectaries; throat light yellow-green (2C). Outside of tepals light greenish yellow (3C) shading to brilliant yellow (7A) towards margins; midribs strong yellow-green (143C) at base shading to light greenish yellow (3C); midribs of outer tepals with a small stripe of strong yellow-green (143B) at top. Spots and papillae absent; nectaries strong yellow-green (143B); pollen strong brown (172A); stigma light greenish yellow (4C). Fls 215 mm wide; tepals 115 × 53 mm, margins smooth, tips recurved. Lvs 153 × 20 mm. Stems 1.2–1.3 m, green with darker markings, with c.6 fls.

'Esperanto' VII(a-b/b)

Parentage unknown
H: Trior Lelie BV, pre-1991 **REG:** Royal Van Zanten Flowerbulbs BV, 2007
 Inside of tepals moderate purplish pink (62B) to light purplish pink (62C); midveins brilliant yellow (10A) to light yellow (10B); throat with a yellowish white (155B) flame. Outside of tepals light purplish pink (65B) on a yellowish white (155B) ground; midribs light yellow (10B). Spots and most papillae dark red (59A), those in throat white, tipped pale yellow; nectaries strong yellow-green (144B/C); pollen strong orange (169C); stigma greyish yellow-green (195B). Fls 300 mm wide; tepals 150 × 70 mm, margins slightly ruffled, tips slightly recurved. Lvs 150 × 40 mm. Stems 0.85 m, green, with up to 7 fls.

'Esprit' VIII(a/b) LA

'Jonita' × seedling LA 93007 T
H: De Jong Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2007
 Inside of tepals strong orange (inner 25A, outer N25B), shading to N25A towards margins; midveins vivid yellowish pink (28A); throat strong orange (25B). Outside of outer tepals strong orange (24A), shading to N25A towards margins; base dark red (183A) to moderate red (184B). Outside of inner tepals strong orange (N25C), shading to N25B towards strong orange (N25A) margins; midribs light olive (152B) at base. Spots few, dark red (59A); papillae absent; nectaries dark red (183A); pollen dark red (187A); stigma vivid reddish orange (32A). Fls 205 mm wide, slightly scented; tepals 125 × 56 mm, margins smooth, tips slightly recurved. Lvs 155 × 26 mm. Stems 1–1.1 m, dark at base, green at top, with up to 6 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Eurydice' I(c/d)

Mis-spelling of 'Eurydike'.

'Evas Feurige' I(c/d)

'Feuer und Rauch' × 'Ophelia'
H: E. Steinbrück, 1991 **G:** E. Steinbrück, 1994
N: E. Steinbrück, 2005 **I:** E. Steinbrück, 2006
REG: E. Steinbrück, 2006
 Tepals moderate red (179A). Spots blackish, of medium size, evenly scattered over basal third of each tepal; papillae absent; nectaries large, purple; filaments and style long, shading to reddish rose-pink in upper two-thirds, pollen greyish reddish brown (200B); stigma moderate reddish brown (175A). Fls 100 mm wide; tepals 55 × 35 mm, margins smooth, tips strongly recurved. Lvs scattered, 100 × 11 mm. Stems 1.1 m, dull dark green, with up to 17 fls. Mid- to late July. Nomenclatural standard: colour print supplied by registrant (WSY 0100435). The name means "Eva's Fiery One" in German. Eva is the first name of the hybridizer.

'Evas Freude' I(b-c/b)

'Golden Lady' × 'Ophelia'
H: E. Steinbrück, 1991 **G:** E. Steinbrück, 1994
N: E. Steinbrück, 2005 **I:** E. Steinbrück, 2006
REG: E. Steinbrück, 2006
 Inside of tepals and throat greyish reddish orange (174B) shading to light yellow (163D) towards margins and tips. Outside of tepals light yellow (163D); midribs broad, beige. Spots sparse, tiny, blackish brown, towards the centre of the basal quarter of each tepal; papillae absent; nectaries green-white; filaments long, slender, recurving, pollen dark brown; style long, tinged reddish rose-pink, stigma purple. Fls 125 mm wide; tepals 65 × 28 mm, margins smooth, tips slightly recurved. Lvs scattered, 70 × 16 mm. Stems 1 m, dark green, but dark violet where pedicels join the stem, with 8–12 fls. Early July. Nomenclatural standard: colour print supplied by registrant (WSY 0100434). The name means "Eva's Joy" in German. Eva is the first name of the hybridizer.

'Evas Traum' I(b-c/b)

'Feuer und Rauch' × 'Ophelia'
H: E. Steinbrück, 1991 **G:** E. Steinbrück, 1994
N: E. Steinbrück, 2005 **I:** E. Steinbrück, 2006
REG: E. Steinbrück, 2006
 Tepals moderate red (181A); midveins pale yellow (158B); throat flushed salmon pink. Spots fairly numerous, tiny, brown, evenly scattered, mostly over lower half of each tepal; papillae sparse, pinkish white; nectaries soft pink, becoming white with age; filaments long, slender, tinged pale rose-pink towards large anthers, pollen strong reddish orange (169A); stigma beige. Fls 140 mm wide; tepals 70 × 42 mm, margins smooth, tips recurved; pedicels dark. Lvs scattered, 92 × 16 mm. Stems 0.9 m, light green, but shiny red-orange where pedicels join the stem, with 9–10 fls. Mid- to late July. Nomenclatural standard: colour print supplied by registrant (WSY 0100433). The name means "Eva's Dream" in German. Eva is the first name of the hybridizer.

'Eyeliner' VIII(a/b) LA

(*longiflorum* 'White Fox' × 'Pulsar') × white-flowered Asiatic mixture

H: Testcentrum voor Siergewassen BV, pre-2004
N: Marklily CV, 2004 **REG:** Testcentrum voor Siergewassen BV, 2006

Tepals yellowish white (155D), midveins tinged strong yellow-green (144C) above throat; throat light yellow-green (144D). Beside midribs of outer tepals a few stripes of moderate red (180A); midribs strong yellow-green (143B) at base with moderate red (180A), shading to light yellow-green (144D) with some strong yellow-green (144A) towards top; midribs of inner tepals strong yellow-green (144C) flushed moderate red (180A), shading to light yellow-green (150D). All tepal-margins closely spotted dark red (59A); papillae yellowish white (155D); nectaries strong yellow-green (143A); pollen moderate reddish brown (166B); stigma greyish red (182B). Fls 192 mm wide; tepals 120 × 60 mm, margins smooth, tips recurved. Lvs 135 × 32 mm. Stems 1.2 m, green, with darker markings towards top, with c.5 fls.

'Fairfield' I(a-b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Inside of tepals vivid reddish orange to vivid orange (N30C/D). Outside of inner tepals vivid orange (N30D). Spots sparse; papillae inconspicuous; nectaries orange; pollen red-brown; stigma a purple-red star. Tepals short and of narrow to medium width, margins slightly ruffled, tips slightly recurved. Lvs of medium length and of narrow to medium width. Stems of average height to tall, green, with darker stripes and spots, with few to average no. of fls. Early.

'Faith' VIII(a-b/a) LA

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2004
N: Marklily CV, 2004 **REG:** Testcentrum voor Siergewassen BV, 2007

Inside of tepals moderate purplish red (70A) to strong reddish purple (70B), shading to nearly white towards base; midveins green at base. Outside of inner tepals moderate purplish red (70A) to strong reddish purple (70B). Spots and papillae absent; nectaries green; pollen orange-yellow; stigma pale yellow-green. Tepals of medium length to long and of medium width, margins (very) slightly ruffled, tips (very) slightly recurved. Lvs of medium length to long and of medium width. Stems of average height to tall, green with darker markings, with few fls. Very late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Firefox' I(a/c-d)

Parentage unknown

H: F. Fellner, 1998 **G:** F. Fellner, 2001 **N:** L. Westfall, 2005 **I:** Valley K Greenhouses, 2006 **REG:** Valley K Greenhouses, 2005

Upper half of inside of each tepal vivid reddish orange (34A), shading from light yellow (11B) at centre of each tepal, to vivid yellow (17B) with narrow margin

of vivid reddish orange (34A) in basal third; throat strong reddish orange (32B). Spots small, strong red (46A), in throat to nectaries; papillae slightly raised; nectaries frosted strong reddish orange (32B); filaments pale, fairly short, pollen strong red (46A); style conspicuous, dark orange-red in upper two-thirds, becoming stout towards pale stigma. Fls 132 mm wide; tepals 68 × 27 mm, margins smooth, tips slightly recurved. Lvs scattered, 115 × 19 mm. Stems 0.9 m, dark green, increasingly mottled purple towards base, with up to 16 fls in a raceme. Late July. Nomenclatural standard: colour print supplied by registrant (WSY 0100432).

'Flevo Surprise' I(a/b)

Parentage unknown

H: Flevo Lilies, pre-2006 **N:** Flevo Lilies, pre-2006
REG: Flevo Lilies, 2007

Inside of tepals brilliant yellow (8A) to light greenish yellow (8B), shading to pale yellow-green (4D) towards tips; base of midveins yellow-green flushed red. Outside of tepals pale yellow (8D). Spots and papillae absent; nectaries green; pollen orange-brown; stigma purple-red. Tepals short to very short and narrow, margins slightly ruffled, tips slightly recurved to recurved. Lvs of short to medium length and of narrow to medium width. Stems short, green, with few to average no. of fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Fort Liard NWT' I(a/b)

Light orange-flowered, brushmarked Asiatic (a/-) seedling × 'Red Artiste'

H: D. Crook, 2000 **G:** D. Crook, 2002 **N:** D. Crook, 2007 **I:** Valley K Greenhouses, post 2008 **REG:** D. Crook, 2007

Tepals vivid reddish orange (42A); midribs moderate reddish orange (42D); throat moderate red (185B). Spots greyish red (176A), evenly distributed in throat and over basal half of each tepal, more sparse towards nectaries; papillae greyish red (176A), sparse, in throat and to either side of nectary groove; nectaries vivid orange (N30D); pollen vivid reddish orange (N30A); stigma light yellowish pink (27A). Fls 100 mm wide; tepals 55 × 25 mm, margins smooth, tips slightly recurved. Lvs 65 × 10 mm, scattered. Stems 0.8 m, moderate reddish brown (176B), with up to 10 fls in a racemose infl. July (NW Territories, Canada). The name is a community in Northwest Territories, Canada.

'Fort Providence NWT' I(b/c)

Light orange-flowered, brushmarked Asiatic (a/-) seedling × 'Red Artiste'

H: D. Crook, 2000 **G:** D. Crook, 2002 **N:** D. Crook, 2007 **I:** Valley K Greenhouses, post 2008 **REG:** D. Crook, 2007

Tepals vivid yellow (16A); midribs flushed rusty orange in lowermost third; throat moderate reddish orange (35B). Spots very conspicuous, greyish brown (166A), small and numerous at base of each tepal, becoming progressively larger, more elongated and more sparsely distributed in upper half of each tepal,

coalescing to form a broad brushstroke either side of each midrib in lowermost third; papillae greyish brown (166A), in throat only; nectaries vivid yellow (13A); pollen vivid yellowish pink (28A); stigma light greenish yellow (8B). Fls 130 mm wide; tepals 75 × 30 mm, margins smooth, tips slightly recurved. Lvs 81 × 12 mm, scattered. Stems 0.9 m, dark reddish orange (178B), with up to 9 fls in a racemose infl. July (NW Territories, Canada).

The name is a community in Northwest Territories, Canada.

'Fort Resolution NWT' I(a/b)

L. × hollandicum × 'Lemon Tree'

H: D. Crook, 2001 **G:** D. Crook, 2003 **N:** D. Crook, 2007 **I:** Valley K Greenhouses, post-2008 **REG:** D. Crook, 2007

Inside of tepals vivid yellow (12A), shading to strong reddish orange (32B) towards margins; margins strong orange-yellow (24B); base pale yellow (11D); throat deep purplish pink (N74C). Outside of tepals strong reddish orange (34C), shading to strong yellowish pink (34D) beside midribs; midribs strong yellowish pink (34D). Spots small, neat, dark purplish red (N79), evenly distributed in throat and lowermost third of each tepal; papillae absent; nectaries vivid yellow (12A) and yellowish white (155B); pollen strong orange (N25A); stigma deep pink (50B). Fls 142 mm wide; tepals 90 × 31 mm, margins smooth, tips straight; with secondary buds. Lvs 110 × 34 mm, dense, scattered. Stems 1.46 m, dark purplish red (N79A), with up to 12 fls in a racemose infl. July (NW Territories, Canada).

The name is a community in Northwest Territories, Canada.

'Fort Simpson NWT' I(a/b)

'Mahogany' open-pollinated

G: D. Crook, 2004 **N:** D. Crook, 2007 **I:** Valley K Greenhouses, post 2008 **REG:** D. Crook, 2007

Inside of tepals brilliant yellow (7A) in basal half of each tepal, moderate red (179A) in upper half; throat moderate red (179A). Outside of outer tepals greyish yellow-green (194B); margins very light greenish blue (117B). Outside of inner tepals brownish orange (171B). All midribs greyish yellow-green (194B). Spots few, greyish brown (166A), sparsely distributed in a band in the lowermost third of each tepal, absent from the base of each tepal; papillae absent; nectaries brilliant yellow (7A); pollen strong orange-yellow (N163C); stigma greyish red (184A). Fls 130 mm wide; tepals 87 × 35 mm, scarcely overlapping and tapering to a claw at base, margins smooth, tips straight. Lvs 105 × 15 mm, very dense, scattered. Stems 1.95 m, strong yellow-green (143A), with up to 10 fls. July (NW Territories, Canada).

Seed collected by Don Allen in 2003. The name is a community in Northwest Territories, Canada.

'Fort Smith NWT' I(b/d)

'Burnished Rose' × pink-flowered Asiatic (b/c) seedling

H: D. Crook, 2002 **G:** D. Crook, 2004 **N:** D. Crook, 2007 **I:** Valley K Greenhouses, post 2008 **REG:** D. Crook, 2007

Inside of tepals deep pink (180D) shading to deep purplish pink (64D); margins moderate reddish orange (179C); throat deep pink (180D). Outside of outer tepals pale purplish pink (62D) shading to strong purplish red (59D) towards margins; midribs light yellow-green (142D). Outside of inner tepals deep purplish pink (58D) beside midribs; margins strong purplish red (59D); midribs light yellow (160B). Spots few, greyish brown (166A); papillae absent; nectaries pale purplish pink (65D); pollen and stigma moderate reddish orange (171A). Fls 60 mm wide, slightly scented; tepals 51 × 19 mm, margins smooth, tips strongly recurved; with secondary buds. Lvs 100 × 9 mm, scattered, bright dark green. Stems 1.09 m, strong yellow-green (143C), with up to 16 fls in a racemose infl. July (NW Territories, Canada). The name is a community in Northwest Territories, Canada.

'Freetown' VII(b-a/b-c)

Seedling RM 94-41 × seedling RW 98-12

H: Vletter en Den Haan Beheer BV, pre-1993

N: Vletter en Den Haan Beheer BV, 1993

REG: Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals strong purplish pink (73B); inside of outer tepals deep purplish pink (68A); all margins, above nectaries and throat, yellowish white (155D). Outside of outer tepals deep purplish pink (68A); midribs strong yellow-green (143A) at base shading to 143C and 144C. Outside of inner tepals moderate purplish pink (186D); midribs strong yellow-green (144C); all margins and base of inner tepals, yellowish white (155D). Spots vivid purplish red (61C); papillae yellowish white (155D); nectaries strong yellow-green (143C); pollen vivid yellowish pink (28A); stigma light grey (N200D). Fls 210 mm wide, scented, becoming paler with age; tepals 122 × 59 mm, margins ruffled, tips recurved. Lvs 197 × 41 mm. Stems 1.1–1.3 m, green, with c.5 fls.

'Galant' I(-/b)

H: W. Pauw & Zn, pre-1990 **REG:** Peacock BV, 2007

Tepals strong orange (30D), inside of tepals with vivid orange-yellow (23A) flame, outside of tepals with broader, light orange-yellow (24C) flame; midribs green-yellow. Spots and papillae absent; strong reddish orange pollen (169A) and stigma (32B). Fls 170 mm wide; tepals 90 × 50 mm. Lvs 170 × 20 mm. Stems 1.1 m, pale green, with up to 12 fls.

Trade designation BEATRIX. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Gander' VII(a/b)

Seedling 93-82 × seedling RW 98-12

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals moderate purplish pink (62B); margins and midveins strong purplish pink (68B); tips and throat yellowish white (155D). Inside of outer tepals moderate purplish pink (62B) shading to moderate purplish pink (68C) towards margins; tips, and beside and above nectaries, yellowish white (155D). Outside of outer tepals moderate purplish pink (186D);

yellowish white (155D) at base; midribs light yellow-green (145D) shading to strong yellow-green (144B) at base and top. Outside of inner tepals very pale purple (69A) shading to moderate purplish pink (70D) towards margins; yellowish white (155D) at base; midribs strong yellow-green (143C) at base shading to light yellow-green (150D), and strong yellow-green (144C) at top. Spots strong purplish red (63B); papillae yellowish white (155D) and strong purplish red (63B); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma very pale green (189D). Fls 260 mm wide; tepals 143 × 86 mm, margins ruffled, tips recurved. Lvs 195 × 58 mm. Stems 1.2–1.4 m, light green, with darker markings at leaf nodes, single flwr.

‘Garden Pleasure’ VIII(a-b/-) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Inside of tepals pale yellowish green (155A), flushed moderate to strong purplish red (59C/59D) at base; midveins darker purplish red. Outside of inner tepals pale yellowish green (155A). Spots purple-red; papillae inconspicuous; nectaries green; pollen red-brown; stigma purple. Tepals of medium length to long and of medium width, margins slightly ruffled, tips recurved. Lvs of medium length to long and of medium width. Stems of average height to tall, green, with few fls. Late to very late.

‘Passionale’ was used as a temporary name prior to registration.

‘Garnet Embers’ I(a/b)

‘Lily Simonet’ × pink-flowered seedling bred by Fred Fellner

H: T.D. Willoughby, 1991 **G:** T.D. Willoughby, 1993 **N:** S.K. Willoughby, 2002 **REG:** T.D. and S.K. Willoughby, 2006

Inside of tepals dark red (187A), shading to deep red (60A) towards margins; throat deep red (60A) with a light greenish yellow (1C), 10-mm blaze. Outside of tepals strong red (51A); midribs brilliant greenish yellow (1B), of more intense colour at top. Spots and papillae absent; nectaries deep red (53A) shading to deep pink (51C), slightly frosted white at base; pollen moderate orange (167A); stigma brilliant greenish yellow (1B). Fls 110 mm wide; tepals 75 × 30 mm, margins smooth, tips slightly recurved. Lvs 100 × 10 mm, scattered, moderate olive-green (137A). Stems 1 m, moderate olive-green (137A), with up to 30 fls in a racemose infl. Mid-July (USDA Zone 3, Alberta, Canada).

‘Gera’ I(a-b/b-c)

‘Feuer und Rauch’ × Asiatic seedling NN C7.44

H: G. Steinbrück, 1989 **G:** G. Steinbrück, 1992 **N:** G. Steinbrück, 2005 **I:** G. Steinbrück, 2006 **REG:** G. Steinbrück, 2006

Tepals and throat pale orange-yellow (24D); midribs greenish yellow. Spots numerous, tiny, violet, towards the centre of each tepal; papillae absent; nectaries large, yellowish white (155B); anthers long, slender, on long, white filaments; pollen greyish reddish

brown (200B); style exceptionally long, upward curving, soft orange; stigma moderate purplish red (58A). Fls 100 mm wide; tepals 72 × 28 mm, margins smooth, tips strongly recurved. Lvs scattered, 110 × 15 mm. Stems 0.75 m, dark green with light red marbling, with c.8 fls. Mid- to late July.

Nomenclatural standard: colour print supplied by registrant (WSY 0100431). Gera, situated in eastern central Germany, is the largest city in Thuringia. In 2007, it hosted the biennial federal horticultural show, BUGA.

‘Ghost Rider’ II(c/c)

‘Helsinki’ × *L. × dalbanonii* ‘Mondschein’ × ‘Vernon White’

H: E.E. Fox, 1994 **G:** E.E. Fox, 2000 **N:** E.E. Fox, 2001 **I:** Fox Lily Ranch, 2006 **REG:** E.E. Fox, 2006

Tepals white, inside of tepals flushed soft creamy yellow over central portion; throat and midribs brilliant yellow-green (149A and 149C respectively). Spots and papillae absent; nectaries and stigma brilliant yellow-green (150C); filaments relatively short, pale, yellowish, pollen vivid yellow (17B); style slightly longer than stamens, stout, yellowish green. Fls 74 mm wide, slightly scented; tepals 37 × 20 mm, margins smooth, tips slightly recurved; pedicels long, apple-green. Lvs whorled, 170 × 34 mm. Stems 1.45 m, light yellowish green (135D), with 14–18 fls. Late June to early July.

Nomenclatural standard: colour print supplied by registrant (WSY 0100429). The name is part of a song title.

‘Giotte’ VIII(b/b) OT

Seedling PG 95-39 × seedling RH 97-09

H: Vletter en Den Haan Beheer BV, pre-2004 **N:** Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals, including midveins, light greenish yellow (4B); upper half of tepals light greenish yellow (4C) to pale yellow-green (4D). Inside of outer tepals brilliant greenish yellow (4A) shading to light greenish yellow (4C) towards margins. Throat pale yellow-green (157D). Outside of outer tepals pale yellow-green (4D) flushed brilliant yellow-green (150C) along either side of midribs; midribs strong to light yellow-green (144C/D), and strong yellow-green (144A) at top. Outside of inner tepals light greenish yellow (3D); pale yellow-green (4D) along either side of midribs; midribs strong yellow-green (144C). Spots absent; papillae light greenish yellow (3C); nectaries strong yellow-green (144A); pollen moderate reddish brown (175A); stigma greyish purple (N77A). Fls 180 mm wide; tepals 115 × 55 mm, margins ruffled, tips recurved. Lvs 205 × 48 mm. Stems 1.2–1.5 m, light green, with c.4 fls.

‘Gluhwein’ VIII(a-b/b) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007

Inside of tepals strong red (51A) to deep pink (51B), paler towards margins and tips; midveins yellow-green at base. Outside of inner tepals pale orange-

yellow (23D) flushed red-purple. Spots absent; papillae few; nectaries white; pollen brown; stigma pale yellow-green. Tepals of medium length to long and of narrow to medium width, margins slightly ruffled, tips recurved to strongly recurved. Lvs of medium length to long and of medium width. Stems 1.45 m, green with darker markings, with up to 6 fls. Early to mid-season. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Go For Gold' VIII(a-b/b) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007
Tepals pale greenish yellow (inside 9D, paler towards tips; outside 10D); midveins vivid yellow (9B); midribs flushed red at base. Spots absent; papillae present; nectaries green; pollen orange-brown; stigma purple. Tepals of medium length and width, margins slightly ruffled to ruffled, tips recurved. Lvs of medium length to long, and broad. Stems 1.25 m, green, with up to 4 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Gold City' VIII(a/b) OT

Seedling PH 95-48 X AUB

H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2007
Inside of tepals brilliant yellow (7A); margins light greenish yellow (8C); tips pale yellow-green (4D); throat brilliant greenish yellow (7C). Outside of outer tepals brilliant yellow-green (150C), shading to pale yellow-green (4D) towards margins; margins light greenish yellow (4C); midribs strong yellow-green (144C). Outside of inner tepals pale greenish yellow (2D), pale yellow-green (4D) either side of midribs; midribs strong yellow-green (144B). Spots absent; papillae brilliant yellow (7A); nectaries strong yellow-green (144A); pollen moderate reddish orange (171A) to brownish orange (171B); stigma dark purplish red (N79A). Fls 220 mm wide; tepals 150 x 80 mm, margins slightly ruffled, tips recurved. Lvs 170 x 45 mm, dark green. Stems 1.05–1.4 m, pale green with darker markings, with up to 3 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Golden Eye' VIII(a/b) OT

Seedling 90.29.44 X seedling 90.126-11

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007
Inside of inner tepals brilliant yellow (10A) to light yellow (10B); inside of outer tepals light yellow (10B/C); paler towards margins and tips; midveins yellow at base. Outside of inner tepals light yellow (10C) to pale greenish yellow (10D). Spots and papillae absent; nectaries yellow; pollen orange; stigma green with purple spots. Tepals of medium length and width, margins slightly ruffled to ruffled, tips strongly recurved. Lvs of medium length to long, and of medium width to broad. Stems 1.5 m, green

with darker markings, with up to 4 small fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Golden Shiraz' VIII(b-c/b-c)

((('Tetra White Henry' x 'Tetra Black Beauty') x tetraploid Trumpets) X ((('Tetra White Henry' x 'Tetra Black Beauty') x tetraploid Trumpets)
H: B. Richardson, 2002 **G:** B. Richardson, post-2002
N: B. Richardson, 2005 **REG:** B. Richardson, 2006
Inside of tepals golden yellow in upper half, margins and base; central portion of basal half of each tepal deep red, extending further along midveins; throat golden yellow with green flares. Outside of tepals golden yellow, strongly flushed dark red, except at margins and tips; base green. Spots and papillae absent; nectaries, filaments and style apple-green; pollen orange; stigma dark grey. Fls 100 mm wide, strongly scented; tepals 80 x 25 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 120 x 25 mm. Stems 1.5 m, dark green with some darker markings towards base and leaf nodes, with c.15 fls. July. Shiraz is a red wine grape which probably originates from the eponymous city in southwest Iran.

'Gonzaga' VIII(a/b) LA

Seedling XC 00-02 X unknown seedling

H: Vletter en Den Haan Beheer BV, pre-2004
N: Vletter en Den Haan Beheer BV, 2004
REG: Vletter en Den Haan Beheer BV, 2007
Tepals, including throat, yellowish white (155D); inside of tepals closely spotted strong purplish red (60B) towards tips; outside of outer tepals with strong purplish red (60B) stripes on young fls, shading to light yellow-green (144D) beside midribs; midribs 144D also; midribs of inner tepals strong yellow-green (144C). Spots and papillae absent; nectaries strong yellow-green (144B); pollen dark reddish orange (175C); stigma brilliant yellow-green (150B). Fls 205 mm wide, slightly scented; tepals 115 x 68 mm, margins smooth, tips slightly recurved. Lvs 170 x 25 mm. Stems 1.3 m, green, with up to 4 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Gracia' VII(a-c-b)

Seedling RM 97-38 X seedling RW 96-05

H: Vletter en Den Haan Beheer BV, pre-2004
N: Vletter en Den Haan Beheer BV, 2004
Inside of tepals strong purplish red (64C) to deep purplish pink (64D); margins, tips, base and throat greenish white (155C). Outside of tepals deep purplish pink (64D); margins, tips and base greenish white (155C); midribs flushed light yellow-green (145B/C), and strong yellow-green at top (144A at top of midribs of outer tepals, 145A at top of inner tepals). Spots and papillae greenish white (155C); nectaries strong yellow-green (144A/B); pollen moderate reddish orange (N172A); stigma brownish grey (N200B) to greyish yellow-green (198A). Fls 215 mm wide; tepals 120 x 70 mm, margins ruffled, tips slightly recurved. Lvs 160 x 50 mm, dark green. Stems 1.2–1.25 m, pale green, with up to 5 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007). An application for registration was turned down in 2007, as the cultivar epithet is too similar to, and likely to cause confusion with, 'Grazia', which was registered by V. Strasser in 1986 (ICNCP, 2004: Art. 19.25). However, as Dutch Plant Breeders' Rights were granted to 'Gracia', the epithet remains the accepted name for this lily (ICNCP, 2004: Art. 28.2).

'Gradisca' VIII(a/c-b) OT

Seedling PH 95-39 × AUB

H: Vletter en Den Haan Beheer BV, pre-1999

N: Vletter en Den Haan Beheer BV, 1999

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals pale greenish yellow (9D), base and midveins brilliant yellow (9C); throat pale yellow-green (4D). Outside of outer tepals pale yellow (11D); midribs strong yellow-green at base (143C) and top (144C). Outside of inner tepals pale yellow-green (4D), shading to light greenish yellow (5D) towards margins; midribs light yellow-green (144D) at base. Spots and papillae brilliant yellow (9C); nectaries strong yellow-green (143C); pollen dark red (183B); stigma greyish reddish purple (N77C). Fls 210 mm wide; tepals 135 × 70 mm, margins ruffled, tips strongly recurved. Lvs 210 × 48 mm. Stems 1–1.1 m, green, with up to 4 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Hacienda' VIII(b-a/b) OT

Seedling PW 94-01 × seedling 92-97

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

Inside of tepals yellowish white (155D), flushed light yellow-green (150D) above nectaries; throat yellowish white (155D). Outside of outer tepals yellowish white (155D), flushed deep yellowish pink (39B) beside base of midribs; midribs light yellow-green (145B) at base and top, shading to 145C between. Outside of inner tepals yellowish white (155B); midribs strong yellow-green (144C) at base, shading to light yellow-green (145D). Spots and papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen moderate reddish orange (171A); stigma greyish reddish purple (N77C). Fls 225 mm wide, scented; tepals 140 × 72 mm, margins ruffled, tips recurved. Lvs 220 × 45 mm. Stems 1.3 m, green with darker markings (especially at base), with up to 3 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007). An application for registration was turned down in 2007, as the cultivar epithet is too similar to, and likely to cause confusion with, Hazienda Group, which was registered by K. Kiel in 1988 (ICNCP, 2004: Art. 19.25).

'Hay River NWT' I(b-c/d)

Pink-flowered Asiatic (b/c) seedling × dark red-mauve Asiatic (a/-) seedling

H: D. Crook, 2002 **G:** D. Crook, 2004 **N:** D. Crook, 2007 **I:** Valley K Greenhouses, post 2008 **REG:** D. Crook, 2007

Tepals, including throat, deep purplish red (61A). Spots small, strong reddish orange (N34B), widely

and quite densely distributed over lower three-quarters of each tepal; papillae absent; nectaries deep purplish red (61A); pollen strong reddish orange (N34B); stigma strong purplish pink (63C). Fls 100 mm wide; tepals 61 × 30 mm, margins smooth, tips strongly recurved. Lvs 100 × 10 mm, scattered. Stems 0.9 m, strong yellow-green (143C), with up to 34 fls in a pyramidal infl. Late July (NW Territories, Canada). Named after the community in which the hybridizer stays.

'Heather's Promise' I(a/c)

Parentage unknown

H: F. Fellner, 1999 **G:** F. Fellner, 2002 **N:** L. Westfall, 2005 **I:** Valley K Greenhouses, 2005 **REG:** Valley K Greenhouses, 2005

Inside of tepals white; a deep purplish pink (70C) flush in basal half extending to narrow, dark purple (79A) margins; throat greenish white (157D). Outside of tepals dark red (59A), fading at margins and tips. Spots large and numerous, evenly distributed, mostly over flushed basal half of each tepal, dark purple (79A); papillae raised; nectaries frosted white; filaments rather short, flushed pale rose-pink, pollen vivid reddish orange (34B); style much longer than stamens, shading to reddish pink towards deep red (53A) stigma. Fls 137–140 mm wide; tepals 77 × 28 mm, margins smooth, tips recurved. Lvs scattered, 122 × 11 mm, dark green. Stems 1.2 m, dark, mottled purple, with 15 or more fls. in a raceme. Buds, pedicels and stems downy. Late July to early Aug.

Nomenclatural standard: colour print supplied by registrant (WSY 0100430). Named for an admiring customer.

'Henri' V(b/a)

Parentage unknown

H: S. en S., V.o.f., pre-2007 **REG:** S. en S., V.o.f., 2007
Tepals greenish white (155C), flushed light yellow-green (145B/C) beside lower part of midveins; midribs yellowish white (155B). Spots and papillae absent; pollen dark yellow; stigma greenish white (157D) to pale green. Fls 180 mm wide, strongly scented; tepals 145 × 50 mm, margins slightly ruffled, tips strongly recurved. Lvs 220 × 36 mm. Stems 1.25 m, green, with up to 6 fls.

'Holland Beauty' VIII(b/-) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Inside of inner tepals moderate red (180A/B) shading to light to pale yellow (12C/D) towards margins; inside of outer tepals strong red (46A), paler towards margins; midveins strong red (46A) at base. Outside of inner tepals pale yellow (161C/D). Spots and papillae present; nectaries green shading to yellow-green; pollen orange-brown; stigma purple, spotted. Tepals long and of medium width to broad, margins slightly ruffled to ruffled, tips recurved to strongly recurved. Lvs of medium length and of medium width to broad. Stems of average height to tall, green, with few fls. Late.

'Honolulu' VIII(a/b) LA

Seedling BG 97-14 × seedling AG 95-10
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006
 Inside of tepals light greenish yellow (5D); tips closely spotted brilliant greenish yellow (5B); midveins brilliant greenish yellow (5B); throat pale greenish yellow (1D). Outside of tepals light greenish yellow (3D) shading to 4B towards margins; all midribs strong yellow-green (143C) at base shading to light greenish yellow (3D); midribs of outer tepals strong yellow-green (143C) at top; midribs of inner tepals light greenish yellow (4B). Spots and papillae absent; nectaries strong yellow-green (143A); pollen moderate reddish brown (166B); stigma light greenish yellow (1C) with deep purplish red (59B) edge. Fls 175 mm wide, younger fls of brighter colour; tepals 133 × 48 mm, margins smooth, tips recurved. Lvs 160 × 20 mm. Stems 1–1.4 m, green with darker markings, with c.6 fls.

'Hosteska' I(a-b/b)

((('Korál' × 'Juno' × ('Appleblossom' × 'Wattle Bird')) × 'Rozárka') × ('Blizzard' × ('Díblík' × 'Red King'))
H: B. Mičulka, pre-2005 **N:** B. Mičulka, pre-2005
I: B. Mičulka, 2006 **REG:** B. Mičulka, 2005
 Inside of tepals pale salmon-beige; central portion apricot, base pink; throat paler pink. Outside of tepals pinkish; midribs darker. Spots over half of each tepal, large, oval, fairly numerous, dark pinkish brown; pale papillae in throat; nectaries green, edged white; pollen ochre-apricot; stigma dark crimson. Fls 115 mm wide; tepals 80 × 40 mm, tips slightly recurved; pedicels 85 mm, with numerous brown spots. Lvs 80 × 8 mm, dull mid-green. Stems 0.85 m, dark brown. Mid-June.

'Hvězda' VI(b-c/c-d)

((('Andrea' × 'Sommertraum') × 'Henrieta') × ('Honeydew' × 'Gold Eagle') × (Silver Sunburst Group × 'Gold Eagle'))
H: B. Mičulka, pre-2007 **N:** B. Mičulka, pre-2007
I: B. Mičulka, 2007 **REG:** B. Mičulka, 2007
 Inside of tepals light yellow; tips whitish; midribs green. Lines of raised and papillate spots and fine furrows over two-thirds of each tepal dusted cinnamon brown; nectaries green; pollen apricot-brown; stigma green. Fls 85 mm wide; tepals 75 × 32 mm, rigid, of thick substance, tips strongly recurved; pedicels 70 mm, quite stout, green. Lvs 110 × 20 mm, scarce, dull, dark green. Stems 0.9 m, stiff, dark green, with average no. of fls. Early July.

'Ice Dancer' VII(a)

Parentage unknown
H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005
 Tepals yellowish white (155D). Spots absent; papillae present; nectaries light green; pollen orange-brown; stigma grey-purple. Tepals of medium length to long and of medium width to broad, margins ruffled to strongly ruffled, tips recurved. Lvs of medium length, broad. Stems of average height to tall, green, with few fls. Late.

'Indian Summerset' VIII(a/b) LA

(Unknown *longiflorum* hybrid × unknown Asiatic) × unknown Asiatic
N: Testcentrum voor Siergewassen BV, pre-2005
N: Marklily, 2005 **REG:** Testcentrum voor Siergewassen BV, 2006
 Inside of tepals strong purplish red (59D/60D); throat pale yellow-green (155A) flushed light yellow-green (144D). Outside of outer tepals moderate purplish red (59C); midribs strong yellow-green at base and top (143C and 144A respectively). Outside of inner tepals strong purplish red (59D); midribs strong yellow-green (143C) at base shading to strong purplish red (60C). Spots and papillae absent; nectaries strong yellow-green (143A); pollen strong reddish orange (169A); stigma greyish reddish purple (N77C). Fls 190 mm wide, becoming paler with age; tepals 132 × 61 mm, margins smooth, tips recurved. Lvs 185 × 23 mm. Stems 1.2 m, dark with a few green markings, with c.5 fls.

'Izmir' VII(a-b/b)

Seedling RW 94-09 × seedling RW 97-01
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006
 Tepals, including throat, yellowish white (155D); inside of tepals with a spot of light greenish yellow (4C) above nectaries. Midribs of outer tepals strong yellow-green at base and top (143B and 144C respectively), shading to strong yellow-green (144C) and pale greenish yellow (2D) between. Midribs of inner tepals strong yellow-green (144B) at base shading to light yellow-green (154D) and yellowish white (155D). Spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144C); stigma dark purplish pink (186C). Fls 195 mm wide, recurved; tepals 130 × 73 mm, margins ruffled, tips recurved. Lvs 175 × 49 mm. Stems 1–1.1 m, light green with darker markings at base, with c.3 fls.

'Jazz it Up' VIII(a/b-c) LA

'Paola' × tetraploid seedling LA 93092-1
H: De Jong Lelies Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2006
 Inside of tepals vivid purplish red (61C); throat light yellow-green (145C). Outside of tepals strong purplish red (60D) shading to 60C towards margins; midribs of outer tepals strong yellow-green (143C) at base shading to 143B at top; midribs of inner tepals strong yellow-green (143B) at base shading to strong purplish red (60C). Spots dark red (59A); papillae absent; nectaries strong yellow-green (143A); pollen dark reddish orange (175B); stigma pale purplish pink (62D). Fls 184 mm wide; tepals 100 × 53 mm, margins smooth, tips slightly recurved. Lvs 170 × 26 mm. Stems 0.9–1.1 m, green with a few darker markings, with c.6 fls.

'Jive' I(a/b-c)

((('Tetra Michella' × 'Saba Jewel') × 'Tetra Stella')
H: De Jong Lelies Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2006

Inside of tepals brilliant greenish yellow (6A), with a large, vivid reddish orange (N30A) blotch above the throat; throat light greenish yellow (3C). Outside of outer tepals light yellow (17D) shading to brilliant greenish yellow (6A) towards margins; midribs strong yellow-green (144C) at base, and deep purplish red (59B) at top. Outside of inner tepals brilliant greenish yellow (6A); midribs strong yellow-green (144C) at base shading to dark pink (182C). Spots few, dark red (59A); papillae brilliant greenish yellow (6A); nectaries strong yellow-green (143A); pollen moderate reddish orange (N172A); stigma dark purple (79A). Fls 195 mm wide; tepals 105 × 65 mm, margins smooth, tips slightly recurved. Lvs 140 × 25 mm. Stems 1–1.3 m, green with darker markings, with *c*.5 fls.

‘Joyce’s Favourite’

H: A.J. Porter, pre-2001 **N:** A.J. Porter, pre-2001
Pollen parent of ‘Yellowknife NWT’. Unregistered lily hybridized by A.J. Porter. No further details known.

‘Joy Hawley’ VI(c/a)

Unknown, open-pollinated Trumpets
S: D. Fielding, 2000 **G:** D. Fielding, 2000
N: D. Fielding, 2001 **I:** D. Fielding, 2004
REG: D. Fielding, 2007

Inside of tepals strong reddish orange (31B); midveins pinkish; throat light orange-yellow (19A). Outside of tepals vivid yellowish pink (28A); midribs red-purple. Spots and papillae absent; nectaries vivid yellowish green (140A); pollen brownish orange (164A). Fls 130 mm wide, slightly scented; tepals 105 × 45 mm, margins smooth, tips slightly recurved; with secondary buds. Lvs 130 × 12.5 mm, scattered, dark green. Stems 1.8 m, green with darker markings, with 30 or more fls. January (southern hemisphere). Nomenclatural standard: colour print supplied by registrant (WSY 0101430). Grown from seed collected in Joy Hawley’s garden.

‘Kastrup’ VIII(a/b) LA

Seedling BG 97-82 × seedling AG 97-15
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006
Inside of tepals vivid yellow (9A); margins and midveins vivid yellow (12A); throat brilliant yellow-green (154C). Outside of outer tepals brilliant yellow (7B) shading to vivid yellow (9A) towards margins; midribs brilliant yellow-green (150C) shading to strong yellow-green (143B) at base and top. Outside of inner tepals vivid yellow (9B) shading to 9A towards margins; midribs strong yellow-green (143C) at base shading to brilliant yellow-green (150B). Spots absent; papillae vivid yellow (9A); nectaries strong yellow-green (143A); pollen greyish brown (166A); stigma brilliant greenish yellow (1A) edged dark red (59A). Fls 210 mm wide; tepals 125 × 60 mm, margins smooth, tips recurved. Lvs 182 × 35 mm. Stems 1–1.3 m, green, with *c*.4 fls.

‘Key West’ VII(b-a/b-c)

Seedling 92-23 × seedling RW 98-03
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006
Inside of inner tepals deep purplish pink (70C); midveins deep purplish pink (N66C). Inside of outer tepals deep purplish pink (N66C), immediate tips and throat yellowish white (155D). Outside of outer tepals deep purplish pink (68A) shading to strong reddish purple (70B) towards margins; midribs strong yellow-green (145A), 144A at top. Outside of inner tepals deep purplish pink (N66D) shading to 72D towards margins; midribs strong yellow-green (144C); all midribs becoming brilliant yellow-green (154C) with age and outside of all tepals yellowish white (155D) at base. Spots strong purplish red (67A); papillae yellowish white (155D) with strong purplish red (67A); nectaries strong yellow-green (143C); pollen moderate to strong reddish orange (171A/169A); stigma very pale green (123D). Fls 220 mm wide, slightly scented; tepals 127 × 58 mm, margins slightly ruffled, tips recurved. Lvs 170 × 40 mm, dark green. Stems 1.1 m, green with darker markings, particularly at base, with *c*.5 fls.

‘Kingdom’ VIII(a/b-c) LA

Seedling BW 96-14 × seedling W 90-07
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006
Tepals yellowish white (155D); throat light yellow-green (145D). Outside of outer tepals with small, vivid purplish red (61C) stripes beside and along midribs; midribs strong greenish yellow (153C) at base. Outside of inner tepals with small, strong yellow-green (144C) stripes at base; midribs vivid purplish red (61C). Inside margins of inner tepals sparsely spotted deep purplish red (61A); elsewhere spots very few, strong purplish red (61B) or absent; papillae absent; nectaries strong yellow-green (143A); pollen brownish orange to dark reddish orange (164A/175C); stigma pale greenish yellow (10D). Fls 220 mm wide, young fls flushed yellow-green; tepals 127 × 70 mm, margins smooth, tips recurved. Lvs 180 × 25 mm. Stems 1.05–1.70 m, dark, with *c*.4 fls.

‘Kolotoč’ I(b-c/b)

‘Strohstern’ × ‘Blizzard’
H: B. Mičulka, pre-2006 **N:** B. Mičulka, pre-2006
I: B. Mičulka, 2006 **REG:** B. Mičulka, 2006
Inside of tepals dark yellow tinted beige; throat yellow. Outside of tepals paler yellow; midribs brownish. Spots and papillae absent; nectaries green, frosted white; pollen brown; stigma yellow. Fls 105 mm wide; tepals 75 × 35 mm, tips recurved; pedicels 70 mm, with numerous dark brown spots. Lvs 90 × 14 mm, dull mid-green. Stems 0.85 m, stiff, brown-black, infl. of medium length. Early July.

‘Koures’ VIII(a/b) LA

Seedling AM 97-14 × seedling BM 497-22
H: Vletter en Den Haan Beheer BV, pre-2004
N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007
Inside of inner tepals strong purplish red (59D), shading to moderate purplish red (59C) towards margins; base light purplish pink (62C); throat pale yellow-green (4D). Inside of outer tepals strong purplish red (60D) shading to (60C) towards margins; a spot of pale purplish pink (62D) at base. Outside of tepals deep purplish pink (N57C), shading to strong purplish red (63A) towards margins; midribs of outer tepals strong yellow-green (144C) at base, moderate purplish red (59C) at top. Inner tepals greenish white (155C) beside midribs at base; midribs tinged strong yellow-green (144C). Spots strong purplish red (58B); papillae absent; nectaries strong yellow-green (144A); pollen moderate reddish orange (N172A); stigma pale pink (49D). Fls 170 mm wide; tepals 115 × 58 mm, margins smooth, tips slightly recurved. Lvs 165 × 23 mm. Stems 1.3–1.4 m, with up to 5 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Kraton' VIII(b-a/b) OT

Seedling PG 95-49 × seedling RH 97-07
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006
Inside of inner tepals brilliant yellow (8A) shading to light greenish yellow (8C) towards tips; midveins vivid yellow (9B); throat pale yellow-green (4D). Inside of outer tepals light greenish yellow (8B); margins and midveins vivid yellow (9B). Outside of outer tepals pale yellow-green (4D); margins brilliant greenish yellow (4A); midribs light yellow-green (142D) at base shading to 145B, and strong yellow-green (143B) at top. Outside of inner tepals light greenish yellow (4C) shading to 6D at margins; pale yellow-green (4D) beside midribs; midribs light yellow-green (145C) at base shading to light greenish yellow (4C), and light yellow-green (154D) at top. Spots absent; papillae vivid yellow (9B); nectaries strong yellow-green (144C); pollen strong orange (N25A); stigma dark purple (79A). Fls 220 mm wide, scented; tepals 135 × 72 mm, margins slightly ruffled, tips strongly recurved. Lvs 225 × 35 mm. Stems 1.3–1.55m, green with darker markings at leaf nodes, with 4–5 fls.

'Krēsla' I(a/b-c)

'Rotaļa' × seedling 220-S-88
H: Ā Zorgevics, 1984 **G:** Ā. Zorgevics, 1988
N: Ā. Zorgevics, 2006 **I:** University of Latvia Botanical Garden, 2006 **REG:** University of Latvia Botanical Garden, 2006
Tepals deep yellowish pink (46D); throat light red. Spots black; papillae absent; nectaries brown; pollen light brown; stigma mahogany. Fls 130 mm wide, weather-resistant; tepals 80 × 40 mm, margins slightly ruffled, tips recurved. Lvs scattered, 80 × 15 mm, glossy. Stems 1.1 m, with c.9 fls. Bulbs white. Early to mid-July.
Published in *Lilium Balticum. Latvijas liliju selekcionāru izaudzētās lilijas*, No. 1 (1995) and *University of Latvia Botanical garden*, Riga (2002). The name means Nightfall in Latvian.

'Kyjov' VI(b-c/a-b)

Parentage unknown
H: B. Mičulka, pre-2007 **N:** B. Mičulka, pre-2007
I: B. Mičulka, 2007 **REG:** B. Mičulka, 2007
Tepals vivid yellow; midribs pale green. Nectaries green; pollen apricot; stigma green. Fls 120 mm wide; tepals 130 × 50 mm, rigid, margins slightly ruffled, tips strongly recurved; pedicels 65 mm, spotted dark brown. Lvs 95 × 14 mm, dark green. Stems 1.3 m, brown with black speckling, with a short infl. Early July.

'Labvakar' VI(b/a)

'Spitniece' × seedling L-11-35
H: Ā Zorgevics, 1990 **G:** Ā Zorgevics, 1994
N: Ā Zorgevics, 1995 **I:** University of Latvia Botanical Garden, 1995 **REG:** University of Latvia Botanical Garden, 2007
Inside of tepals brilliant yellow (9C); throat brilliant yellow-green (149B). Outside of tepals flushed pinkish brown; midribs dark. Spots and papillae absent; pollen dark brown; stigma grey. Fls 150 mm wide, strongly scented; tepals 130 × 40 mm, margins slightly ruffled, tips slightly recurved. Lvs 120 × 15 mm, scattered. Stems 1.7 m, green with darker markings, with up to 9 fls. Late July to early Aug.
Published in *University of Latvia Botanical garden*, Riga (2002). The name means Good Evening in Latvian.

'La Carita' VII(a-b/b)

Parentage unknown
H: Testcentrum voor Siergewassen BV, pre-2003
N: Hobaho, 2003 **REG:** Testcentrum voor Siergewassen BV, 2007
Inside of tepals yellowish white (155B), brilliant yellow (8A) beside midveins and at base. Outside of inner tepals yellowish white (155B). Spots yellow; papillae present; nectaries green; pollen orange-brown; stigma purple. Tepals of medium length and of medium width to broad, margins ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad to very broad. Stems tall, green with some darker markings, with few to average no. of fls. Mid-season to late.
Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Lake Manitoba' VII(a/b)

Red/pink-flowered seedling × 'Ascari'
H: Bischoff Tulleken Lelies BV, pre-2003 **N:** Bischoff Tulleken Lelies BV, 2003 **REG:** Mak Breeding BV, 2006
Inside of tepals vivid purplish red (61C) shading to yellowish white (155D) at margins, except towards the basal portion of margins of outer tepals, which are deep purplish pink (N57D); midveins strong red (53B); throat yellowish white (155D). Outside of outer tepals strong purplish red (61B) shading to deep purplish pink (N57D) towards margins; outside of inner tepals strong purplish red (64C). Outside of all tepals yellowish white (155D) at base; midribs strong yellow-green (144C) at base shading to yellowish white (155D), which is flushed strong yellow (153D) in inner tepals; top of outer midribs strong yellow (153D) with strong yellow-green (143A); top of inner midribs strong yellow-green (143B). Spots and papillae absent;

nectaries strong yellow-green (144C); pollen strong reddish orange (169A); stigma light purple (84B). Fls 195 mm wide, scented; tepals 115 × 65 mm, margins slightly ruffled, tips recurved. Lvs 155 × 40 mm. Stems 1 m, green with darker markings, with c.7 fls.

‘Lake Michigan’ VII(a/b)

Pink-flowered seedling × ‘Souvenir’

H: Bischoff Tulleken Lelies BV, pre-2003 **N:** Bischoff Tulleken Lelies BV, 2003 **REG:** Mak Breeding BV, 2006

Inside of tepals light purplish pink (63D), basal portion with large, yellowish white (155D) flame; throat yellowish white (155D). Outside of tepals strong purplish pink (63C); margins in basal portion of outer tepals yellowish white (155D); midribs light yellow-green (144D) at base shading to yellowish white (155D), and strong yellow-green (143A) at top; basal portion of inner tepals with large, yellowish white (155D) flame; midribs light yellow-green (144D) at base shading to strong purplish pink (63C), and brilliant yellow-green (150C) at top. Spots absent; papillae yellowish white (155D); nectaries light yellow-green (144D); pollen strong reddish orange (169A); stigma very pale green (189D). Fls 225 mm wide, scented; tepals 125 × 83 mm, margins ruffled, tips recurved. Lvs 150 × 49 mm. Stems 1.05 m, green, with c.7 fls.

‘Lake Winnipeg’ VII(a/b-c)

‘Color Parade’ × un-named seedling

H: Bischoff Tulleken Lelies BV, pre-2003 **N:** Bischoff Tulleken Lelies BV, 2003 **REG:** Mak Breeding BV, 2006

Inside of all tepals light purplish pink (62C); inside of outer tepals shading to pale purplish pink (65C) towards margins; all midveins with yellowish white (155D) towards top. Above and beside nectaries of inner tepals, brilliant yellow (10A) with some yellowish white (155D); outer tepals vivid yellow (9B) above nectaries, and yellowish white (155D) beside; throat yellowish white (155D). Outside of outer tepals dark purplish pink (186C) shading to pale purplish pink (65D) towards margins; some yellowish white (155D) beside midribs; midribs strong yellow-green (143C) at base shading to pale greenish yellow (2D), and strong yellow-green (143B) at top. Outside of inner tepals light purplish pink (65B), yellowish white (155D) at base; midribs strong yellow-green (143C) at base shading to pale yellow-green (155A) flushed light purplish pink (65B), and light yellow-green (145B) at top. Spots deep purplish pink (N57C); papillae brilliant yellow (9C); nectaries strong yellow-green (144B); pollen moderate reddish orange (N172A); stigma greyish yellow-green (148D). Fls 225 mm wide, slightly scented; tepals 130 × 65 mm, margins ruffled, tips recurved to strongly recurved. Lvs 160 × 63 mm, dark green. Stems 0.9–1.1 m, green, with 4–7 fls.

‘Laksmi’ VIII(a/b) LA

‘Amarone’ × seedling LA 93007-8T

H: De Jong Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2007

Inside of tepals deep red (53A); margins of outer tepals

dark red (187B); throat strong yellowish pink (37A/B). Outside of outer tepals dark red (187C), shading to deep red (60A) towards margins; midribs greyish red (178A) at base, strong yellowish green (N144A) at top. Outside of inner tepals moderate red (184B), shading to strong pink (50C) beside midribs in basal half; midribs dark red (187C). Spots and papillae absent; nectaries moderate olive-green (146A); pollen strong brown (172A); stigma dark purplish red (N79B). Fls 210 mm wide, darker when young; tepals 120 × 50–60 mm, margins smooth, tips recurved. Lvs 190 × 30 mm, glossy, dark green. Stems 1.1 m, green with darker markings, esp. in basal half, with up to 3 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Lecco’ VIII(a/b) LA

Seedling BG 97-14 × seedling AG 95-10

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals brilliant greenish yellow (5A); base light greenish yellow (4C) beside nectaries; throat light greenish yellow (4C). Outside of tepals light greenish yellow (5C); margins brilliant greenish yellow (5A); midribs strong yellow-green (143C) at base, shading to moderate yellow-green (143D). Spots and papillae absent; nectaries strong yellow-green (143A); pollen strong orange (25A); stigma brilliant yellow-green (150B). Fls 180 mm wide; tepals 134 × 52 mm, margins smooth, tips recurved. Lvs 200 × 30 mm. Stems 1.4–1.5 m, green with darker markings, with up to 6 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Lesotho’ VIII(b-a/b) OT

Seedling PG 96-43 × seedling RH 97-09

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of inner tepals brilliant yellow (7A/B), central portion towards tips light greenish yellow (7D); tips 5D; throat strong yellow-green (143B). Central portion of inside of outer tepals brilliant greenish yellow (7C), shading to 6C towards margins and light greenish yellow (5D) at tips. Outside of outer tepals pale yellow-green (4D), shading to brilliant yellow (10A) towards margins; central portion vivid yellow-green (154A) at base and tips; tips tinged strong yellow-green (144B). Outside of inner tepals brilliant yellow (10A), shading to 11A at margins; central portion pale yellow-green (4D). Spots absent; papillae vivid yellow (9A); nectaries strong yellow-green (143B); pollen dark red (183A); stigma dark purple (79B). Fls 220 mm wide; tepals 130 × 68 mm, margins slightly ruffled, tips recurved. Lvs 200 × 34 mm, dark green. Stems 1.3 m, pale green with darker markings, with up to 5 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Levi' I(a/c)

Seedling AM 94-08 × seedling AW 94-09

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2007

Inside of tepals yellowish white (155D) in basal two-thirds, deep purplish pink (61D) in upper third; margins narrow, deep purplish pink (61D); throat deep purplish pink (61D) with yellowish white (155D); outer tepals with a spot of light yellow-green (145C) above nectaries. Outside of outer tepals moderate purplish red (185C), shading to moderate purplish pink (186D) towards margins; margins vivid purplish red (61C). Outside of inner tepals deep purplish pink (58D); margins strong purplish red (58C); all midribs strong yellow-green (143C) at base, shading to moderate purplish red (186B). Spots and papillae absent; nectaries strong yellow-green (143B); pollen moderate reddish brown (166B); stigma pale yellowish pink (159D). Fls 165 mm wide, becoming paler with age; tepals 95 × 55 mm, margins smooth, tips slightly recurved. Lvs 155 × 35 mm. Stems 1.2–1.3 m, green, with up to 10 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Lianne' VII(a/b)

Seedling 91-79 × seedling 93-7

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals strong purplish red (64B) shading to moderate purplish red (186B) towards margins; light greenish yellow (6D) at base beside nectaries; midveins vivid purplish red (N57B), yellowish white (155D) at top; throat yellowish white (155D). Inside of outer tepals vivid purplish red (67B) shading to strong purplish red (64B) towards margins; yellowish white (155D) edged light greenish yellow (7D) at base beside nectaries; midveins strong purplish red (58B), yellowish white (155D) at top. Outside of outer tepals moderate purplish red (186B) shading to strong purplish red (63A) towards margins; yellowish white (155D) at base; midribs strong yellow-green (144C) at base shading to light yellow-green (150D), and strong yellow-green at top (144C with 144A). Outside of inner tepals dark purplish pink (186C) shading to strong reddish purple (70B) towards margins; yellowish white (155D) beside midribs; midribs strong yellow-green (144C) at base shading to light yellow-green (154D) with some strong reddish purple (70B). Spots strong purplish red (61B); papillae deep purplish pink (N57C); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma light yellowish green (136D). Fls 185 mm wide, scented; tepals 115 × 70 mm, margins ruffled, tips recurved. Lvs 190 × 42 mm. Stems 1.2 m, green with darker markings, with c.6 fls.

'Lodi' VII(b-a/b)

Seedling 92-100 × seedling RF 99-100

H: Vletter en Den Haan Beheer BV, pre-2004**N:** Vletter en Den Haan Beheer BV, 2004**REG:** Vletter en Den Haan Beheer BV, 2007

Inside of tepals deep purplish pink (N57D), shading

to strong purplish pink (73B) towards margins; inner tepals with little stripe of light greenish yellow (1C) at base above nectaries; midveins vivid purplish red (N57B); base of outer tepals yellowish white (155D) beside midveins and light greenish yellow (1C) above nectaries; throat yellowish white (155D). Outside of tepals deep purplish pink (73A), shading to strong purplish pink (73B) towards margins, yellowish white (155D) beside midribs at base; midribs strong yellow-green (144C) at base, shading to yellowish white (155D), and moderate yellow-green (146C) at top of outer tepals and strong yellow-green (144C) at top of inner tepals. Spots and papillae vivid purplish red (N57A); nectaries strong yellow-green (144C); pollen moderate reddish orange (171A); stigma very pale green (189D). Fls 220 mm wide, scented; tepals 125 × 58 mm, margins slightly ruffled, tips recurved. Lvs 230 × 47 mm. Stems 1.25–1.35 m, green, with up to 7 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Long Island' VIII(b-a/a) LO

'White Elegance' × seedling LO 93003-2B T

H: De Jong Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2007

Tepals yellowish white (155D), flushed light yellow-green (150D); lower two-thirds of midveins strong purplish red (58C) from above nectaries; throat light yellow-green (150D). Midribs of outer tepals light yellow-green (145B) shading to strong yellow-green (143A) at top; midribs of inner tepals light yellow-green (145D). Spots and papillae absent; nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma light greenish grey (190D). Fls 250 mm wide, slightly scented; tepals 195 × 75 mm, margins smooth, tips recurved. Lvs 250 × 41 mm. Stems 1.2 m, green, with up to 4 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Lucka' I(a/b)

'Montreux' × 'Latvia'

H: K. Braun, pre-2006 **N:** K. Braun, pre-2006**I:** K. Braun, 2006 **REG:** K. Braun, 2006

Tepals pink; throat brown. Spots over half of each tepal, dark brown; papillae absent; nectaries pinkish; pollen pale brown; stigma dark brown. Fls 130 mm wide; tepals 73 × 30 mm, tips slightly recurved; pedicels 80 mm, green. Lvs 80 × 17 mm, dull mid-green. Stems 0.6 m, stiff, green, infl. short. Late June.

'Lugums'

Unregistered seed parent of 'Sniega Meita'. No further details known.

'Mabel' VII(a)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2001**N:** Marklily CV, 2001 **REG:** Testcentrum voor Siergewassen BV, 2007

Tepals deep purplish pink (N66D) on pale background; inside of tepals paler towards margins and base; midveins white at base. Spots white; papillae present;

nectaries green; pollen reddish brown; stigma grey-green. Fls becoming paler with age; tepals of medium length to long, and of medium width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad to very broad. Stems of average height to tall, green with darker markings, with few to average no. of fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Macbeth' VIII(b-a/b) OT

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of inner tepals brilliant yellow (8A) to vivid yellow (9A), shading to light greenish yellow (4C) towards margins; greenish white (157D) at tips; throat strong yellow-green (143A). Inside of outer tepals brilliant yellow (8A) to vivid yellow (9A), shading to greenish white (157D) towards margins; light greenish yellow (4C) at tips. Outside of outer tepals pale greenish yellow (2D), shading to pale yellow-green (4D) towards margins; midribs vivid yellow-green (154A), tinged strong yellow-green (144B) at top. Outside of inner tepals pale yellow-green (4D); greenish white (157D) towards tips; midribs strong yellow-green (144B) at base and top. Spots absent; papillae vivid yellow (9A); nectaries strong yellow-green (143A); pollen greyish reddish brown (200B); stigma dark greyish reddish brown (200A). Fls 240 mm wide; tepals 135 × 76 mm, margins slightly ruffled, tips recurved. Lvs 215 × 52 mm, dark green. Stems 1.1–1.2 m, pale green with darker markings, with up to 4 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Mambo' VII(a-b/b)

Un-named seedling × un-named seedling

H: Sande BV, pre-2004 **N:** Sande BV, 2004

REG: Sande BV, 2007

Inside of tepals dark red (187C) to deep red (60A). Outside of inner tepals strong purplish red (60B/C). Spots few to average number, darker than tepal colour; papillae present; nectaries dark purple-red; pollen reddish brown; stigma grey-green with small purple spots. Fls 120–150 mm wide; tepals of medium length and of medium width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length, and of medium width to broad. Stems tall, 1–1.2 m, green with a few darker markings, with up to 7 fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Marchesa' VIII(a/b) LA

Seedling BW 96-19 × seedling AW 95-07

H: Vletter en Den Haan Beheer BV, pre-2001

N: Vletter en Den Haan Beheer BV, 2001

REG: Vletter en Den Haan Beheer BV, 2006

Tepals yellowish white (155D); inside of tepals with a small, strong yellow-green (144C) stripe above throat; throat light yellow-green (145C). Midribs of tepals

strong yellow-green (143B) at base shading to light yellow-green (144D); midribs of outer tepals strong orange-yellow (163B) at top. Spots and papillae absent; nectaries strong yellow-green (143A); pollen greyish brown (166A); stigma light yellow-green (154D). Fls 185 mm wide; tepals 110 × 60 mm, margins smooth, tips recurved. Lvs 160 × 20 mm. Stems 1.1–1.4 m, green with darker markings, with c.4 fls.

'Marco Polo' VII(b/b)

Syn. of 'Vedea'.

In 1993, 'Vedea' was granted a US Plant Patent under the cultivar epithet 'Marco Polo', which is the accepted name in countries in which 'Marco Polo' has legal protection. In *ILR&C 2007* 'Marco Polo' appears as a trade designation.

'Marlon' VII(a-b/b)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2001

N: Marklily, 2001 **REG:** Testcentrum voor Siergewassen BV, 2006

Inside of inner tepals deep purplish pink (N57D) shading to yellowish white (155D) towards margins and along edges of nectaries; midveins deep purplish pink (N57C); throat yellowish white (155D). Inside of outer tepals deep purplish pink (N57C/D); yellowish white (155D) at narrow margins. Outside of outer tepals dark purplish pink (186C); yellowish white (155D) at narrow margins; midribs light yellow-green (144D) at base shading to yellowish white (155D) flushed dark purplish pink (186C), and light yellow-green (144D) with strong yellow-green (144A) at top. Outside of inner tepals dark to moderate purplish pink (186C/D) shading to yellowish white (155D) towards margins; midribs dark purplish pink (186C) shading to light yellow-green (150D) at base and top. Spots and most papillae vivid purplish red (N57B), a few papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen strong reddish orange (169A); stigma light yellow-green (138D). Fls 215 mm wide; tepals 123 × 57 mm, margins ruffled, tips recurved. Lvs 155 × 45 mm. Stems 1.2 m, light green with darker markings, with c.6 fls.

'Mastermind' VIII(a/b) LA

Seedling BG 97-14 × seedling AG 95-10

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals brilliant yellow (12B); throat brilliant yellow-green (154B). Outside of outer tepals brilliant greenish yellow (7C) shading to brilliant yellow (7A) towards margins; outside of inner tepals brilliant yellow (8A); all midribs strong yellow-green at base (143C), with a small flush of red; midribs of outer tepals shading to brilliant greenish yellow (7C), and strong yellow-green (143B) at top; midribs of inner tepals shading to brilliant yellow (8A). Spots and papillae absent; nectaries strong yellow-green (143A); pollen strong orange (N25B); stigma greyish red (182B). Fls 180 mm wide; tepals 132 × 56 mm, margins smooth, tips recurved. Lvs 181 × 22 mm. Stems 1.15–1.5 m, dark with green markings, with c.5 fls.

'Mataro' I(a-b/-)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005
 Inside of tepals vivid yellowish orange to strong orange (30C/N25A), flushed purple-red at base; midveins strong orange (N25B) at base. Outside of inner tepals vivid reddish orange (33A/B). Spots absent; papillae present; nectaries orange-red; pollen red-brown; stigma purple-red. Tepals short, of narrow to medium width, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length to long and of narrow to medium width. Stems of average height to tall, green with darker stripes and spots, with few to average no. of fls. Mid-season to late.

'Matelotte' VII(a/c-b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2006
N: Vletter en Den Haan Beheer BV, 2006
REG: Vletter en Den Haan Beheer BV, 2007
 Tepals, including throat, greenish white (155C). Outside of outer tepals flushed pale greenish yellow (1D) beside midribs; midribs strong yellow-green at base (144C) and light greenish yellow (1C) towards strong yellow-green (145A) top. Midribs of inner tepals brilliant yellow-green (154C). Spots and papillae greenish white (155C); nectaries strong yellow-green (144B); pollen dark reddish orange (175B); stigma pale green (189C). Fls 250 mm wide; tepals 135 × 87 mm, margins ruffled, tips recurved. Lvs 235 × 75 mm, dark green. Stems 1.2–1.3 m, pale green, with up to 3 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'May Tay' VII(a-b/b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005
 Inside of tepals moderate purplish pink (65A), inner tepals shading to yellowish white (155D) towards base; midveins light greenish yellow (8B) at base. Outside of inner tepals light purplish pink (65B). Spots yellow; papillae present; nectaries green; pollen orange-brown; stigma purple. Fls 220 mm wide; tepals 152–160 × 52–80 mm, margins ruffled, tips slightly recurved to recurved. Lvs 89–212 × 19–56 mm. Stems 1–1.2 m, green with darker stripes and spots, with up to 6 fls. Late.

'Maywood' VIII(a/b-c) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2006
 Inside of tepals greyish purplish red (N77B) shading to very pale purple (76D) towards margins; outer tepals with narrower margins; inner tepals with a broad edge of pale yellow-green (155A) above and beside nectaries; outer tepals yellowish white (155D) beside nectaries and pale greenish yellow (2D) above; throat pale yellow-green (155A). Outside of tepals greyish purplish red (N77D); outer tepals shading to a narrow margin of very pale purple (76D); inner tepals on a white ground; all pale yellow-green (155A)

at base; all midribs strong yellow-green (144C) at base; midribs of outer tepals shading to dark greenish yellow (152D), and strong yellow-green (143A) at top; midribs of inner tepals shading to pale greenish yellow (2D) flushed moderate purplish pink (186D). Spots few, moderate purplish red (58A); papillae absent; nectaries strong yellow-green (144B); pollen strong orange (N163B); stigma very pale green (130D). Fls 280 mm wide, scented, becoming paler with age; tepals 145 × 65 mm, margins smooth to slightly ruffled, tips strongly recurved. Lvs 225 × 35 mm. Stems 1.3–1.7 m, green, with c.6 fls.

'Méd'ák' VI(c/b-d)

((('Damson' × unknown) × Golden Splendor Group) × unknown) × ((Royal Gold Group × unknown) × 'Africanka')

H: B. Mičulka, pre-2006 **N:** B. Mičulka, pre-2006
I: B. Mičulka, 2006 **REG:** B. Mičulka, 2006
 Tepals beige-orange; outside of tepals paler; midribs brownish pink at base; throat yellowish. Spots and papillae absent; nectaries green; pollen apricot-brown; stigma red-brown. Fls 135 mm wide; tepals 135 × 55 mm, margins slightly ruffled, tips recurved; pedicels 105 mm, with numerous, dark brown spots. Lvs scattered, 110 × 10 mm, dull mid-green. Stems 1.05 m, slender but stiff, with short infl. Late July.

'Merlet' VIII(a/b) LA

Seedling BW 96-33 × seedling AF 96-08

H: Vletter en Den Haan Beheer BV, pre-1999
N: Vletter en Den Haan Beheer BV, 1999
REG: Vletter en Den Haan Beheer BV, 2006
 Inside of tepals moderate purplish pink (186D); midveins deep purplish pink (N66D); throat light yellow-green (142D). Outside of tepals moderate purplish pink (186D) shading to dark purplish pink (186C) towards margins; midribs of outer tepals moderate yellow-green (146D) shading to strong yellow-green at base and top (143B and 143A respectively), with a little yellowish white (155D) at top; midribs of inner tepals strong yellow-green at base (143C) shading to dark purplish pink (186C), and moderate yellow-green (146D) at top. Spots and papillae absent; nectaries strong yellow-green (143B); pollen brownish orange (164A); stigma pale yellow (8D). Fls 195 mm wide, becoming paler with age; tepals 115 × 52 mm, margins slightly ruffled, tips recurved. Lvs 163 × 21 mm. Stems 1.1–1.65 m, dark with green markings, with c.4 fls.

'Metod' I(a-b/b)

'Rena' × ('Windahloo' × 'Jupiter')

H: B. Mičulka, pre-2005 **N:** B. Mičulka, pre-2005
I: B. Mičulka, 2006 **REG:** B. Mičulka, 2005
 Inside of tepals orange-red; tips paler. Outside of tepals pale orange; midribs yellowish. Brushmarks brown-red, slightly raised; spots numerous, over half of each tepal, of medium size, dark pink-brown; nectaries feathered white; pollen and stigma orange. Fls 140 mm wide; tepals 80 × 38 mm, tips slightly recurved; pedicels 55 mm, stout, green. Lvs 100 × 11 mm, dull mid-green. Stems 0.75 m, green, hairy at leaf nodes, infl. short. Late June.

'Mia' I(a/b)

'Romance' × 'Tango'

H: K. Braun, pre-2006 **N:** K. Braun, pre-2006

I: K. Braun, 2006 **REG:** K. Braun, 2006

Tepals yellow; throat yellow-green. Spots dense, over two-thirds of each tepal, brown; nectaries yellow; pollen brown; stigma yellow. Fls 160 mm wide; tepals 78 × 44 mm, tips slightly recurved; pedicels 100 mm, brown, with secondary buds. Lvs 60 × 15 mm, dark green, shiny. Stems 1 m, spotted brown, infl. long. Late June.

'Mistress' VIII(a-b/b) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV,

2003 **REG:** World Breeding BV, 2007

Inside of tepals light greenish yellow (8C) to pale yellow (8D), paler towards tips; midveins vivid yellow (9B) at base. Outside of inner tepals pale yellow (8D). Spots absent; papillae present; nectaries green; pollen brown; stigma grey-green. Tepals very short to short and of narrow to medium width, margins (very) slightly ruffled, tips slightly recurved. Lvs of short to medium length, and broad. Stems short, green, with up to 8 fls. Late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Mojeemoo' I(a/b-c)

Rust-red-flowered Asiatic (a/-) seedling bred by Fred Fellner × 'Marlène'

H: D. Crook, 2000 **G:** D. Crook, 2002 **N:** D. Crook, 2007 **I:** Valley K Greenhouses, post 2008 **REG:** D. Crook, 2007

Inside of inner tepals light yellow (12C) in basal third to half, upper portion strong yellowish pink (33C) over a whitish background, becoming darker towards tips; throat strong yellowish pink (33C). Outside of tepals strong yellowish pink (33C). Spots fairly numerous, greyish purple (N77A), evenly distributed over yellow portion of inner tepals, those nearest throat small and round, becoming slightly larger and more elongated towards centre of tepals; papillae few, fine, greyish purple (N77A), just above throat; nectaries vivid yellow (12A); pollen vivid orange (28B); stigma light yellowish pink (27A). Fls 155 mm wide; tepals 83 × 40 mm, margins smooth, tips slightly recurved; with secondary buds. Lvs 92 × 16 mm, scattered. Stems 0.8 m, strong yellow-green (144B), with up to 8 fls. Late July (NW Territories, Canada).

The nickname of the hybridizer's grand-daughter.

'Moneta' VII(a/b)

(Dwarf, white-flowered seedling × 'Time Out') × dwarf, white-flowered seedling

H: Bischoff Tulleken Lelies BV, pre-2003 **N:** Bischoff Tulleken Lelies BV, 2003 **REG:** Mak Breeding BV, 2006

Inside of tepals pale purplish pink (65D) on a yellowish white (155D) ground, with a spot of brilliant yellow (10A) above nectaries; throat yellowish white (155D). Outside of tepals yellowish white (155D) with pale purplish pink (65D); midribs strong yellow-green at base and top (143C and 144C

respectively), with some 144A at top of outer tepals, shading to light yellow-green (145C) between. Spots absent; papillae yellowish white (155D) and brilliant yellow (10A); nectaries strong yellow-green (144C) edged brilliant greenish yellow (6A); pollen dark orange (N163A); stigma light greenish grey (N189D). Fls 170–240 mm wide, scented, the pale purplish pink (65D) flush fading with age; tepals 104 × 46 mm, margins slightly ruffled, tips recurved. Lvs 170 × 43 mm. Stems 0.45–0.65 m, green, with c.6 fls.

'Monsela' VII(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Tepals, including throat, yellowish white (155D); midribs of outer tepals strong yellow-green (144C) at base shading to yellowish white (155D) at about mid-way and light yellow-green (150D) and strong yellow-green (144C) above; outer tepals with strong yellow-green (143B) at top. Spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen orange-brown; stigma greenish white (192D). Fls 220 mm wide, slightly scented; tepals 135 × 70 mm, margins ruffled, tips recurved. Lvs 192 × 55 mm. Stems 0.9–1 m, green with darker markings, with c.3 fls.

'Mont de Marsan' VII(b-a/b)

Seedling RW 95-20 × seedling RW 99-31

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Tepals, including throat, yellowish white (155D). Midribs light yellow-green (144D) at base, shading to yellowish white (155D); those of outer tepals light yellow-green (150D) at top, tinged strong yellow-green (143C); midribs of inner tepals light yellow-green (154D) at top. Spots and papillae yellowish white (155D); nectaries light yellow-green (144D); pollen moderate reddish orange (N172B); stigma dark purplish red (N79B). Fls 220 mm wide, slightly scented; tepals 120 × 70 mm, margins slightly ruffled, tips recurved. Lvs 182 × 36 mm. Stems 1.25–1.3 m, pale green, with up to 5 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Monte Cassino' I(a/b)

Parentage unknown

H: P. Hoff, pre-1987 **REG:** P. Hoff, 2007

Inside of inner tepals greyish red (178A) to moderate red (179A), shading to strong orange (169B) at base; throat moderate red (181C). Inside of outer tepals strong red (46A) to moderate red (181A); base and lower portion of midveins strong orange (168A).

Outside of outer tepals greyish red (184A), tips light green; inner tepals greyish red (178A) to moderate red (179A); midribs moderate red (181A), 180A/B at base. Spots dark brown; papillae present; pollen dark brown; stigma purple-red. Tepals of medium length and of medium width, margins smooth, tips recurved to strongly recurved. Lvs of medium length to long,

and of medium width, pale green. Stems tall, pale green with some darker markings.

Thought to be no longer in cultivation.

'Moonwalker' I(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals vivid yellow (9A), shading to (12A) towards tips and margins of inner tepals; throat pale greenish yellow (1D). Outside of tepals brilliant greenish yellow (5A), outer tepals tinged strong yellow-green towards tips (144B) and base (144C), inner tepals light greenish yellow (5C) beside midribs; midribs of inner tepals light yellow-green (144D) at base, shading to brilliant greenish yellow (5A). Spots absent; papillae few, vivid yellow (9A); nectaries strong yellow-green (144B); pollen moderate reddish brown (175A); stigma deep red (60A). Fls 163 mm wide; tepals 102 × 49 mm, margins smooth, tips slightly recurved. Lvs 130 × 25 mm. Stems 1.2–1.55 m, basal two-thirds dark, upper one-third green with darker markings, with up to 7 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Mother's Day' I(a/b)

(('Gran Paradiso' × 'Compass') × un-named seedling) × un-named seedling

H: Bischoff Tulleken Lelies BV, pre-1999 **N:** Bischoff Tulleken Lelies BV, 1999 **REG:** Mak Breeding BV, 2006

Inside of tepals vivid yellow: inner 17B, outer 16A, shading to vivid orange-yellow (21A) towards margins; with a vivid reddish orange (33B) spot above throat, larger on outer tepals; throat brilliant yellow (20A). Outside of tepals pale yellow (20C); outer tepals flushed moderate red (180B) on one side only; inner tepals shading to brilliant yellow (20A) towards margins; midribs of outer tepals strong yellow-green (143B) at base and top, shading to moderate red (180B) between; midribs of inner tepals light yellow-green (144D). Spots few, strong red (53B); papillae vivid yellow (16A); nectaries strong yellow-green (143B); pollen moderate reddish orange (171A); stigma deep purplish red (59B). Fls 190 mm wide; tepals 110 × 56 mm, margins smooth, tips recurved. Lvs 175 × 21 mm. Stems 0.9 m, dark with green markings, with c.7 fls.

'Mount Dragon' I(a-b/b)

Parentage unknown

H: Mak 't Zand BV, pre-2005 **N:** Mak 't Zand BV, 2005 **REG:** Mak Breeding BV, 2006

Inside of tepals vivid reddish orange (34A/B) to strong reddish orange (42B), shading to moderate reddish orange (42D) towards base and vivid red (45B) towards margins; midveins strong red (39A). Outside of tepals vivid reddish orange (34B) to moderate reddish orange (35B), shading to (35A) towards base and vivid reddish orange (44C) towards margins. Spots few; papillae present; nectaries orange; filaments moderate yellowish pink (39C) towards

base, strong red (39A) above; anthers and pollen strong orange (169B); style strong reddish orange (34C); stigma vivid reddish orange (33B). Buds very downy, vivid reddish orange (34A/B). Fls 150 mm wide, slightly scented; tepals 70 × 35 mm, margins slightly ruffled, tips slightly recurved to recurved; pedicels stout, moderate olive-green (146A). Lvs 75–100 × 15–17 mm, scattered, deep yellowish green (141A). Stems 0.4 m, strong yellow-green (144A) with darker markings, with up to 10 fls. Early: June. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Mount Duckling' I(a-b/b)

Parentage unknown

H: Mak 't Zand BV, pre-2005 **N:** Mak 't Zand BV, 2005 **REG:** Mak Breeding BV, 2006

Tepals deep purplish pink (N57D) to strong purplish pink (63C), inside of tepals shading to yellowish white (155D) towards base and becoming much paler with age. Spots few, dark red (187A); papillae present; nectaries pink, base green; filaments yellowish white (155B) towards base, strong purplish pink (63C) above; anthers moderate reddish brown (166B); pollen strong orange (26B); style strong purplish pink (63C); stigma strong purplish red (60D). Fls 140 mm wide, slightly scented; tepals 75 × 38 mm, margins (very) slightly ruffled, tips slightly recurved; pedicels moderate yellow-green (146B). Lvs 72–80 × 7–15 mm, scattered, deep yellowish green (141A). Stems 0.5 m, moderate olive-green (137B) with darker markings, with up to 10 fls. Early June.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Mueßer Stern' I(a/b)

'Alena' × orange-flowered, brushmarked seedling St. 154/2/89

H: G. Steinbrück, 1992 **G:** G. Steinbrück, 1995

N: G. Steinbrück, 2006 **I:** G. Steinbrück, 2006

REG: G. Steinbrück, 2007

Tepals moderate orange-yellow (168D), inside of tepals becoming darker towards base; midveins strong orange (169D); throat moderate orange-yellow (168D). Spots few, medium to large, red-violet, scattered over basal one-third of each tepal; nectaries greenish white; pollen dark reddish orange (172B); stigma beige. Fls 135 mm wide, scented; tepals 75 × 33 mm, margins smooth, tips slightly recurved. Lvs 90 × 10 mm, scattered. Stems 0.7 m, dark green, with up to 7 fls. Mid-July. Name means Star of Müß in German. Müß is a district of Schwerin, the capital city of the state of Mecklenburg-Vorpommern in northern Germany.

'Nashville' VIII(b-a/b) LA

Seedling BG 95-60 × seedling G 01-10

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals vivid yellow (12A); throat light yellow-green (2C). Outside of tepals brilliant greenish yellow (6C) to vivid yellow (9B), shading to vivid yellow (12A) at margins; midribs tinged green. Spots absent; papillae few; nectaries strong yellow-green

(143A/B); pollen strong orange (25A); stigma vivid yellow (9A). Fls 200 mm wide; tepals 110 × 55 mm, margins smooth, tips slightly recurved. Lvs 190 × 30 mm. Stems 1.2–1.5 m, green, with up to 4 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Navarrosse’ VII(a/b)

Seedling RW 99-30 × seedling RW 96-05

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2007

Tepals, including throat, yellowish white (155D).

Midribs light yellow-green (144D) at base, shading to

145D, midribs of outer tepals strong yellow-green

(143C) at top; those of inner tepals light yellow-green

(145B) at top. Spots and papillae few, yellowish white

(155D); nectaries strong yellow-green (144C); pollen

moderate reddish orange (171A); stigma very pale

green (189D). Fls 230 mm wide, scented; tepals 134 ×

58 mm, margins ruffled, tips recurved. Lvs 54 × 18 mm.

Stems 1.2–1.3 m, green, with up to 4 fls.

Published in *Registraties* week 44-2006 t/m 45-2007,

KAVB (2007).

‘Negritta’ I(a/b)

Seedling OR 91-15 × seedling AG 97-15

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals strong orange (N25C); inner tepals

with a narrow margin at tips and a large spot above

throat of strong orange (N25A); outer tepals shading to

strong orange (N25B) towards margins; throat strong

orange (26B). Outside of outer tepals vivid orange-

yellow (23A) shading to strong orange (25A) towards

margins; midribs moderate olive-green (146A) at base,

deep yellowish green (141A) at top. Outside of inner

tepals strong orange-yellow (24B) shading to strong

orange (25B) towards margins; midribs dark greenish

yellow (152C) at base shading to strong greenish yellow

(153C). Spots and papillae absent; nectaries strong

yellow-green (143A); pollen moderate red (179A);

stigma vivid reddish orange (42A). Fls 160 mm wide;

tepals 98 × 46 mm, margins smooth, tips recurved.

Lvs 142 × 11 mm. Stems 1.2–1.4 m, dark, with c.6 fls.

‘Nemo’ VIII(a/b) LA

‘Jonita’ × seedling LA 93007 T

H: De Jong Beheer BV, pre-2005 **N:** De Jong Lelies

Holland BV, 2005 **REG:** De Jong Lelies Holland

BV, 2007

Inside of tepals strong orange (25A); inner tepals

shading to N25B towards margins, outer to N25A;

midveins with a stripe of moderate red (N34A) above

nectaries which shades to strong orange (N25A) in

inner tepals; throat strong orange-yellow (24B).

Outside of tepals strong orange (outer 25B, inner

24A); outer tepals shading to vivid orange (28B)

towards margins; basal third and midribs greyish red

(178A); midribs strong yellow-green (144A) towards

top; inner tepals shading to vivid yellowish pink

(28A) towards margins; all of midribs greyish red

(178A). Spots and papillae absent; nectaries moderate

olive-brown (N199A); pollen greyish red (178A);

stigma vivid reddish orange (30A). Fls 200 mm wide,

scented; tepals 117 × 53 mm, margins smooth, tips

recurved. Lvs 190 × 38 mm. Stems 1–1.2 m, green

with darker markings at the base, with up to 6 fls.

Published in *Registraties* week 44-2006 t/m 45-2007,

KAVB (2007).

‘Missouri Garnet’ I(a/c)

Red-flowered Asiatic (a/-) seedling × magenta-

flowered Asiatic (a/-) seedling

H: Hartle-Gilman Gardens, 1992 **G:** E. Horner, 1995

N: E. Horner, 2007 **I:** Horner Lilies, 2007

REG: E. Horner, 2007

Inside of tepals deep red (60A); throat dark red (187A).

Outside of tepals strong purplish red (60B). Spots

absent; papillae white; nectaries deep red (60A); pollen

rust-coloured; style red, stigma cream. Fls 130 mm

wide; tepals 85 × 35 mm, margins slightly ruffled, tips

slightly recurved. Lvs 240 × 17 mm, scattered, strong

yellow-green (144A). Stems 1.1 m, dark red-brown,

with up to 18 fls. July (Ontario, Canada).

Nomenclatural standard: colour print provided by

registrant (WSY 0101433). Nissouri is an Indian

word meaning “running water” to describe an area

drained by creeks and streams, and was the former

name of the township in which the registrant stays.

‘Missouri Ice Wine’ I(a/c)

‘Juanita’ × Dark wine-red-flowered Asiatic (a/-)

seedling hybridized by V. Bowen

H: D. Schaefer, 1992 **G:** E. Horner, 1994

N: E. Horner, 2007 **I:** Horner Lilies, 2007

REG: E. Horner, 2007

Tepals glossy, dark red (187A inside, 59A outside);

throat moderate red (179A). Spots few, dark red;

papillae white; nectaries dark red (187A); pollen

rust-brown; stigma dark brown. Fls 110 mm wide;

tepals 70 × 25 mm, scarcely overlapping and tapering

to a claw at base, margins slightly ruffled, tips slightly

recurved. Lvs 100 × 12 mm, scattered, moderate

olive-green (146A). Stems 0.9 m, dark red-brown,

with up to 12 fls. July (Ontario, Canada).

Nomenclatural standard: colour print provided by

registrant (WSY 0101432). Nissouri is an Indian

word meaning “running water” to describe an area

drained by creeks and streams, and was the former

name of the township in which the registrant stays.

‘Missouri Sunflare’ I(a/b)

‘Yellow Princess’ × ‘Apricot Supreme’

H: D. Schaefer, 1993 **G:** E. Horner, 1995

N: E. Horner, 2007 **I:** Horner Lilies, 2007

REG: E. Horner, 2007

Tepals vivid yellow (17C), outside of tepals reddish

brown at base; throat strong yellow-green (144B).

Brushmarks short, dense, creating a patch of dark red

(59A) in lowermost quarter of each tepal; spots few,

small, dark red (59A), mostly either side of brush-

marks; papillae absent; nectaries strong yellow-green

(144B); pollen brown; stigma rust-coloured. Fls 120

mm wide; tepals 75 × 40 mm, margins slightly

ruffled, tips straight; with secondary buds. Lvs 100 ×

18 mm, scattered, strong yellow-green (144B). Stems

1.3 m, strong yellow-green (144B), with up to 24 fls. July (Ontario, Canada).

Nomenclatural standard: colour print provided by registrant (WSY 0101431). Missouri is an Indian word meaning "running water" to describe an area drained by creeks and streams, and was the former name of the township in which the registrant stays.

'Norah' VIII(a/b) LA

(Seedling 3473-vi × seedling 2008RS zl) × seedling LA 93064-14T

H: De Jong Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2007

Inside of tepals strong purplish pink (63C), base of inner tepals shading to deep purplish pink (64D) towards margins; midveins and stripes at the base of outer tepals strong purplish red (64C); throat pale yellow (8D). Outside of tepals dark purplish pink (186C), outer tepals shading to moderate purplish red (186B) towards margins, base of inner tepals yellowish white (155D) beside midribs; midribs strong yellow-green (144B) at base, shading to purplish pink tinged pale green. Spots few, strong purplish red (60D); papillae absent; nectaries moderate olive-green (137A); pollen dark reddish orange (175B); stigma pale purplish pink (65C). Fls 210 mm wide, scented; tepals 120 × 62 mm, margins smooth, tips recurved. Lvs 195 × 28 mm. Stems 1.2–1.3 m, green, with up to 5 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Northwest Territories' I(a/c)

Un-named seedling hybridized by F. Fellner × 'Joyce's Favourite'

H: D. Crook, 2001 **G:** D. Crook, 2003 **N:** D. Crook, 2007 **I:** Valley K Greenhouses, post-2008 **REG:** D. Crook, 2007

Tepals vivid reddish orange: inside of tepals 32A on opening, 33A by second day; outside of tepals 32A; throat N30A. Midribs brilliant orange (25C). Spots very sparse, greyish reddish brown (200B), mostly on basal third of each tepal; papillae absent; nectaries strong reddish orange (32B) to strong yellowish pink (32C); pollen dark reddish orange (173A); stigma moderate orange-yellow (168D). Fls 130 mm wide; tepals 70 × 26 mm, margins smooth, tips slightly recurved. Lvs 122 × 11 mm, scattered, mid-green. Stems 1 m, moderate reddish brown (177A), with up to 13 fls. in a racemose infl. Mid-July (NW Territories, Canada).

Named after the territory of Canada in which the hybridizer stays.

'Novatore' VIII(a/a-b) OT

Seedling PG 96-43 × seedling RH 97-09

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals brilliant yellow: inner 11A, outer 12B; tips of inner tepals light yellow (10B); midveins of outer tepals vivid yellow (12A); throat pale yellow-green (4D). Outside of outer tepals pale yellow-green (4D) shading to light greenish yellow (5D) towards margins;

midribs light yellow-green (144D) at base shading to pale yellow-green (4D), and strong yellow-green (144A) at top. Outside of inner tepals light greenish yellow (5C); beside and along midribs pale yellow-green (4D); midribs light yellow-green (144D) at base. Spots absent; papillae brilliant yellow (11A); nectaries strong yellow-green (144C); pollen greyish reddish brown (200B); stigma deep purple (79C). Fls 240 mm wide, scented; tepals 148 × 85 mm, margins smooth, tips recurved. Lvs 240 × 51 mm, dark green. Stems 1.25–1.4 m, green with darker markings at top, usually twin-flwd.

NOVITA VIII(b-a/b) LO

Trade designation of 'Zanlovita'.

'Nymph' VIII(a/-) OT

Parentage unknown

H: World Breeding BV, pre-2001 **N:** World Breeding BV, 2001 **REG:** World Breeding BV, 2005

Tepals light greenish yellow (5D); inside of inner tepals flushed deep purplish red (59B) at base; base of midveins of outer tepals 59B also. Spots few, papillae present; nectaries green; pollen orange-brown; stigma grey. Tepals long, of narrow to medium width, margins slightly ruffled to ruffled, tips recurved. Lvs of medium length to long and of medium width. Stems tall, green, with few fls. Late season.

'Oberto' VII(a/c-b)

Seedling RM 96-52 × seedling RM 99-73

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of inner tepals deep purplish pink (N66D), shading to yellowish white (155D) towards margins and light greenish yellow (3C) above throat; midveins deep purplish pink (N66C). Inside of outer tepals deep purplish pink (N66C), shading to yellowish white (155D) towards margins and brilliant greenish yellow (3B) above throat; midveins vivid purplish pink (N66B); throat yellowish white (155D). Outside of outer tepals deep purplish pink (70C), shading to strong reddish purple (70B) towards margins and yellowish white (155D) beside midribs; midribs strong yellow-green (144C) at base, shading to light yellow-green (145D) and moderate olive-green (146A) towards top. Outside of inner tepals dark purplish pink (186C), shading to yellowish white (155D) towards margins and at base beside midribs; midribs strong yellow-green (144C) at base and top, shading to light yellow-green (150D) between. Spots and papillae vivid purplish red (N57B); nectaries strong yellow-green (N144C); pollen strong orange (N163B); stigma greenish white (192D). Fls 220 mm wide, scented; tepals 117 × 70 mm, margins ruffled, tips recurved. Lvs 205 × 47 mm. Stems 1.1–1.2 m, green, with up to 5 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Obsession' VII(a/-)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2005

N: Marklily CV, 2005 **REG:** Testcentrum voor

Siergewassen BV, 2007

Tepals greenish white (155C); inside of tepals flushed vivid purplish red (N57A) except at base. Spots absent; papillae present; nectaries green; pollen reddish brown; stigma grey-green. Tepals long and of medium width to broad, margins ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and of medium width to broad. Stems of average height to tall, green with darker markings, with few fls. Mid-season to late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Ohara’ VIII(a-b/-) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** World Breeding BV and Eota/Koster, 2003 **REG:** World Breeding BV, 2005

Tepals light greenish yellow (8B/C); inside of inner tepals 8C towards tips; midribs of inner tepals flushed red. Spots yellow, papillae present, nectaries green; pollen light red-brown; stigma grey-green. Tepals of medium length and width, margins slightly ruffled, tips recurved. Lvs of medium length to long and of medium width to broad. Stems of average height to tall, green, with few to average no. of fls. Mid-season to late. Named in honour of Mrs Wakako Ohara, Headmistress of the Ohara School of Ikebana.

‘Olivia’s Moondance’ I(a/b)

(‘Unique’ × ‘Melissa Jamie’) × Yellow-flowered seedling with very dense spotting at centre of each tepal
H: T.D. Willoughby, 1995 **G:** T.D. Willoughby, 1997
N: T.D. Willoughby, 2005 **I:** J. Harley, 2007

REG: T.D. and S.K. Willoughby, 2006

Tepals pale yellow-green (4D), bleaching to white in sun; midribs brilliant yellow-green (142B); throat vivid yellow (14B). Spots absent; nectaries deep yellowish green (141B); pollen moderate orange (167B); stigma strong yellow-green (142A). Fls 150 mm wide; tepals 100 × 30 mm, margins smooth, tips slightly recurved; in a racemose infl. Lvs 105 × 25 mm, scattered, moderate olive-green (137A). Stems 0.6 m, moderate olive-green (137B), with up to 8 fls. Mid-July (USDA Zone 3, Alberta, Canada). Named after the registrants’ daughter.

‘Onidin’ VIII(a/b) LA

Seedling BG 94-10 × seedling AG 94-01

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals vivid yellow (9A); throat light greenish yellow (1C). Outside of tepals brilliant greenish yellow: outer 6C, inner 7C; all shading to brilliant yellow (7A) towards margins; midribs light yellow-green (144D) at base, strong yellow-green (144A) at top in outer tepals. Spots few, strong purplish red (60B); papillae vivid yellow (9A); nectaries strong yellow-green (144A); pollen moderate reddish orange (N172A); stigma vivid yellow (13A). Buds slightly hairy. Fls 170 mm wide; tepals 105 × 51 mm, margins smooth, tips recurved. Lvs 143 × 25 mm. Stems 1.05 m, green, with c.8 fls.

‘Optimist’ VII(a-b/-)

Parentage unknown

H: Licomba, pre-2007

Tepals greenish white (155C). Spots absent; papillae present; nectaries green; pollen red-brown; stigma grey-green. Fls 250 mm wide; tepals of medium length to long and of medium width to broad, margins ruffled to strongly ruffled, tips recurved to strongly recurved. Lvs of medium length and broad to very broad. Stems 1.1 m, green, with few fls. Mid-season. An application for registration was turned down in 2007, as the cultivar epithet was registered by Licomba in 1985 for another lily (ICNCP, 2004: Art. 27.1). However, as European and Dutch Plant Breeders’ Rights have been granted to this lily under the epithet ‘Optimist’, and a description published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007), this name is established and its use accepted (ICNCP, 2004: Art. 24.4).

‘Orange Cocotte’ I(a/b)

Parentage unknown

H: Kaneko Seeds, Japan, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2006

Inside of tepals, including throat, strong orange (30D). Outside of outer tepals brilliant orange (29A) beside and along midribs, shading to strong orange (30D) towards margins; midribs faintly flushed strong yellow-green (144C) in upper half, and strong yellow-green (143B) at top. Outside of inner tepals strong orange (30D); midribs faintly flushed strong yellow-green (144C). Spots and papillae absent; nectaries strong yellow-green (143C); pollen absent; stigma light orange (28C). Fls 92 mm wide; tepals 54 × 25 mm, margins smooth, tips slightly recurved. Lvs 110 × 15 mm, dark green. Stems 0.9 m, dark with green markings; with c.9 fls.

‘Orange Dazzle’ VIII(a-b/-) LA

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Tepals strong orange: inside of tepals 25A; outside of inner tepals 25B. Spots and papillae absent; nectaries green, pollen orange-brown; stigma orange. Tepals short, of narrow to medium width, margins slightly ruffled, tips slightly recurved. Lvs of medium length and width. Stems of average height to tall, green with darker stripes and spots, with few to average no. of fls. Mid-season.

‘Orange Jewel’ VIII(a-b/-) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Inside of tepals light to pale orange-yellow (16C/D); midveins light orange-yellow (16C) at base. Outside of tepals pale orange-yellow (16D). Spots and papillae present; nectaries green; pollen red-brown; stigma dark purple. Tepals of medium length to long and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length to long, broad. Stems of average

height, green with darker stripes and spots, with few fls. Late to very late.

'Orange Pixels' I(a/b)

'Cha Cha Cha' × 'Brandaris'

H: Laan Flora Facilities BV, pre-2005 **N:** Laan Flora Facilities BV, 2005 **REG:** Laan Flora Facilities BV, 2006

Inside of tepals vivid yellowish orange (30C); throat moderate red (180C). Outside of outer tepals moderate red (181B); margins moderate reddish orange (35A); midribs strong yellow-green (143A) at top. Outside of inner tepals moderate red (180B) shading to moderate reddish orange (N34D) towards margins. Spots numerous, dark red (187A), at base and along margins in lower two-thirds of each tepal; papillae moderate red (180C); nectaries moderate olive-green (146A); pollen moderate reddish brown (175A); stigma dark red (59A). Fls 150 mm wide; tepals 90 × 45 mm, margins smooth, tips slightly recurved. Lvs 130 × 23 mm. Stems 1–1.2 m, green with darker markings at base, with c.8 fls.

'Original Love' VIII(a-b/b) LA

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2003 **N:** Coöp. Kwekersvereniging "The Originals" UA, 2003 **REG:** Coöp. Kwekersvereniging "The Originals" UA, 2006

Tepals moderate red (N34A), paler towards tips; base of midveins dark red. Spots and papillae present; nectaries purple-red; pollen red-brown; stigma purple-red. Fls 200 mm wide; tepals of medium length to short and of narrow to medium width, margins slightly ruffled to ruffled, tips slightly recurved. Lvs of medium length to long and of medium width. Stems 0.9–1.1 m, green with darker stripes and spots, with 4–6 fls. Mid-season.

'Orriolo' VIII(a-b/b) LA

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2005 **N:** Marklily CV, 2005 **REG:** Marklily CV, 2006
Inside of tepals vivid yellow (14A); throat light greenish yellow (7D). Outside of tepals vivid yellow (12A); margins brilliant yellow (13B); base flushed strong yellow-green (144A); midribs of outer tepals vivid yellow (15B) along one side only; midribs of inner tepals brilliant yellow-green (154B). Spots few to absent; papillae vivid yellow (14A); nectaries strong yellow-green (143A); pollen strong orange (N25A); stigma light yellow (18A) edged deep purplish red (71A). Fls 180–200 mm wide; tepals 130 × 50–70 mm, margins smooth to slightly ruffled, tips slightly recurved. Lvs 120–150 × 20 mm, dark green. Stems 1.25–1.7 m, green, with 5–6 fls. Early to mid-season.

'Ovatic' VIII(a-b/b) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007
Inside of tepals yellowish white (155B); outside of inner tepals pale yellow-green (155A). Spots and papillae absent; nectaries green; pollen brown. Tepals

of medium length to long and of medium width, margins ruffled, tips recurved. Lvs of medium length and of medium width to broad. Stems 1.5 m, green with darker markings, with up to 4 fls. Very late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Overlooked' I(a/c)

Parentage unknown

H: A. Burnett, pre-1995 **G:** A. Burnett, pre-1995 **N:** L. Westfall, 1999 **I:** Valley K Greenhouses, 2004 **REG:** Valley K Greenhouses, 2005

Inside of tepals in basal half vivid yellow (14B) under cool conditions, orange under warmer conditions; margins and upper half of tepals strong reddish orange (34C); throat vivid reddish orange (34B). Brushmarks few, on basal half of each tepal, darker orange; spots small, raised, very sparse, towards margins in basal quarter of outer tepals, orange-red; papillae faint; nectaries vivid yellow (14B); filaments orange shading to orange-red at tapered top; pollen strong reddish orange (169A); stigma club-shaped, vivid reddish orange (34A). Buds deep reddish orange. Fls 145 mm wide; tepals 75 × 42 mm, margins smooth, tips recurved. Lvs scattered, 114 × 25 mm. Stems 1 m, bright green, glossy, with 6–10 fls. Mid-July. Nomenclatural standard: colour print supplied by registrant (WSY 0100427).

'Pallada' VIII(a/b) LA

Seedling LARS 93-04 × seedling RS 91-10

H: Vletter en Den Haan Beheer BV, pre-2003 **N:** Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals deep pink (51B) shading to 51C towards margins. Inside of outer tepals strong red (51A) shading to deep pink (51B) towards margins; throat moderate pink (51D). Outside of outer tepals strong pink (54C) shading to deep purplish pink (54B) towards margins; midribs strong yellow-green (143C) at base shading to strong yellow (153D). Outside of inner tepals moderate purplish pink (54D), pale yellow (8D) at base; midribs strong yellow-green (143C) at base shading to strong red (53D). Spots few, deep purplish red (59B); papillae moderate pink (51D); nectaries strong yellow-green (144A); pollen greyish red (176A); stigma strong reddish purple (70B). Buds slightly hairy. Fls 160 mm wide; tepals 93 × 49 mm, margins smooth, tips recurved. Lvs 190 × 22 mm. Stems 1.15–1.5 m, dark with green markings, with c.8 fls.

'Palmares' VIII(b/b) OT

Seedling PG 96-43 × seedling RH 97-09

H: Vletter en Den Haan Beheer BV, pre-2003 **N:** Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals brilliant greenish yellow (3A/B), tips and throat pale greenish yellow (1D and 2D respectively). Outside of outer tepals pale yellow-green (4D); top of midribs light greenish yellow (4B) with strong yellow-green (144A). Outside of inner tepals light greenish yellow (3C); midribs pale yellow-green (4D). Spots absent; papillae brilliant greenish yellow (3A); nectaries strong yellow-green (144A/B); pollen

dark red (183A); stigma dark red (187A). Fls 230 mm wide; tepals 124 × 68 mm, margins slightly ruffled, tips recurved. Lvs 217 × 32 mm, dark green. Stems 1.1–1.4 m, light green, with c.7 fls.

‘Palmira’ VIII(a-b/b-c) OT

Seedling PH 95-39 × AUB

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Inside of tepals light greenish yellow (6D); midvein of each tepal and margins of outer tepals brilliant greenish yellow (6C); throat pale yellow-green (4D). Outside of outer tepals pale yellow-green (4D); margins light greenish yellow (4C); midribs strong yellow-green (143C) at base, uppermost third light yellow-green (150D), top strong yellow-green (143B). Outside of inner tepals light greenish yellow (5D), pale yellow-green (4D) beside midribs; midribs strong yellow-green (143C) at base shading to light yellow-green (150D), and strong yellow (153D) at top. Spots absent; papillae brilliant greenish yellow (6C); nectaries strong yellow-green (143C); pollen dark orange (N163A); stigma greyish purple (N77A). Fls 270 mm wide, scented; tepals 155 × 70 mm, margins of outer tepals ruffled, tips recurved. Lvs 205 × 44 mm. Stems 1.1–1.4 m, green, usually twin-flwd.

‘Pantanal’ I(a/-)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Tepals strong orange: inside N25C, outside of inner tepals 25B. Spots few; papillae inconspicuous; nectaries green; pollen red-brown; stigma orange. Tepals of short to medium length and of narrow to medium width, margins slightly ruffled, tips slightly recurved. Lvs of medium length to long and of narrow to medium width. Stems of average height to tall, green, with few to average no. of fls. Early to mid-season.

‘Papiljo’ VII(a-b/b-c)

Parentage unknown

H: World Breeding BV, pre-2001 **N:** World Breeding BV, 2001 **REG:** World Breeding BV, 2007

Inside of tepals deep purplish pink (N57D) to strong purplish pink (63C), base and midveins brilliant yellow (9C). Outside of inner tepals light purplish pink (73C). Spots yellow; papillae present; nectaries pale green; pollen red-brown; stigma grey-green. Tepals of medium length and width, margins ruffled, tips recurved to strongly recurved. Lvs of medium length to long and of medium width to broad. Stems 1.75 m, green, with up to 10 fls. Late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Parasol’ VII(b/c)

Seedling RM 96-52 × seedling RM 99-43

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals deep purplish pink (N57C), inner tepals shading to 61D towards margins; margins

yellowish white (155D), narrower in outer tepals; throat yellowish white (155D). Outside of tepals deep purplish pink (N57D), shading to yellowish white (155D) towards margins and at base beside midribs; midribs of outer tepals light yellow-green (145B) at base, shading to 150D and strong yellow-green (144B) towards top; midribs of inner tepals strong yellow-green (144C) at base, shading to brilliant yellow-green (150C) and light yellow-green (144D) towards top. Spots and papillae strong purplish red (58B); nectaries strong yellow-green (145A); pollen moderate reddish brown (166B); stigma pale yellow-green (193C). Fls 215 mm wide, scented, becoming darker with age; tepals 110 × 65 mm, margins ruffled, tips recurved. Lvs 200 × 55 mm, dark green. Stems 1–1.2 m, green with darker markings at base, with up to 3 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Pārslīņa’ I(a/b)

‘Saulstarīte’ × seedling 990-11

H: Ā Zorgevics, 1996 **G:** Ā Zorgevics, 2001

N: Ā Zorgevics, 2004 **I:** University of Latvia

Botanical Garden, 2004 **REG:** University of Latvia Botanical Garden, 2007

Inside of tepals pale greenish yellow (2D); throat light yellow-green (145B). Outside of tepals pale yellow flushed brown. Spots few, brown, mostly in the centre of each tepal at the base; nectaries clear; pollen brown; stigma pale brown. Fls 140 mm wide; tepals 80 × 40 mm, margins smooth, tips moderately recurved. Lvs 80 × 10 mm. Stems 1.1 m, green, with up to 15 fls. Late July.

The name means Flake in Latvian.

‘Passionale’ VIII(a-b/-) OT

A temporary name for ‘Garden Pleasure’ prior to registration.

‘Pazzo’ VIII(b-a/c-b) OT

Seedling PG 95-49 × seedling RH 97-09

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals brilliant yellow (10A), shading to light greenish yellow (5D) beside midveins towards tips; midveins vivid yellow (9A); throat pale yellow-green (4D). Outside of outer tepals pale yellow-green (4D), shading to light greenish yellow (6D) towards margins; midribs moderate yellow-green (143D) at base. Outside of inner tepals light yellow (10C), shading to pale yellow-green (4D) beside midribs; midribs moderate yellow-green (143D) at base, shading to light yellow (10C). Spots and papillae vivid yellow (9A); nectaries strong yellow-green (143C); pollen moderate reddish orange (N172A); stigma dark purple (79A). Fls 245 mm wide, scented; tepals 140 × 75 mm, margins slightly ruffled, tips recurved. Lvs 250 × 48 mm. Stems 1.3 m, green with darker markings at base, with up to 4 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Pergamon' I(a/b)

Seedling AJ 98-08 × seedling AJ 95-34

H: Vletter en Den Haan Beheer BV, pre-2004**N:** Vletter en Den Haan Beheer BV, 2004**REG:** Vletter en Den Haan Beheer BV, 2007

Inside of tepals vivid orange (28B); midveins vivid yellowish pink (28A) above nectaries; throat light orange (28D). Outside of outer tepals vivid orange-yellow (23A), shading to strong orange (N25A) towards margins; midribs strong yellow (153D) at base, strong yellow-green (143A) towards top. Outside of inner tepals brilliant orange (29A), shading to light orange (29B) beside midribs; midribs strong orange-yellow (24B). Spots few, dark red (59A); papillae few, vivid yellowish pink (28A); nectaries moderate olive-green (137A); pollen vivid reddish orange (N30B); stigma strong orange (25A). Fls 200 mm wide; tepals 107 × 60 mm, margins smooth, tips slightly recurved. Lvs 170 × 20 mm. Stems 1.1–1.2 m, green with darker markings, especially at base, with up to 6 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Pesaro' VII(b/b)

Syn. of 'Pisaro'.

This lily was registered as 'Pesaro' in 1990 and a description under this cultivar epithet was published in *ILR 1982 Ninth Supplement*. In the same year, Dutch Plant Breeders' Rights were granted to this lily under the cultivar epithet 'Pisaro', which becomes the spelling accepted by the ICRA. In 1993, 'Pisaro' was granted a US Plant Patent under the cultivar epithet 'Pesaro', which remains the accepted name in countries in which 'Pesaro' has legal protection. In *ILR&C 2007*, 'Pesaro' is spelled with an acute accent over the "e". However, this diacritic mark does not appear in either *ILR 1982 Ninth Supplement* or in US Plant Patent documentation.

'Petit Urk' I(a/b)

'Petit Urk' was registered by Mak-Leek Inc. in 2007. Details and a description of 'Petit Urk' were published in *ILR&C 2007*.

'Pink Angel' I(a/b)

Un-named seedling × pink-flowered seedling

H: Bischoff Tulleken Lelies BV, pre-2005 **N:** Bischoff Tulleken Lelies BV, 2005 **REG:** Mak Breeding BV, 2007

Inside of tepals strong purplish pink (68B), shading above nectaries to a yellowish white (155D) ground streaked light orange-yellow (22B); throat yellowish white (155D). Outside of outer tepals moderate purplish pink (186D), shading to deep purplish pink (68A) towards margins; midribs moderate yellow-green (143D) at base, shading to dark greenish yellow (152D) and strong yellow-green (144A) at top. Outside of inner tepals deep purplish pink (68A), tinged pale purplish pink (65C) beside midribs; midribs strong yellow-green (143C) at base, shading to dark greenish yellow (152D). Spots deep purplish red (59B); papillae yellowish white (155D); nectaries strong yellow-green (143A); pollen vivid orange-yellow (23A); stigma pale purplish pink (62D). Fls

140 mm wide; tepals 75 × 45 mm, margins smooth, tips slightly recurved. Lvs 120 × 17 mm. Stems 0.5 m, green with darker markings, with up to 5 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Pinkerton' VI(b/a)

('Fatima' × 'Richard Latty') × unknown

H: B. Mičulka, pre-2005 **N:** B. Mičulka, pre-2005**I:** B. Mičulka, 2006 **REG:** B. Mičulka, 2005

Inside of tepals apricot marbled bright pink; margins pale apricot; throat apricot yellow. Outside of tepals pink; midribs darker. Spots and papillae absent; nectaries green; pollen apricot; stigma dark brown-crimson. Fls a short trumpet, 125 mm wide, 85 mm long; tepals 125 × 44 mm, tips recurved; pedicels 65 mm, lightly spotted brown. Lvs 95 × 10 mm, dull dark green. Stems 0.95 m, stout, infl. racemose. Late June.

'Pink Heaven' VIII(b/a) LO

'White Elegance' × Oriental mixture

H: Testcentrum voor Siergewassen BV, pre-2006**N:** Testcentrum voor Siergewassen BV, 2006**REG:** J.P. van Veen BV, 2006

Inside of inner tepals moderate purplish red (59C); inside of outer tepals deep purplish red (59B) shading to moderate purplish red (59C) towards margins; all margins narrow, very pale purple (76D); with a small spot of yellowish white (155D) at each tip and beside base of midveins of inner tepals; all midveins strong yellow-green (144B); in basal third only of inner tepals, shading to deep purplish red (59B); throat strong yellow-green (144B). Outside of tepals moderate purplish red (186B), with a broad stripe of very pale purple (69C) along midribs, with smaller, darker stripes (69A), and a spot of strong yellow-green (144B) at top. Spots and papillae absent; pollen strong orange (N163B); stigma pale yellow-green (157C). Fls 200 mm wide, scented; tepals 160 × 75 mm, margins smooth, but slightly ruffled towards recurved tips. Lvs 235 × 30 mm. Stems 1.22 m, dark with green markings, with c.3 fls.

'Pink Pride' VII(a/-)

Parentage unknown

H: P.J. de Waard & Zn, pre-1990 **N:** Bloothoofd, 1990 **REG:** P.J. de Waard & Zn, 2007

Inside of tepals moderate purplish pink (74D); margins, tips, midveins and throat yellowish white (155D). Outside of tepals yellowish white (155D) flushed light purple (75B); midribs brilliant yellow-green (150C) shading to strong yellow-green (144C) at top. Spots and papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen strong orange (169C); stigma dark purple (79A). Fls 200 mm wide; tepals 100 × 40 mm, margins smooth, tips strongly recurved. Lvs 160 × 45 mm. Stems 1.05 m, dark, with up to 7 fls.

'Pirandello' VIII(a/b) LA

Seedling BM 94-12 × seedling AM 95-07

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of tepals moderate purplish red (58A) shading to deep purplish red (59B) towards margins; throat pale greenish yellow (1D) with small strong yellow-green (141D) stripe beside nectaries. Outside of tepals moderate purplish red (186B) shading to deep purplish red (71A) towards margins; midribs strong yellow-green (143C) at base shading to strong yellow (153D); midribs of outer tepals strong yellow-green (143B) at top. Spots dark red (59A); papillae absent; nectaries strong yellow-green (143A); pollen moderate reddish brown (177A); stigma light greenish yellow (1C). Fls 190 mm wide, becoming paler with age; tepals 112 × 58 mm, margins smooth, tips slightly recurved. Lvs 200 × 270 mm. Stems 1.1–1.25 m, green with darker markings, with c.5 fls.

'Pleasure' I(a-b/-)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Inside of inner tepals more yellow than vivid yellow (13A), flushed light red at base. Inside of outer tepals more yellow than vivid yellow (14A), flushed light red at base; midveins yellow-green at base. Outside of inner tepals vivid yellow (14A). Spots few; papillae present; nectaries green; pollen brown; stigma purple-red. Tepals short, of narrow to medium width, margins slightly ruffled, tips slightly recurved. Lvs of medium length and of narrow to medium width. Stems of average height, green, with few to average no. of fls. Early.

'Plougana' V(b/a)

Parentage unknown

H: Institut National de la Recherche Agronomique (INRA), pre-1990 **REG:** INRA, 2007

Inside of tepals greenish white (155C) to yellowish white (155D); throat and lower portion of midveins light yellow-green (145B and C respectively). Outside of tepals greenish white (155C); midribs pale yellow-green (155A). Spots and papillae absent; pollen strong orange-yellow (17A); stigma pale yellow-green (157C). Fls 180 mm wide; tepals 165 × 45 mm, margins smooth to slightly ruffled, tips recurved. Lvs 200 × 28 mm. Stems 1.05 m, pale green, with up to 4 fls.

'Ploumidia' V(b/a)

Parentage unknown

H: Institut National de la Recherche Agronomique (INRA), pre-1988 **REG:** INRA, 2007

Tepals, including throat, yellowish white (155D); midveins pale green. Spots and papillae absent; pollen orange; stigma pale green. Fls of average diameter, strongly scented; tepals long to very long and of medium width to broad, margins slightly ruffled, tips recurved. Lvs very long and narrow. Stems of average height, green.

'Plouzarel' V(b/a)

Parentage unknown

H: Institut National de la Recherche Agronomique (INRA), pre-1990 **REG:** INRA, 2007

Tepals yellowish white (155B); midveins brilliant yellow-green (149B). Spots and papillae absent; nectaries pale yellow-green (155A); pollen orange; stigma grey-blue. Tepals long and of narrow to medium width, margins slightly ruffled, tips strongly recurved. Lvs of medium length and of narrow to medium width. Stems of average height to tall, green.

'Pomero' VII(a/b-c)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals strong purplish pink (73B); narrow margins and large spot above nectaries, yellowish white (155D). Inside of outer tepals strong purplish pink (68B); margins, above nectaries, and throat, yellowish white (155D). Outside of outer tepals dark purplish pink (186C), margins yellowish white (155D); midribs moderate yellow-green (148A) at base shading to moderate purplish red (186A), and 186B with moderate yellow-green (148A) at top. Outside of inner tepals moderate purplish pink (62B), margins yellowish white (155D); midribs dark red (59A) at base with a little strong yellow-green (144C), shading to moderate purplish red (186B), and yellowish white (155D) at top. Spots absent; papillae yellowish white (155D) and strong purplish pink (73B); nectaries light yellow-green (145B); pollen dark orange (N163A); stigma yellowish white (156D). Fls 250 mm wide; tepals 137 × 65 mm, margins ruffled, tips recurved. Lvs 235 × 55 mm. Stems 1.3 m, dark, with c.9 fls.

'Pordenone' I(a/b)

Seedling AW 95-05 × seedling AM 96-17

H: Vletter en Den Haan Beheer BV, pre-2004**N:** Vletter en Den Haan Beheer BV, 2004**REG:** Vletter en Den Haan Beheer BV, 2007

Tepals yellowish white (155D), midribs light yellow-green (144D); some tepals flushed strong purplish red (64B) towards tips; throat yellowish white (155D). Spots few, deep purplish red (59B); papillae absent; nectaries strong yellow-green (143B); pollen brownish orange (N167B); stigma moderate purplish red (59C). Fls 183 mm wide; tepals 95 × 50 mm, margins smooth, tips recurved. Lvs 195 × 17 mm. Stems 1.3 m, dark, with up to 5 fls.

Published in *Registraties week 44-2006 t/m 45-2007*, KAVB (2007).

'Portovenere' VIII(b-a/b) OT

Seedling PG 96-43 × seedling 93-78

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2007

Inside of inner tepals brilliant greenish yellow (6C); midveins 6B; throat pale yellow-green (4D). Inside of outer tepals light greenish yellow (8B), yellowish white (155D) at base beside nectaries; midveins brilliant yellow (11A). Outside of outer tepals pale yellow-green

(4D), shading to brilliant greenish yellow (5A) towards margins; midribs and area beside midribs flushed brilliant yellow (11A); base and top of midribs strong yellow-green (144C and 143C respectively). Outside of inner tepals light greenish yellow (4C), shading to 6D towards margins and pale yellow-green (4D) beside midribs; midribs light yellow-green (144D) at base shading to pale yellow-green (4D). Spots and papillae brilliant greenish yellow (6B); nectaries strong yellow-green (144C); pollen dark red (187A); stigma light purplish grey (N187D) tinged purplish grey (N187C). Fls 190 mm wide, scented; tepals 110 × 63 mm, margins ruffled, tips strongly recurved. Lvs 244 × 38 mm. Stems 1.2–1.3 m, green, with up to 7 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Prairie Jewel' I(b-a/b-c)

Parentage unknown

H: F. Fellner, 1997 **G:** F. Fellner, 1999 **N:** L. Westfall, 2004 **I:** Valley K Greenhouses, 2006 **REG:** Valley K Greenhouses, 2005

Inside of tepals vivid red (45B); base with a narrow patch of white under hot conditions; small spot of white at tips; throat white, heavily frosted. Outside of tepals greyish purple (N187B). Spots deep red (185A), small, fairly numerous, mostly over basal third of each tepal; papillae and nectaries white; pollen strong orange (168A); stigma red. Fls 137 mm wide, colour rarely fading; tepals 58 × 37 mm, margins smooth, tips slightly recurved; pedicels scurfy; with secondary buds, all buds with short, bristly hairs. Lvs scattered, 134 × 11 mm. Stems 0.8 m, bright green, with c.17 or more fls. Late July. Resistant to botrytis. Nomenclatural standard: colour print supplied by registrant (WSY 0100428).

'Precious Peggy' I(b/c)

Sutter's Gold Group × 'Unique'

H: S. & N. Dunville, 1993 **G:** S. & N. Dunville, 1995 **N:** N. Dunville, 2000 **I:** S. & N. Dunville, 2005 **REG:** S. & N. Dunville, 2005

Inside of tepals shading from whitish pink in basal half to moderate purplish pink (68C) in upper third; small spot of strong purplish pink (68B) at tips; base and throat white. Outside of tepals strong purplish pink (68B); outer tepals shading to yellowish green beside greenish midribs; midribs of inner tepals more prominent, light green. Spots strong red (46A), regularly distributed over basal third of each tepal and throat, fairly large, becoming progressively smaller towards nectaries; papillae whitish and moderate purplish pink (68C); nectaries strong purplish pink (68B); pollen moderate orange (167B); style short, stout, pale apple green; stigma reddish brown. Fls c.130 mm wide; inner tepals 72 × 37 mm, outer tepals 71 × 25 mm, margins smooth, tips slightly recurved. Lvs scattered, 116 × 11 mm. Stems 1.36 m, mid-green with brown markings, with c.15 fls. Mid-July.

'Presidente' VII(a/b-c)

(Un-named seedling × (*L.* × *parkmanii* 'Allegra' × 'Star Gazer')) × (Un-named seedling × ('White Delight' × 'Star Gazer'))

H: Bischoff Tulleken Lelies BV, pre-2001 **N:** Bischoff Tulleken Lelies BV, 2001 **REG:** Mak Breeding BV, 2006

Inside of tepals deep purplish pink (70C); throat and flare above nectaries yellowish white (155D). Outside of outer tepals moderate purplish pink (186D) with dark purplish pink (186C) stripes; yellowish white (155D) at base beside midribs; midribs strong yellow-green (143A) at base and top, shading to light yellow-green (145D) and strong yellow (153D) between. Basal third of outside of inner tepals yellowish white (155D), upper two-thirds moderate purplish pink (186D); midribs strong yellow-green (143A) at base shading to light yellow-green to pale greenish yellow (145D/2D) flushed dark purplish pink (186C), and strong yellow-green (144B) at top. Spots absent; papillae yellowish white (155D) and deep purplish pink to strong reddish purple (70C/B); nectaries light yellow-green (145B); pollen dark orange (N163A); stigma light grey (188D). Fls 225 mm wide, scented; tepals 135 × 75 mm, margins slightly ruffled, tips recurved. Lvs 125 × 48 mm. Stems 1–1.1 m, green with darker markings, with c.3 fls.

'Pretty Pearl' VII(a/b-c)

Seedling 9-13-SP × 'Kissproof'

N: Rooyakkers Lelie BV, 2007 **REG:** Rooyakkers Lelie BV, 2007

Inside of inner tepals strong purplish red (64C), shading to yellowish white (155D) towards margins; midveins strong purplish red (64B); throat yellowish white (155D). Inside of outer tepals strong purplish red (64C); margins towards tips and base beside nectaries yellowish white (155D); midveins strong purplish red (64C) with light yellow-green (145B). Outside of outer tepals deep purplish pink (68A) and yellowish white (155D); midribs yellowish white (155D) at base shading to light yellow-green (145D). Outside of inner tepals deep purplish pink (68A) and yellowish white (155D); margins towards tips, and up to half of midribs, yellowish white (155D). Spots moderate purplish red (58A); papillae absent; nectaries light yellow-green (154D). Fls double, 180 mm wide; tepals 105 × 36 mm, margins ruffled, tips slightly recurved. Lvs 250 × 40 mm. Stems 0.8–1.35 m, green, with up to 4 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Priedula' VI(b/a)

Seedling 169-S-F × 'Atvasara'

H: Ā. Zorgevics, 1987 **G:** Ā. Zorgevics, 1992 **N:** Ā. Zorgevics, 2006 **I:** University of Latvia Botanical Garden, 2006 **REG:** University of Latvia Botanical Garden, 2006

Inside of tepals flushed moderate yellowish pink (33D), particularly towards margins and beside midveins; base and throat brilliant yellow (20A). Outside of tepals light violet; midribs darker, shading to white towards margins. Spots and papillae absent; pollen brown; stigma brownish green. Fls 150 mm wide, strongly scented, long-lasting; tepals 110 × 40 mm, margins almost smooth, tips strongly recurved. Lvs scattered, 110 × 7 mm, glossy. Stems 1.3 m, violet. Bulbs purple. Published in *Lilium Balticum. Latvijas liliju*

selektionāru izaudzētās liliņas, No. 1 (1995).

Nomenclatural standard: colour print supplied by registrant (WSY 0100425).

‘Prime Ice’ V(b/a)

Parentage unknown

H: J.W.A. van der Wereld BV, pre-2003 **N:** Liliande BV, 2003 **REG:** J.W.A. van der Wereld BV, 2007
Tepals greenish white (155C); midveins green at base. Spots and papillae absent; nectaries green; pollen orange-yellow; stigma white. Tepals long and of medium width, margins (very) slightly ruffled, tips recurved. Lvs of medium length to long and of medium width. Stems 2 m, green, with up to 7 small fls. Very late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007). ‘Rime Ice’ was used as a temporary name prior to registration.

‘Private Dancer’ VII(a-b/b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007

Inside of inner tepals strong purplish red (60C/D), white beside nectaries. Inside of outer tepals strong purplish red (60D) to deep purplish pink (67C). Outside of inner tepals strong reddish purple (72C) to strong purplish red (60D). Spots and papillae present; nectaries green; pollen orange-brown; stigma green, speckled. Fls of small to average dimensions; tepals of short to medium length and of medium width to broad, margins slightly ruffled to ruffled, tips recurved to strongly recurved. Lvs of medium length and of medium width to broad. Stems 1.25 m, green with darker spots and stripes, with up to 4 fls. Mid-season to late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Promenade’ VIII(a/b) LA

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007

Inside of tepals vivid red (45A), shading to strong reddish orange (169A) towards tips. Outside of inner tepals strong reddish orange (169A) to strong orange (169B), colouring more intense towards margins. Spots absent; papillae inconspicuous; nectaries green; pollen red-brown; stigma purple-red. Tepals of short to medium length and of narrow to medium width, margins (very) slightly ruffled, tips slightly recurved to recurved. Lvs of medium length and of narrow to medium width. Stems of average height to tall, green with some darker spots and stripes, with average no. of fls. Early to mid-season.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Prorosa’ I(a/b)

Un-named ‘Peachblush’ × ‘Carol Jean’ seedling × pink-flowered seedling

H: Bischoff Tulleken Lelies BV, pre-2002 **N:** Bischoff Tulleken Lelies BV, 2002 **REG:** Mak Breeding BV, 2006

Inside of inner tepals light purplish pink (63D), strong purplish pink (63C) above throat with a small yellowish white (155D) stripe. Inside of outer tepals strong purplish pink (63C), light purplish pink (63D) at top and a small yellowish white (155D) stripe above throat; throat pale yellow (19C). Outside of outer tepals pale purplish pink (65C), margins moderate purplish pink (65A); midribs yellowish white (155D) shading to light yellow-green (144D), and strong yellow-green (144C) at top. Outside of inner tepals light purplish pink (65B) shading to moderate purplish pink (65A) towards margins. Spots and papillae absent; nectaries moderate olive-green (137A); pollen dark orange (N163A); stigma light yellow (21D). Fls 160 mm wide; tepals 85 × 50 mm, margins smooth, tips very slightly recurved. Lvs 125 × 24 mm. Stems 0.75 m, dark with a little green marking, with c.8 fls.

‘Puerto Cortes’ I(a-b/-)

Parentage unknown

H: Fa Jaap Boon, pre-1999 **N:** Trior Lelie BV/Koninklijke Van Zanten BV, 1999 **REG:** Fa Jaap Boon, 2007

Tepals more yellow than vivid reddish orange (32A); midveins orange at base. Spots and papillae present; nectaries orange; pollen dark brown; stigma orange. Tepals short to very short and narrow, margins slightly ruffled, tips slightly recurved to recurved; infl. very hairy. Lvs of medium length and of narrow to medium width. Stems of average height, green with darker spots and stripes, with few to average no. of fls. Very early.

‘Pulse’ VII(a/b)

Un-named seedling × un-named seedling

H: Sande BV, pre-2005 **N:** Sande BV, 2005 **REG:** Sande BV, 2007

Inside of tepals vivid purplish red (67B) to deep purplish pink (67C), becoming more blue with age; paler towards margins, tips and base; midveins deep red (60A) at base. Outside of inner tepals deep purplish pink (N66D). Spots few to average no.; papillae present; nectaries green edged yellow; pollen reddish brown; stigma purple-red with small spots. Fls 180 mm wide; tepals of medium length to long and of medium width, margins ruffled, tips slightly recurved to recurved. Lvs of medium length and broad. Stems 1 m, green, with c. 6 fls. Late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Puma’ VIII(a/b) LA

Mixture of Asiatic tetraploid hybrids × mixture of Div. VIII LA hybrids

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies Holland BV, 2006 **REG:** De Jong Lelies Holland BV, 2007

Inside of tepals strong orange (25B); narrow margins, tips and basal half of midveins N25A; throat light orange-yellow (25D). Outside of tepals strong orange-yellow (24B), shading to strong orange (25A) towards margins; midribs strong yellow-green (144C) at base. Spots and papillae absent; nectaries light olive (152A); pollen greyish reddish brown (200B); stigma deep purplish red (59B). Fls 195 mm wide; tepals 120 × 55 mm, margins smooth, tips recurved. Lvs 205 ×

33 mm. Stems 1–1.2 m, green with darker markings, with up to 3 fls.
Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Purple Diamond' VIII(a/b) LA

'Amarone' × seedling LA 93012-IT

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies Holland BV, 2006 **REG:** De Jong Lelies Holland BV, 2007

Inside of tepals strong purplish red (60B); margins deep red (60A); throat strong purplish pink (63C), strong yellow-green (144A) beside nectaries. Outside of tepals strong purplish red (60C); margins 60B; light purplish pink (63D) towards base beside midribs; tips and base tinged strong yellow-green (144A). Spots and papillae absent; nectaries strong yellow-green (144A); pollen moderate orange (167A); stigma dark purplish red (N79A). Fls 230 mm wide; tepals 120 × 40–60 mm, margins smooth, tips recurved to strongly recurved. Lvs 190 × 30 mm, dark green, glossy. Stems 1.1 m, green with darker markings, with up to 3 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Purple Prince' VIII(a-b/a-b) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007

Tepals moderate purplish red (64A) to strong purplish red (64B); inside of tepals paler towards base; midveins pale purple-red at base. Spots present; papillae inconspicuous; nectaries green; pollen orange-brown; stigma dark purple. Tepals of medium length and width, margins slightly ruffled to ruffled, tips slightly recurved. Lvs of medium length and of medium width to broad. Stems 1.45 m, green with some darker spots and stripes, with up to 5 fls. Very late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Pyroman' I(b-c/d)

Parentage unknown

H: B. Mičulka, pre-2007 **G:** B. Mičulka, pre-2007
N: B. Mičulka, pre-2007 **I:** B. Mičulka, pre-2007
REG: B. Mičulka, 2007

Inside of tepals deep orange-red, tinted crimson in centre; throat paler orange. Spots numerous, large, black, over three-fifths of each tepal; papillae present in throat; nectaries white, frosted; pollen and stigma dark brown. Fls 120 mm wide; tepals 80 × 30 mm, margins ruffled, tips strongly recurved; pedicels 75 mm, with dark brown spots. Lvs 115 × 11 mm. Stems 1.05 m, stiff, slightly hairy, infl. short. Early July.

'Queen Mother' VII(a/b)

('Star Gazer' × un-named seedling) × un-named seedling

H: Bischoff Tulleken Lelies BV, pre-2001 **N:** Bischoff Tulleken Lelies BV, 2001 **REG:** Mak Breeding BV, 2006

Inside of tepals deep purplish pink (N57D); up to half the length of margins yellowish white (155D) in inner tepals and in upper third of outer; midveins

vivid purplish red (N57B); throat, and along nectaries of outer tepals, yellowish white (155D). Outside of outer tepals dark purplish pink (186C), with some yellowish white (155D) at base; midribs strong yellow-green (144B) at base shading to yellowish white (155D), and strong yellow-green (144A) with 144B at top. Outside of inner tepals light purplish pink (63D) shading to strong purplish pink (63C) towards tips, very pale purple (69D) in basal third, some yellowish white (155D) at base beside midribs; midribs strong yellow-green (144C) at base and top, shading to brilliant yellow-green (154C) between. Spots and papillae deep purplish red (61A); nectaries light yellow-green (144D); pollen strong reddish orange (169A); stigma very pale green (130D). Fls 230 mm wide; tepals 135 × 70 mm, margins smooth, tips recurved. Lvs 155 × 40 mm. Stems 1.1–1.3 m, green, with darker markings, with c.6 fls.

'Radar Love' I(a-)

Parentage unknown

H: P. Hoff, 1987 **G:** Hoffgaarde BV, 1989
N: Hoffgaarde BV, 1989 **I:** Hoffgaarde BV, pre-1993
REG: Hoffgaarde BV, 2007

Inside of tepals brilliant yellow (11A) with small vivid orange-yellow (21A) stripes, strong yellowish pink (37A) stripes along margins; midveins vivid orange-yellow (21A), up to half flushed strong yellowish pink (37A); throat pale yellowish pink (29D). Outside of tepals pale yellow (20C), flushed pale purplish pink (65D) towards margins and midribs (65C/D); margins vivid orange-yellow (21A), fading towards the tip; base strong yellow-green (144B). Spots and papillae dark red (187A); nectaries pale orange-yellow (24D) edged deep yellowish pink (39B); pollen deep red (60A); stigma vivid reddish orange (34B). Fls 170 mm wide; tepals 90 × 40 mm, margins slightly ruffled, tips slightly recurved. Lvs 90 × 10 mm, moderate olive-green (137A). Stems 0.65 m, pale green with darker markings, with up to 11 fls.

This lily might have been distributed as 'Romance' prior to registration. An application to register this lily as 'Romance' was turned down in July 1993, as the cultivar epithet was registered by K. Braun in 1986 for another lily (ICNCP, 2004: Art. 27.1). Furthermore, this lily was granted Dutch Plant Breeders' Rights in October 1993 under the name 'Radar Love'.

'Raisance' VIII(a-b/a) LA

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007

Tepals strong purplish red: inside and outside of inner tepals 60C, inside of outer tepals 60B/C. Spots and papillae absent; nectaries green; pollen orange-brown; stigma orange. Tepals of medium length and width, margins (very) slightly ruffled, tips slightly recurved to recurved. Lvs long and of narrow to medium width. Stems 1.15 m, green with some darker markings, with up to 4 small fls. Mid-season. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

RED RUM VII(a-b/b)
Trade designation of 'Zanlorum'.

'Red Sensation' I(a-b/-)
Parentage unknown

REG: M.A. van den Berg en Zn, 2006
Inside of tepals vivid red (44A) towards tips, shading to 45A at base; midveins orange-red at base. Outside of tepals vivid red to vivid reddish orange (44A/B), vivid to strong red (45A/46A) at base. Spots and papillae present; nectaries green, becoming red; pollen red-brown; stigma purple. Tepals short to very short, narrow, margins smooth to very slightly ruffled, tips slightly recurved. Lvs of medium length and of narrow to medium width. Stems of average height to tall, green, with darker spots and stripes, with few to average no. of fls. Early.

'Revolution' VIII(a/-) OT
Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005
Inside of inner tepals vivid yellow (9B) shading to brilliant yellow (9C) towards tips. Inside of outer tepals brilliant yellow (9C). Outside of inner tepals brilliant yellow to pale greenish yellow (9C/D). Spots absent; papillae present; nectaries green; pollen orange-brown; stigma green flushed red. Tepals of medium length to long and of medium width, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length to long, broad. Stems of average height to tall, green. Mid-season to late.

'Rhodos' (Bischoff Tulleken) VII(a-b/-)
Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-1992
Inside of inner tepals light purplish pink (65B); margins yellowish white (155B). Inside of outer tepals moderate purplish pink (65A); margins yellowish white (155B). Outside of tepals yellowish white (155B). Spots red; papillae present; nectaries green; pollen dark brown; stigma purple. Tepals of medium length and of narrow to medium width, margins ruffled to strongly ruffled, tips strongly recurved. Lvs of medium length and width. Stems of average height, green with darker markings, with average no. of fls. An application for registration was turned down in 2007, as the cultivar epithet was registered by V. Strasser in 1986 for another lily (ICNCP, 2004: Art. 27.1). However, as Dutch Plant Breeders' Rights were granted to this lily under the epithet 'Rhodos' in 1992, it remains the accepted name (ICNCP, 2004: Art. 28.2).

'Rhodos' (Van der Salm) I(a/b)
Un-named seedling × un-named seedling

H: G.C. Van der Salm, pre-1983
Tepals vivid red (44A), darker towards margins and base. Pollen orange-brown. Fls 140–150 mm wide; tips recurved; pedicels 89–114 mm long, reddish. Lvs 60–90 × 13 mm. Stems 0.7–0.9 m, with up to 9 fls in a well-spaced pyramidal raceme. As this lily was granted a US Plant Patent under the cultivar epithet 'Rhodos' in 1986, this epithet remains the accepted name (ICNCP, 2004: Art. 28.2).

'Rodilana' I(a/b)

Seedling 3646-RO × seedling AZ TC008

H: De Jong Lelies Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2006

Inside of tepals vivid reddish orange (34A); inner tepals flushed vivid red (45B) towards margins, along midveins and above throat; outer tepals shading to vivid red (46B) towards margins; throat strong reddish orange (31A). Outside of outer tepals moderate red (180A) shading to vivid reddish orange (34A) towards margins; midribs light olive (152B) at base. Outside of inner tepals vivid reddish orange (34B); midribs dark red (183A). Spots and papillae absent; nectaries dark yellowish green (139A); pollen greyish to dark red (178A/183A); stigma dark red (59A). Fls 165 mm wide; tepals 85 × 47 mm, margins smooth, tips very slightly recurved to straight. Lvs 135 × 17 mm. Stems 0.95–1.1 m, dark, with c.8 fls.

'Romance' (Hoff) I(a/-)

'Radar Love' might have been distributed as 'Romance' prior to registration.

'Romero Star' VII(b/b)

Parentage unknown

H: A.C. Imanse, pre-1991 **REG:** A.C. Imanse, 2007
Inside of tepals and top of throat moderate purplish pink (62B); margins, tips, base and most of throat yellowish white (155D); midveins flushed strong red (53C/D). Outside of outer tepals pale purplish pink (62D); margins, tips and base yellowish white (155D); midribs light purplish pink (62C). Outside of inner tepals greenish white (155C); midribs spotted pale yellow (11C) and flushed light purplish pink (62C). Spots numerous, dark red (59A); papillae present; nectaries strong yellow-green (145A); pollen brownish orange (171B); stigma deep reddish purple (77A). Fls 230 mm wide; tepals 125 × 65 mm, margins strongly ruffled, tips strongly recurved. Lvs 200 × 48 mm. Stems 1.3 m, green with some darker markings at base, with up to 6 fls.

'Ronaldo' VII(a/b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007
Inside of inner tepals moderate purplish red (64A), inside of outer tepals strong purplish red (60B); tips and base paler; midveins yellow at base. Outside of inner tepals strong purplish red (64B/C). Spots and papillae present; nectaries pale green; pollen red-brown; stigma grey-green. Tepals of medium length to long and of medium width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad. Stems of average height to tall, green with some darker spots and stripes, with few to average no. of fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Rosande' I(a/b)

Pink-flowered seedling × pink-flowered seedling
H: Bischoff Tulleken Lelies BV, pre-2004 **N:** Bischoff Tulleken Lelies BV, 2004 **REG:** Mak Breeding BV, 2006

Inside of tepals deep purplish pink (N57D), margins closely spotted strong purplish red (64C), a spot of pale purplish pink (65D) above throat; throat pale yellow-green (193D). Outside of outer tepals light yellow-green (145C) with strong purplish red (59D), shading to 59D towards margins; tips and midribs strong yellow-green (144A and 144C respectively). Outside of inner tepals deep purplish pink (58D) shading to N57C towards margins; midribs strong yellow-green (144B). Spots and papillae absent; nectaries strong yellow-green (143B); pollen vivid orange-yellow (23A); stigma with strong reddish purple (70B) stripe on moderate yellow-green (143D) ground. Fls 180 mm wide; tepals 103 × 45 mm, margins smooth, tips slightly recurved. Lvs 145 × 20 mm. Stems 0.63–0.8 m, green, with c.6 fls.

'Roxette' VIII(a/b) LA

Seedling BW 96-19 × seedling AW 95-07
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006

Tepals, including throat, yellowish white (155D); outside of outer tepals with a broad, light yellow-green (144D) stripe beside and along midribs; midribs strong yellow-green (143C) at base and top; midribs of inner tepals 143C throughout. Spots very sparse, strong purplish red (60B); papillae yellowish white (155D); nectaries strong yellow-green (143C); pollen moderate reddish orange (171A); stigma strong yellow (N144B). Fls 195 mm wide, buds and young fls flushed reddish; tepals 105 × 60 mm, margins smooth, tips slightly recurved. Lvs 130 × 18 mm, dark green. Stems 0.9–1.3 m, dark, with green markings, with c.5 fls.

'Rozapet' VI(b-c/a)

Parentage unknown
H: B. Mičulka, pre-2006 **N:** B. Mičulka, pre-2006 **I:** B. Mičulka, 2006 **REG:** B. Mičulka, 2006

Inside of tepals deep pink, with pale yellow rays in throat. Outside of tepals darker pink. Spots absent; nectaries pale olive-green; pollen apricot yellow; stigma greyish. Fls 125 mm wide; tepals 110 × 38 mm, tips recurved; pedicels 90 mm long, stout, spotted. Lvs 100 × 10 mm, dull dark green. Stems 1 m, stout, greyish and slightly spotted, infl. short. Early July.

'Sabaneta' VIII(a-b/b) OT

Parentage unknown
H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007
 Tepals light purplish pink (55C) on pale greenish yellow (10D) ground; inside of tepals paler towards margins and tips; midveins light greenish yellow (5C) at base, shading to purple-red. Spots and papillae present; nectaries green; pollen red-brown; stigma purple, speckled. Tepals of short to medium length and of medium width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium

length to long and of medium width to broad. Stems 1.3 m, green, with average no. of fls. Very early to early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Sabrosa' VII(b-a/b)

'Gentile' × 'Willeke Alberti'
H: De Jong Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2007

Tepals dark purplish pink (186C), shading to moderate purplish pink (186D) towards margins; base pinkish white (N155B); inside of tepals with narrow margins of pinkish white (N155B); midribs N155B. Spots absent; papillae pinkish white (N155B); nectaries strong yellow-green (144A); pollen vivid orange (28B); stigma light yellowish green (135D/136D). Fls 200 mm wide; tepals 130 × 75 mm, margins ruffled, tips recurved. Lvs 205 × 70 mm, dark green. Stems 1–1.15 m, green with darker markings, with up to 4 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Salinas' VII(a/b)

Parentage unknown
H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Inside of inner tepals deep red (53A/60A), paler towards tips. Inside of outer tepals deep red (60A); midveins more reddish and deep to strong red (53A/46A) at base. Outside of inner tepals strong purplish red (60C/D). Spots and papillae present; nectaries green, becoming yellow; pollen orange-brown; stigma purple. Tepals of medium length and of narrow to medium width, margins ruffled, tips slightly recurved to recurved. Lvs of medium length to long, broad. Stems of average height to tall, green, with darker stripes and spots, with few fls. Mid-season to late.

'San Bernardo' VII(b-a/b)

Seedling RW 96-05 × seedling RW 99-06
H: Vletter en Den Haan Beheer BV, pre-2004
N: Vletter en Den Haan Beheer BV, 2004
REG: Vletter en Den Haan Beheer BV, 2007

Tepals, including throat, yellowish white (155D); midveins flushed pale greenish yellow (2D); midribs of outer tepals strong yellow-green (N144C) at base, shading to yellowish white (155D); light yellow-green (145D) tinged strong yellow-green (144A) at top; midribs of inner tepals light yellow-green (144D) at base and top, shading to pale greenish yellow (2D) between. Spots absent; papillae few, yellowish white (155D); nectaries strong yellow-green (N144D); pollen strong reddish orange (169A); stigma very pale green (189D). Fls 215 mm wide, scented; tepals 120 × 67 mm, margins ruffled, tips recurved. Lvs 170 × 52 mm. Stems 1.2–1.3 m, green with darker markings at base, with up to 5 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Santhia' VIII(a-b/b) OT

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of tepals light greenish yellow (5D), midveins and above nectaries of inner tepals, 5C; shading to pale yellow-green (155A) towards margins of outer tepals and beside top of midveins of inner tepals. Outside of outer tepals pale yellow-green (155A), light yellow-green (145D) beside midribs; midribs strong yellow-green (144B) at base and top, shading to 144C between. Outside of inner tepals pale yellow-green (4D), margins closely spotted light greenish yellow (5D); pale yellow-green (155A) beside midribs; midribs strong yellow-green (144C) at base and top, shading to light yellow-green (145C) between. Spots absent; papillae light greenish yellow (5C); nectaries strong yellow-green (144C); pollen brownish orange to dark reddish orange (172C/B); stigma dark greyish purple (N92A). Fls 215 mm wide, slightly scented; tepals 150 × 75 mm, margins slightly ruffled, tips recurved. Lvs 220 × 57 mm. Stems 1.1–1.3 m, green, with darker markings at base and top, usually twin-flwr.

'San Vito' VIII(a/b-c) LA

Seedling BW 96-14 × seedling AJ 96-27

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

All tepals pale yellow (11D), with pale yellowish pink (27D) margins; outside of tepals sometimes more pinkish red; throat yellowish white (N155D). Outside of outer tepals with small stripes of deep pink (48A) at base; midribs strong yellow-green (143C), brownish orange (164A) at top; outside of inner tepals closely spotted deep purplish red (59B) at base. Spots very sparse, dark red (59A); papillae absent; nectaries strong yellow-green (N144D); pollen moderate reddish brown (175A) flushed strong reddish orange (169A); stigma light orange-yellow (22D). Fls 210 mm wide, predominantly yellowish white (155D) under warm and bright conditions; tepals 120 × 65 mm, margins smooth, tips recurved. Lvs 175 × 20 mm. Stems 1.1–1.5 m, dark, with green markings at top, with *c.* 4 fls.

'Satin Sheets' Div.?(b-c/c)

Parentage unknown

H: F. Gartleman, 1988 **G:** F. Gartleman, 1992**N:** W. Nicholls, 2000 **REG:** Memorial University of Newfoundland Botanical Garden, 2005

Inside of tepals shading from dark red (187A) at base to deep red (53A) at tips; throat greyish purple to dark red (N77A/187A). Outside of tepals deep purplish red (59B). Spots sparse, evenly distributed over basal third of each tepal, and papillae, in throat only, all dark red (187A); nectaries greyish purple (N77A); pollen dark red (183A); stigma deep red (53A). Fls 125 mm wide, with a satin sheen; tepals 73 × 33 mm, margins smooth, tips recurved. Lvs scattered, 113 × 12 mm. Stems 1.3 m, dark red (187A), with *c.* 15 fls. July to late Aug. Nomenclatural standard: colour print supplied by registrant (WSY 0100426).

'Satisfaction' VIII(a-b/-) OT

Parentage unknown

H: World Breeding BV, pre-2001 **N:** World Breeding BV, 2001 **REG:** World Breeding BV, 2005

Inside of inner tepals between moderate red (185B) and more red than strong red (53B), paler towards tips; becoming lighter brown-purple with age. Inside of outer tepals strong red (53B); midveins brilliant greenish yellow (6C) at base. Outside of tepals deep pink (185D). Spots and papillae absent; nectaries green; pollen red-brown; stigma purple. Tepals of medium length to long and of medium width to broad, margins ruffled, tips recurved. Lvs of medium length to long and of medium width. Stems tall, green with darker stripes and spots, with few fls. Late.

'Scalini' VII(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2004**N:** Vletter en Den Haan Beheer BV, 2004**REG:** Vletter en Den Haan Beheer BV, 2007

Inside of tepals deep purplish pink (N74C) to moderate purplish red (186B); margins very pale purple (76D) on inner tepals, narrow and greenish white (155C) on outer tepals; midveins above nectaries strong reddish orange (31A) to light yellow (20B), tinged vivid reddish purple (N74B) at top; throat light orange-yellow (16C) to pale orange-yellow (16D). Outside of outer tepals vivid reddish purple (N74B) to deep purplish pink (N74C) on paler ground, shading to pale yellow-green (155A) beside midribs at base and at margins; midribs pale greenish yellow (1D) at base shading to brilliant yellow-green (154C) towards top; top strong yellow-green (143B). Outside of inner tepals deep purplish pink (N74C) on paler ground, shading to pale yellow-green (155A) beside midribs at base; margins greenish white (155C); midribs light yellow-green (154D) shading to vivid yellow-green (154A) towards top. Spots dark red (187C); papillae greenish white (155C); nectaries strong yellow-green (144B); pollen strong brown (172A); stigma dark purplish red (N79B). Fls 206 mm wide; tepals 127 × 66 mm, margins ruffled, tips slightly recurved. Lvs 265 × 110 mm, dark green. Stems 1.1 m, pale green, with up to 3 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Sefanja' I(a/b)

('Grand Prix' × ('Yellow Blaze' × 'Connecticut King')) × short, red/orange-flowered seedling

H: Bischoff Tulleken Lelies BV, pre-2004 **N:** Bischoff Tulleken Lelies BV, 2004 **REG:** Mak Breeding BV, 2006

Inside of tepals vivid yellowish pink (28A) shading to vivid yellowish orange (30C) towards margins; throat green with darker markings. Outside of outer tepals strong orange (N25B) shading to N25A towards margins; midribs strong yellow-green at base and top (143C and 143A respectively), shading to light yellow-green (145C) between. Outside of inner tepals vivid orange (28B) shading to vivid yellowish pink (28A) towards margins; midribs strong yellow-green (143C). Spots and papillae absent; nectaries moderate

olive-green (137A); pollen strong reddish orange (169A); stigma strong purplish red (63A). Fls 175 mm wide; tepals 106 × 49 mm, margins smooth, tips recurved. Lvs 155 × 20 mm, dark green. Stems 0.55–0.65 m, green with darker markings, with c.5 fls.

‘Selecta’ I(a-b/-)

Parentage unknown

H: Peacock BV, pre-1992 **REG:** Peacock BV, 2007
Tepals vivid yellow (9B), shading to brilliant yellow (9C) beside midribs; midribs flushed brilliant greenish yellow (1B). Spots very few; papillae absent; nectaries strong yellow-green (144C); pollen strong orange (169C); stigma greenish white (157D) edged moderate red (182A). Fls 185 mm wide; tepals 90 × 38 mm, margins slightly ruffled, tips strongly recurved. Lvs of short to medium length and of narrow to medium width. Stems of average height to tall, green with darker markings, with average no. to many fls.

‘Semafor’ I(b-c/c-d)

(‘Heloma’ × ‘Mánes’) × (‘Rusalka’ × unknown)

H: B. Mičulka, pre-2005 **N:** B. Mičulka, pre-2005
I: B. Mičulka, 2006 **REG:** B. Mičulka, 2005
Inside of tepals pale apricot, flushed pinkish; flushed beige at tips; throat pale apricot-beige. Outside of tepals white; midribs pinkish and brownish. Brushmarks faint, brown, fine; nectaries beige, frosted; pollen brown; stigma dark crimson. Fls 115 mm wide; tepals 75 × 44 mm, margins slightly ruffled, tips recurved; pedicels 60 mm, with numerous brown spots. Lvs 85 × 13 mm, dull pale green. Stems 0.95 m, stout. Mid-July.

‘Serano’ VIII(a/b) OT

Oriental/Trumpet hybrid mixture × Oriental mixture

H: Testcentrum voor Siergewassen BV, pre-2004
N: Marklily CV, 2004 **REG:** Testcentrum voor Siergewassen BV, 2006
Inside of inner tepals light yellow (10C), tips and beside midveins vivid yellow (9A and 12A respectively). Inside of outer tepals pale greenish yellow (10D), vivid yellow (12A) beside midveins; midveins light yellow (10B), top of midveins and throat vivid yellow (9A and 12A respectively). Outside of tepals pale yellow-green to pale greenish yellow (4D/10D); midribs of outer tepals strong yellow-green at base and top (144B and 144A respectively), and light yellow (10C) between; midribs of inner tepals strong yellow-green (144C) at base. Spots and papillae absent; nectaries strong yellow-green (144A); pollen dark greyish red (N186C); stigma dark purple (79B). Fls 215 mm wide, scented; tepals 140 × 70 mm, margins slightly ruffled, tips recurved. Lvs 175 × 37 mm, dark green. Stems 1.05 m, light green with darker spots and stripes, with c.3 fls.

‘Serena Madonna’ VII(a/b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007
Inside of tepals and outside of inner tepals greenish white (155C). Spots absent; papillae present; nectaries green; pollen orange-brown; stigma green, spotted purple. Tepals of medium length to long and of medium width to broad, margins ruffled, tips slightly

recurved. Lvs of medium length and broad to very broad. Stems of average height to tall, green, with few to average no. of fls. Late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Serene Angel’ VII(a/c)

Seedling 9-13-SP × ‘White Stargazer’

N: Rooyakkers Lelie BV, 2007 **REG:** Rooyakkers Lelie BV, 2007

Inside of inner tepals greenish white (155C); midveins strong yellow-green (N144C), shading to strong yellow (N144B) towards top. Inside of outer tepals greenish white (155C); midveins vivid greenish yellow (2A) in basal half, fading towards top. Outside of outer tepals greenish white (155C), shading to strong yellow-green (144C) beside midribs. Outside of inner tepals tinged light yellow-green (145B). Spots greenish white (155C); papillae absent; nectaries strong yellow-green (N144C). Fls double, 160 mm wide; tepals 75–90 × 25–35 mm, margins slightly ruffled, tips recurved. Lvs 210 × 60 mm, dark green. Stems 0.9–1 m, green, with up to 4 fls.
Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Serrona’ VIII(a-b/-) LO

longiflorum ‘Avita’ × *longiflorum* ‘White Fox’

H: Testcentrum voor Siergewassen BV, pre-2006

N: Marklily CV, 2006 **REG:** Testcentrum voor Siergewassen BV, 2007

Inside of tepals greenish white (155C), with a patch of strong purplish red (60D) at base. Outside of inner tepals mainly greenish white (155C). Spots and papillae absent; nectaries green; pollen orange-brown; stigma grey-green. Fls 200 mm wide; tepals long and of medium width to broad, margins very slightly to slightly ruffled, tips slightly recurved to recurved. Lvs long, and of medium width to broad. Stems 1.5 m, green, with few fls. Very late.
Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Siesta’ VIII(a-b/b) OT

Seedling K 89-10-007 × seedling 89-22-1

H: World Breeding BV, pre-2002 **N:** World Breeding BV, 2002 **REG:** World Breeding BV, 2006

Inside of inner tepals brilliant yellow (9C) shading to brilliant greenish yellow (6A) towards margins. Inside of outer tepals light greenish yellow (6D); throat brilliant yellow (9C). Outside of outer tepals light yellow (10C) shading to 10B towards margins; midribs strong yellow-green (143C) at base and top shading to light yellow-green (144D) between. Outside of inner tepals light yellow (10B), beside and along midribs pale yellow (8D); midribs light yellow-green (144D) at base. Spots and papillae absent; nectaries strong yellow-green (143C); pollen strong reddish orange (169A); stigma moderate yellow-green (138C). Fls 180 mm wide, slightly scented; tepals 110 × 48 mm, margins smooth, tips recurved. Lvs 175 × 30 mm, dark green. Stems 0.6 m, green with darker spots, with c.3 fls.

‘Silvano’ VIII(b-a/b) OT

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2006

Inside of inner tepals light greenish yellow (7D) shading to 6D towards margins; tips and throat pale yellow-green (4D). Inside of outer tepals brilliant greenish yellow (3A) shading to pale yellow-green (4D) towards margins. Outside of outer tepals pale yellow-green (4D); midribs strong yellow-green at base and top (143C and 144B respectively), shading to light yellow-green (154D) flushed moderate red (179B) between.

Outside of inner tepals pale greenish yellow (1D), pale yellow-green (4D) beside midribs; midribs strong yellow-green (144C) at base shading to pale greenish yellow (1D), and light yellow-green (154D) at top. Spots absent, papillae light greenish yellow (7D); nectaries strong yellow-green (143C); pollen dark reddish orange (172B); stigma dark greyish purple (N92A). Fls 255 mm wide, scented; tepals 150 × 75 mm, margins ruffled, tips recurved. Lvs 210 × 50 mm. Stems 1–1.3 m, green with darker markings, with *c.* 4 fls.

‘Sky Treasure’ VIII(b/a) LO

‘White Explosion’ × seedling LO 95008-N

H: De Jong Beheer BV, pre-2004 **N:** De Jong Lelies Holland BV, 2007

Tepals yellowish white (155D); basal third of midveins, from above nectaries, light green, shading to deep purplish pink (N57C); midribs light yellow-green (144D) at base shading to 150D; throat light yellow-green (145C). Spots and papillae absent; nectaries strong yellow-green (144B); pollen dark reddish orange (175B); stigma greenish white (192D). Fls 270 mm wide, scented; tepals 190 × 70 mm, margins smooth to slightly ruffled, tips recurved. Lvs 217 × 42 mm. Stems 1.2 m, green and darker markings in equal measure, with 1 or 2 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Sniega Meita’ VI(b/a)

‘Lugums’ × seedling 236-S-3

H: Ā. Zorgevics, 1990 **G:** Ā. Zorgevics, 1994**N:** Ā. Zorgevics, 1995 **I:** University of Latvia**REG:** University of Latvia Botanical Garden, 2007

Tepals yellowish white (155D), flushed strong purple (77B), especially towards margins on the inside, and towards midribs; base and at least lowermost third of midveins vivid yellow (12A); throat green. Midribs greenish at base with a spot of dark purplish red, shading to yellowish green; top flushed rosy purple. Nectaries clear; pollen pale brown; stigma pale purple. Fls 140 mm wide, strongly scented; tepals 120 × 40 mm, margins smooth, tips slightly recurved. Lvs 80 × 15 mm, scattered. Stems 1.2 m, green, with up to 9 fls. Late July.

Published in *Lilium Balticum. Latvijas liliju selekcionāru izaudzētās lilijas*, No. 1 (1995). The name means Snow Daughter in Latvian.

‘Socrates’ VIII(b-a/b-c) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV,2003 **REG:** World Breeding BV, 2006

Inside of inner tepals pale purplish pink (65C); inside of outer tepals light purplish pink (65B); all shading to yellowish white (N155D) towards margins and brilliant yellow-green (150C) above nectaries; midveins of outer tepals strong purplish pink (68B); throat yellowish white (N155D). Outside of outer tepals very pale purple (69D) shading through 69B to yellowish white (N155D) margins; midribs strong yellow-green at base and top (143C, and 144C with 144A, respectively), shading to light yellow-green (145C) between. Outside of inner tepals very pale purple (69D) shading to yellowish white (N155D) towards margins; midribs strong yellow-green (144C) at base shading to light yellow-green (150D), and brilliant yellow-green (149C) at top. Spots sparse, strong purplish red (58B); papillae absent; nectaries strong yellow-green (144B); pollen strong orange (N25A); stigma light greenish grey (190D). Fls 250 mm wide, scented, dark purplish pink (186C) when young; tepals 140 × 62 mm, margins slightly ruffled, tips strongly recurved. Lvs 180 × 26 mm. Stems 1.2–1.6 m, green with darker markings, with *c.* 4 fls.

‘Sokolka’ VI(c/c-d)

(‘Oneida’ × Apricot Trumpets Group) × (‘Lady Alice’ × (*henryi* × Palmer seedling 56227))

H: B. Mičulka, pre-2007 **N:** B. Mičulka, pre-2007**I:** B. Mičulka, 2007 **REG:** B. Mičulka, 2007

Inside of tepals beige-apricot; tips pale beige; base and throat beige-orange. Outside of tepals paler, shading to pale pink. Nectaries dark green; pollen ochre-apricot; stigma green. Fls 140 mm wide; tepals 100 × 29 mm, margins strongly ruffled, tips recurved; pedicels 65 mm, spotted brown. Lvs 85 × 18 mm, dark green. Stems 1.15 m, stiff, brown-red, infl. short. Early July.

‘Soledo’ VII(a/b)

Parentage unknown

H: Mak ’t Zand BV, pre-2003 **N:** Liliande BV, 2003**REG:** Mak Breeding BV, 2006

Inside of inner tepals strong purplish red (60B to 64C), shading to greenish white (155C) towards margins.

Inside of outer tepals deep red (53A) to strong purplish red (60D), shading to greenish white (155C) towards margins. Outside of tepals deep purplish pink (N66D). Spots and papillae absent; nectaries pale green; pollen orange-brown; stigma purple with small spots. Tepals of medium length and width, margins ruffled, tips recurved. Lvs of medium length and broad to very broad. Stems 1 m, green with some darker markings, with up to 4 fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Spektákl’ I(a/b-c)

(‘Connecticut Beauty’ × ‘Maléřečka’) × (‘Max und Moritz’ × ‘Dagmar’)

H: B. Mičulka, pre-2005 **N:** B. Mičulka, pre-2005**I:** B. Mičulka, 2006 **REG:** B. Mičulka, 2005

Inside of tepals, including throat, yellowish; tips pink. Outside of tepals pale pink, greenish at base; midribs

darker pink. Brushmarks small, fine, mahogany; spots large, fairly numerous, mahogany, over two-fifths of each tepal; nectaries pink, frosted; pollen and stigma dark brown. Fls 132 mm wide; tepals 75 × 37 mm, tips very slightly recurved; pedicels 55 mm, pale green. Lvs 105 × 15 mm, dull pale green. Stems 0.85 m, green. Late June.

'Spinetta' VIII(a/b) LA

Seedling BG 97-14 × seedling AG 95-10
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006
Inside of tepals brilliant greenish yellow (3A), margins and midveins 5B; throat pale greenish yellow (2D). Outside of tepals light greenish yellow (3C) shading to brilliant greenish yellow (5A) towards margins; midribs strong yellow-green (143C) at base shading to light greenish yellow (3C); midribs of outer tepals strong yellow-green (143A) at top. Spots strong purplish red (60B); papillae absent; nectaries strong yellow-green (143B); pollen dark orange (N163A); stigma brilliant greenish yellow (1B) and moderate purplish red (59C). Fls 180 mm wide, becoming paler with age; tepals 140 × 58 mm, margins smooth, tips recurved. Lvs 170 × 32 mm. Stems 1.05–1.30 m, dark with green markings, with 3–4 fls.

'Stainless Steel' I(a/b)

'Fata Morgana' F₂ × 'Fata Morgana' F₂
H: Laan Flora Facilities BV, pre-2005 **N:** Laan Flora Facilities BV, 2005 **REG:** Laan Flora Facilities BV, 2006
Inside of tepals vivid orange (N30D) shading to vivid reddish orange (N30C) towards margins; throat strong orange-yellow (24B). Outside of outer tepals strong orange (N25A); midribs moderate olive-brown (199A) at base, becoming a broad, strong orange (24A) stripe, and strong yellow-green (143B) with some moderate purplish red (59C) at top. Outside of inner tepals strong orange (25A) shading to N25A towards margins; midribs moderate orange (170D) at base shading to strong orange (N25C). Spots and papillae absent; nectaries light olive (152B); pollen absent; stigma vivid orange (28B). Fls 145–180 mm wide; tepals 82 × 40 mm, margins smooth, tips straight. Lvs 150 × 17 mm. Stems 1.2–1.4 m, green with some darker markings at top, with c.14 fls.

'Sugar Diamond' VIII(a/b) LA

'Paola' × seedling LA 93064-32 tetraploid
H: De Jong Lelies Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2006
Inside of tepals dark purplish pink (186C), with a large spot of pale yellow-green (155A) above throat; throat pale greenish yellow (1D). Outside of outer tepals moderate purplish red to dark purplish pink (186B/186C); midribs strong yellow-green (144C). Outside of inner tepals dark purplish pink (186C); midribs strong yellow-green (144B) at base. Spots and papillae absent; nectaries strong yellow-green (143B); pollen dark orange (N163A); stigma light yellow-green (145C). Fls 185 mm wide; tepals 110 × 51 mm, margins smooth,

tips slightly recurved. Lvs 185 × 19 mm. Stems 1–1.1 m, green, basal half with darker markings, with c.5 fls.

'Summercarnival' I(a/b-c)

Seedling G 93-22 × seedling AG 95-14
H: Vletter en Den Haan Beheer BV, pre-2004
N: Vletter en Den Haan Beheer BV, 2004
REG: Vletter en Den Haan Beheer BV, 2006
Inside of tepals vivid yellow (14A), 17B above throat; throat light greenish yellow (1C). Outside of tepals vivid yellow (12A) shading to 14A towards margins; midribs strong yellow-green at base and top (144C and 144A respectively). Spots absent; papillae vivid yellow (17B); nectaries moderate olive-green (137A); pollen strong orange (N163B); stigma deep purplish red (59B). Fls 185 mm wide; tepals 115 × 61 mm, margins smooth, tips slightly recurved. Lvs 135 × 20 mm. Stems 1.05–1.45 m, green, with c.4 fls.

'Sunny Crown' VIII(a/b) OA

'Sorbonne' × 'Esra'
H: De Jong Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2006
Inside of tepals pale yellow-green (4D), with large deep red (60A) blotch extending from above nectaries to up to half of each tepal (larger on outer tepals); margins brilliant greenish yellow (5A); throat greenish white (157D). Outside of tepals greenish white (157D); margins brilliant greenish yellow (5A); midribs strong yellow-green (144B) at base shading to pale yellow-green (4D); midribs of outer tepals strong yellow-green (144B) at top. Spots deep red (60A); papillae greenish white (157D); nectaries strong yellow-green (144A); pollen dark reddish orange (175B); stigma deep purple (83B). Fls 170 mm wide; tepals 110 × 48 mm, margins slightly ruffled, tips recurved. Lvs 160 × 22 mm, dark green. Stems 1.2–1.3 m, pale green with darker markings, with up to 7 fls.
Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Suzuka' I(a-b/b)

Parentage unknown
H: Gebr. Mak BV, pre-1991 **REG:** Gebr. Mak BV, 2007
Inside of tepals vivid yellow (12A). Outside of tepals brilliant yellow (9C), shading to vivid yellow (9B) towards margins; midribs strong yellow-green (143C). Spots dark red (187A); papillae absent; nectaries strong yellow-green (143A); pollen moderate brown (165A); stigma dark red (187A). Fls 200 mm wide; tepals 95 × 45 mm, margins ruffled, tips slightly recurved. Lvs 140 × 13 mm. Stems 0.85 m, pale green, with up to 8 fls.

'Suzzarra' I(a/b)

Seedling AM 96-17 × seedling AW 96-04
H: Vletter en Den Haan Beheer BV, pre-1991
N: Vletter en Den Haan Beheer BV, 1991
REG: Vletter en Den Haan Beheer BV, 2007
Tepals yellowish white (155D). Inside of tepals tinged light yellow-green (145D) above nectaries; midveins greenish white (157D) on inner tepals, light yellow-green (145D) on outer; throat light yellow-green (145C).

Outside of outer tepals strong yellow-green (144C) beside midribs; all midribs 144C; midribs of outer tepals tinged strong purplish red (60B) at base and top. Spots and papillae absent; nectaries strong yellow-green (143B); pollen dark orange (N163A); stigma deep purplish red (59B) with strong yellow-green (N144D). Fls 195 mm wide; tepals 114 × 55 mm, margins smooth, tips slightly recurved. Lvs 175 × 25 mm. Stems 1.3 m, green with darker spots, with up to 6 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Taranto’ I(a/-)

Seedling R.S. 516 × seedling R.S. 843

H: Vletter & Den Haan, 1982 **G:** Vletter & Den

Haan, 1985 **N:** Vletter & Den Haan, 1989

I: Vletter & Den Haan, c.1990 **REG:** Vletter & Den Haan, 1990

Inside of tepals strong reddish purple (70B) to deep purplish pink (70C), with a chrome-yellow flame.

Outside of tepals strong reddish purple (70B); margins moderate purplish red (186A). Spots few, small, dark red (187A), on inner tepals only; nectaries light yellow-green (145C/D); pollen vivid orange-yellow (23A). Fls 160 mm wide; tepals 90 × 30 mm, margins ruffled, tips strongly recurved. Lvs 150 × 15 mm, scattered, mid-green. Stems 1.4 m, green with darker markings, with up to 15 fls. May.

‘Tasmania’ I(a/-)

Parentage unknown

H: Mak-Leek Inc., pre-2006 **REG:** Mak-Leek Inc., 2006

Inside of tepals vivid yellow (14A), with patches and spots of greyish purple to dark greyish purple (N77A/202A). Outside of inner tepals vivid yellow (13A). Papillae present; nectaries yellow-green; pollen orange-brown; stigma dark purple. Tepals very short to short, narrow, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of short to medium length, narrow. Stems of average height to tall, green, with few to average no. of fls. Early to mid-season.

‘Tenorio’ I(a-b/-)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2005

Inside of tepals strong orange (N25B), paler towards tips; midveins strong orange (25B) at base. Outside of inner tepals strong orange (N25B/C). Spots absent, papillae faint; nectaries orange; pollen red-brown; stigma purple-red. Tepals short and of narrow to medium width, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length to long and of narrow to medium width. Stems of average height to tall, dark, with few fls. Early to mid-season.

‘Terni’ VIII(b/b) LO

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2006

Tepals, including throat, yellowish white (155D); midveins pale greenish yellow (2D) above nectaries;

midribs of outer tepals strong yellow-green at base and top (144C and 143C respectively), shading to pale yellow-green (4D) between; midribs of inner tepals strong yellow-green (141D) at base shading to pale yellow-green (155A), and pale greenish yellow (2D) at top. Spots absent, papillae present; nectaries strong yellow-green (144C); pollen strong brown (172A); stigma pale yellow-green (193D). Fls 250 mm wide; tepals 144 × 65 mm, margins slightly ruffled, tips recurved. Lvs 240 × 65 mm. Stems 1.2–1.7 m, green, with c.4 fls.

‘The Legend’ VII(a-b/-)

Parentage unknown

H: World Breeding BV, pre-2001 **N:** World Breeding BV, 2001 **REG:** World Breeding BV, 2005

Tepals yellowish white (155D); inside of inner tepals brilliant yellow (8A) in basal portion; midveins of outer tepals yellow at base. Spots few, more conspicuous on older fls, yellow-brown; papillae present; nectaries green; pollen light brown; stigma purple. Tepals of medium length to long and of medium width, margins ruffled, tips recurved to strongly recurved. Lvs of medium length, broad. Stems of average height, green with darker stripes and spots, with few fls. Mid-season to late.

Syn. ‘Legend’

Although granted Dutch Plant Breeders’ Rights under the cultivar epithet ‘Legend’, this lily was granted European P.B.R. under the epithet ‘The Legend’.

‘Thomas’s Giant’ VI(b/a-b)

(*regale* × Pink Perfection Group) × (*regale* × Pink Perfection Group)

H: R. Thomas, 1982 **G:** N. Martschinke, 1988 **N:** N. Martschinke, 1999 **I:** G. Ewald, 1999 **REG:** N. Martschinke, 2006

Tepals yellowish white (155D); throat brilliant greenish yellow (5A/B); midribs of outer tepals moderate yellowish brown (N199C) flushed light yellow-green (145B); midribs of inner tepals 145B. Spots and papillae absent; nectaries dark greyish reddish brown (200A); pollen strong orange (168A); style strong yellow-green (145A), dark red (59A) below the stigma; stigma greyish yellow-green (194B) flushed moderate purplish red (186B). Fls 145 mm wide, slightly scented; tepals 120 × 47 mm, margins slightly ruffled, tips recurved; with secondary buds. Lvs scattered, 185 × 15 mm. Stems 1.85 m, mid-green, with c.33 fls. Late July.

Published in Gūdrūn Ewald’s 1999 bulb list.

Nomenclatural standard: colour print supplied by registrant (WSY 0100424). Name commemorates the breeder, Dick Thomas, who died in 1997.

‘Tiny Athlete’ I(a-b/b)

Parentage unknown

H: Mak ’t Zand BV, pre-2003 **N:** Liliande BV, 2003 **REG:** Mak Breeding BV, 2006

Inside of tepals pale purplish pink (56B) on opening, maturing to light pink (49C) to light purplish pink (62C to 65B), and paling to light yellowish pink (38C) with age; midveins light purple-red at base. Outside of tepals pale purplish pink (56C) on opening, pale pink

(49D) at maturity. Spots very few; papillae present; nectaries green; filaments pale yellowish pink (36D); anthers strong brown (172A); pollen dark reddish orange (173A); style light yellow-green (145C); stigma light yellowish pink (36A). Buds strong yellow-green (144C) towards base, shading to pale purplish pink (62D) towards top. Fls 150 mm wide, slightly scented; tepals 87–89 × 35–43 mm, margins slightly ruffled to ruffled, tips recurved. Lvs 68–71 × 11–12 mm, scattered, dark green (135A). Stems 0.6 m, green with dark red (187A) spots and stripes, with up to 10 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Bee' I(a-b/b)

Parentage unknown

H: Mak 't Zand BV, pre-2003 **N:** Liliande BV, 2003 **REG:** Mak Breeding BV, 2006
Inside of tepals vivid yellow (16A to 17B) to strong orange-yellow (17A); more yellow towards margins; midveins strong orange-yellow (17A). Outside of tepals vivid yellow (14A/B) to light orange-yellow (16B); midribs greenish. Spots few to average no., dark red (187A), towards base of each tepal; papillae present; nectaries orange-yellow; filaments brilliant yellow (20A); anthers moderate red (183C); pollen strong orange (168B); style brilliant greenish yellow (1B); stigma light orange-yellow (24C) with red star. Buds brilliant yellow (13B). Fls 140 mm wide, slightly scented; tepals 92–94 × 41–42 mm, margins (very) slightly ruffled, tips slightly recurved to recurved. Lvs 70–130 × 12–19 mm, scattered, dark yellowish green (139A). Stems c.0.45 m, strong yellow-green (143B), with up to 9 fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Bell' I(a/b)

Parentage unknown

H: Mak 't Zand BV, pre-2003 **N:** Liliande BV, 2003 **REG:** Mak Breeding BV, 2006
Tepals strong purplish red (54A) to deep purplish pink (54B); inside margins, and blotch surrounding nectaries, pale yellow (20C/D); midveins vivid yellow (17C) at base. Spots few; papillae present; nectaries green, shading to orange-yellow; pollen orange; stigma purple-red. Tepals short to very short, and narrow, margins slightly ruffled, tips slightly recurved to recurved. Lvs of short to medium length and of medium width. Stems 0.55 m, green, with few to average no. of fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Dessert' I(a/b)

Parentage unknown

H: Mak 't Zand BV, pre-2005 **N:** Mak 't Zand BV, 2005 **REG:** Mak Breeding BV, 2006
Inside of tepals vivid yellow (inner 14B, outer 14A); midveins vivid orange-yellow (23A) at base. Outside of tepals vivid yellow (13A) to brilliant yellow (13B). Spots absent; papillae inconspicuous; nectaries green; pollen orange; stigma yellow. Tepals short and narrow, margins slightly ruffled, tips slightly recurved to

recurved. Lvs of short to medium length and of narrow to medium width. Stems 0.55 m, green with darker spots and stripes, with up to 6 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Diary' I(a-b/b)

Parentage unknown

H: Mak 't Zand BV, pre-2005 **N:** Mak 't Zand BV, 2005 **REG:** Mak Breeding BV, 2006
Inside of tepals strong orange (25A) to vivid yellowish pink (28A) or vivid orange (N30D); base of midveins vivid reddish orange (N30C) to vivid orange (N30D). Outside of tepals vivid yellowish pink (28A) to vivid orange (28B), becoming vivid reddish orange (30B) with age. Spots absent; papillae inconspicuous; nectaries orange; filaments strong orange (25B/26B); anthers moderate reddish brown (175A); pollen dark reddish orange (175B); style strong orange (169C); stigma moderate red (179A). Buds strong orange (25A) to vivid yellowish pink (28A). Fls 120 mm wide, slightly scented; tepals 95–100 × 34–48 mm, margins slightly ruffled to ruffled, tips recurved. Lvs 90 × 16–20 mm, scattered, moderate olive-green (137B). Stems 0.45 m, moderate yellow-green (146B) with darker markings, with up to 8 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Dino' I(a-b/b)

Parentage unknown

H: Mak 't Zand BV, pre-2005 **N:** Mak 't Zand BV, 2005 **REG:** Mak Breeding BV, 2006
Inside of inner tepals vivid reddish orange (N30C) to vivid orange (N30D) or vivid yellowish pink (28A); inside of outer tepals vivid orange (28B), more red towards margins; midveins strong orange (N25B) at base. Outside of tepals strong orange (26B) or vivid orange (28B), more red towards margins. Spots absent; papillae present; nectaries orange; filaments light orange-yellow (24C); anthers dark red (183A); pollen strong brown (172A); style brilliant orange (25C); stigma dark red (187A). Buds strong orange (25B), midribs greenish. Fls 180 mm wide, slightly scented; tepals 90–95 × 30–50 mm, margins slightly ruffled, tips slightly recurved. Lvs 77–155 × 15–24 mm, moderate olive-green (137B). Stems 0.6 m, green with dark red (187A) markings, with up to 9 fls. Early to mid-season. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Ghost' I(a/b-c)

Parentage unknown

H: Mak 't Zand BV, pre-2003 **N:** Liliande BV, 2003 **REG:** Mak Breeding BV, 2006
Inside of tepals deep red (60A), paler towards top; base of midveins and outside of tepals deep red (53A) to dark red (59A). Spots few; papillae present; nectaries deep red; filaments deep purplish red (187D); anthers deep red (185A); pollen strong orange (168A); style moderate red (180A); stigma moderate yellow-green (146B). Fls 160 mm wide, slightly scented; tepals 75–80 × 40 mm, margins slightly ruffled, tips slightly

recurved. Lvs 75–110 × 13–21 mm, scattered, moderate olive-green (137A). Stems short, moderate yellow-green (146B), with up to 7 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Tiny Hope’ I(a-b/-)

Parentage unknown

H: Mak ‘t Zand BV, pre-2003 **N:** Liliande BV, 2003 **REG:** Mak Breeding BV, 2006

Inside of inner tepals moderate red (N34A) to strong red (46A); inside of outer tepals strong red (46A); paler towards tips; midveins strong yellowish pink (35C), purple-red at base. Outside of tepals moderate red (N34A) to vivid reddish orange (42A). Spots dark red (187A), towards base of each tepal; papillae present; nectaries orange-purple-red; filaments moderate reddish orange (35B); anthers strong reddish orange (169A); pollen vivid orange (28B); style vivid reddish orange (44C); stigma dark red (187A). Buds strong red (46A), greenish towards tip. Fls 180 mm wide, slightly scented; tepals 100–110 × 30–40 mm, margins slightly ruffled, tips slightly recurved to recurved. Lvs 40–62 × 12–18 mm, scattered, deep yellowish green (141A). Stems of short to average height, moderate yellow-green (146B) with some darker spots and stripes, with up to 9 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Tiny Icon’ I(a-b/b-c)

Parentage unknown

H: Mak ‘t Zand BV, pre-2003 **N:** Mak ‘t Zand BV, 2003 **REG:** Mak Breeding BV, 2006

Inside of tepals strong purplish pink (55B) or strong purplish red (58C), maturing to deep purplish pink (55A/58D) before fading to strong pink (54C); paler towards tips; midveins greenish white at base. Outside of tepals light purplish pink (55C) to deep purplish pink (58D). Spots very few; papillae few; nectaries green; filaments yellowish white (155B); anthers strong orange (24A); pollen strong orange-yellow (24B); style light yellow-green (145C); stigma strong reddish purple (72B). Buds pale purplish pink (65C). Fls 125 mm wide, slightly scented; tepals 80 × 36–47 mm, margins slightly ruffled, tips slightly recurved. Lvs 67–82 × 12–29 mm, scattered, strong yellow-green (143A). Stems 0.7 m, moderate yellow-green (146B) with darker spots and stripes, with up to 10 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Tiny Invader’ I(a-b/b-c)

Parentage unknown

H: Mak ‘t Zand BV, pre-2004 **N:** Mak ‘t Zand BV, 2004 **REG:** Mak Breeding BV, 2006

Tepals open strong orange (25A); inner tepals mature to vivid yellowish pink (28A), becoming vivid reddish orange (30B) or vivid orange (N30D) with age; outer tepals mature to vivid yellowish pink (28A) to brilliant orange (29A) or vivid orange (N30D). Spots few, dark red (187A), towards base; papillae present; nectaries orange; filaments and anthers dark reddish orange (172B); pollen moderate reddish orange (171A);

style strong orange (169D); stigma strong reddish orange (169A). Buds light orange (28C). Fls 140 mm wide, slightly scented; tepals 108 × 32–42 mm, margins smooth to very slightly ruffled, tips recurved. Lvs 83–92 × 15 mm, scattered, strong yellow-green (143B). Stems 0.45 m, strong yellow-green (143C), with up to 9 fls. Early.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Tiny Kissed’ I(a/b)

Parentage unknown

H: Mak ‘t Zand BV, pre-2003 **N:** Liliande BV, 2003 **REG:** Mak Breeding BV, 2006

Inside of tepals deep red (53A) at base, shading to vivid red (44A) at tips. Outside of tepals vivid red (44A) to vivid reddish orange (44B). Spots few, darker red than tepal colour; papillae present; nectaries red; pollen reddish brown; stigma dark purple. Tepals short to very short and narrow, margins (very) slightly ruffled, tips slightly recurved; infl. very downy. Lvs of medium length and width. Stems 0.75 m, green with darker spots and stripes, with few to average no. of fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Tiny Nanny’ I(a-b/b-c)

Parentage unknown

H: Mak ‘t Zand BV, pre-2003 **N:** Liliande BV, 2003 **REG:** Mak Breeding BV, 2006

Inside of tepals open yellowish white (155D), maturing to yellowish white (155B) or greenish white (155C). Outside of tepals yellowish white (155B/155D), light yellow-green (150D) towards base. Spots and papillae absent; nectaries green; filaments brilliant yellow-green (150C); anthers greyish red (178A); pollen dark red (183A); style strong yellow-green (144C); stigma deep red (185A). Buds yellowish white (155D). Fls 140 mm wide, slightly scented; tepals 100–111 × 30–40 mm, margins slightly ruffled, tips slightly recurved. Lvs 76–90 × 12 mm, scattered, brilliant yellow-green (149A). Stems 0.45 m, strong yellow-green (144B) with darker spots and stripes, with up to 8 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Tiny Puppet’ I(a-b/b)

Parentage unknown

H: Mak ‘t Zand BV, pre-2004 **N:** Mak ‘t Zand BV, 2004 **REG:** Mak Breeding BV, 2006

Inside of tepals deep red (185A) to dark red (187A/B/C). Outside of tepals deep red (53A) to moderate purplish red (185C). Spots moderate red (185B); papillae present; nectaries orange-red; filaments strong purplish red (60B); anthers moderate reddish orange (171A); pollen strong orange (170A); style strong orange (168B); stigma greyish red (178A). Buds greyish red (182B). Fls 140 mm wide, slightly scented; tepals 85 × 29–41 mm, margins very slightly to slightly ruffled, tips slightly recurved. Lvs 70 × 16 mm, scattered, moderate yellowish green (139B). Stems 0.6 m, moderate yellow-green (146B), with up to 10 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Sensation' I(a-b/b-c)

Parentage unknown

H: Mak 't Zand BV, pre-2003 **N:** Liliande BV,

2003 **REG:** Mak Breeding BV, 2006

Inside of tepals vivid yellow (13A/14B), basal half with large, deep red (185A) or dark red (187B/C) spots and smears; margins narrow, vivid yellow (13A); midveins yellow at base. Outside of tepals vivid yellow (9B) to brilliant yellow (9C/13B/13C). Papillae present; nectaries yellow-green; filaments light greenish yellow (8B); anthers and pollen moderate orange (167A and 167B respectively); style vivid yellow (9B) to brilliant yellow (9C); stigma moderate orange (168C). Buds brilliant greenish yellow (1B) to light yellow-green (145C). Fls 160 mm wide, slightly scented; tepals 101–104 × 32–45 mm, margins slightly ruffled, tips slightly recurved. Lvs 67–74 × 13 mm, scattered, moderate olive-green (137B). Stems 0.55 m, moderate yellow-green (146B), with up to 10 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Skyline' I(a-b/b)

Parentage unknown

H: Mak 't Zand BV, pre-2003 **N:** Liliande BV,

2003 **REG:** Mak Breeding BV, 2006

Inside of tepals open vivid orange-yellow (23A), shading to strong orange (25A) towards margins; becoming strong orange (24A) to vivid orange (28B) towards centre; vivid yellowish pink (28A) towards margins, and strong orange (30D) towards base; midveins strong orange (N25C), vivid orange-yellow (23A) at base. Outside of tepals becoming vivid yellowish orange (30C) at maturity. Outside of tepals open light orange-yellow (23C) to strong orange (25A); becoming vivid yellowish pink (28A) to brilliant orange (29A) or strong orange (N25A/B). Spots dark red (187A), random, towards base of each tepal; papillae present; nectaries green; filaments strong yellowish pink (32C); anthers moderate reddish orange (171A); pollen and style strong orange (170A and 168B respectively); stigma greyish red (178A). Buds light orange-yellow (22B) to brilliant orange (25C). Fls 160 mm wide, slightly scented; tepals 105–110 × 33–48 mm, margins slightly ruffled, tips slightly recurved. Lvs 80–110 × 10–30 mm, scattered, moderate yellowish green (139B). Stems 0.5 m, strong yellow-green (143A) with darker spots and stripes, with up to 9 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Snowflake' I(a-b/b)

Parentage unknown

H: Mak 't Zand BV, pre-2003 **N:** Liliande BV, 2003

REG: Mak Breeding BV, 2006

Inside of tepals open yellowish white (158D); outside pale yellow-green (157C). All tepals develop to yellowish white (155B) to pale yellow-green (157A), becoming yellowish white (155B/158C) at maturity. Spots few; papillae present; nectaries green; filaments yellowish white (155D); anthers strong orange (170A); pollen moderate orange (170B); style strong yellow-green (145A); stigma light purple (78C). Buds light yellow-green (145C) towards base, brilliant

yellow (14C) towards tip. Fls 140 mm wide, slightly scented; tepals 93 × 30–45 mm, margins (very) slightly ruffled, tips slightly recurved. Lvs 81–102 × 10–14 mm, scattered, strong yellow-green (143A). Stems 0.8 m, moderate olive-green (146A) with darker spots and stripes, with up to 10 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Star' I(a-b/b)

Parentage unknown

H: Mak 't Zand BV, pre-2003 **N:** Liliande BV,

2003 **REG:** Mak Breeding BV, 2006

Inside of tepals vivid reddish orange (N30C) to vivid orange (N30D); midveins strong orange (N25C). Outside of tepals strong orange (30D) with a greenish flush. Spots and papillae absent; nectaries green; pollen reddish brown; stigma orange-red. Tepals short and narrow, margins (very) slightly ruffled, tips slightly recurved to recurved. Lvs of medium length and of narrow to medium width. Stems 0.55 m, green, with few fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Todd' I(a-b/c)

Parentage unknown

H: Mak 't Zand BV, pre-2003 **N:** Liliande BV,

2003 **REG:** Mak Breeding BV, 2006

Inside of tepals yellowish white (155B), shading to light purplish pink (55C/65B) towards margins and tips; midveins light yellowish pink (26D). Outside of tepals light purplish pink (55C) to pale purplish pink (55D) or very pale purple (69A). Spots few, dark red (187A); papillae present; nectaries green at base, shading to pink; filaments very pale purple (69B); anthers brownish orange (165B); pollen strong orange-yellow (163B); style strong greenish yellow (153B); stigma greyish yellow-green (191B) or light purple-red. Buds pale purplish pink (65C). Fls 170 mm wide, slightly scented; tepals 75–82 × 30–40 mm, margins slightly ruffled to ruffled, tips recurved. Lvs 85–95 × 18 mm, scattered, deep yellowish green (141A). Stems of short to average height, strong yellow-green (144A), with up to 9 fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Tiny Toes' I(a-b/b-c)

Parentage unknown

H: Mak 't Zand BV, pre-2004 **N:** Mak 't Zand BV,

2004 **REG:** Mak Breeding BV, 2006

Tepals open strong yellowish pink (37B), becoming 37A or vivid reddish orange (44C) to deep yellowish pink (44D); paling with age to strong red (39A); midveins orange-yellow at base. Spots and papillae absent; nectaries green; filaments moderate pink (49B); anthers strong reddish orange (169A); pollen strong orange (169C); style pale yellow (161C); stigma strong reddish purple (78B). Buds light yellow-green (145B) towards base, pale purplish pink (62D) towards tip. Fls 145 mm wide, slightly scented; tepals 92–95 × 32–43 mm, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs 88–93 × 11–21 mm, scattered, strong yellow-green (143A). Stems

0.75 m, strong yellow-green (144A) with some darker spots and stripes, with up to 10 fls. Mid-season. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Tobin’ I(b/c)

‘Melissa Jamie’ × ‘Unique’

H: T.D. Willoughby, 1991 **G:** T.D. Willoughby, 1993 **N:** T.D. Willoughby, 2003 **REG:** T.D. and S.K. Willoughby, 2006

Inside of tepals light yellow (19B) (bleaching to white), shading to deep pink (52C) towards margins; tips vivid purplish red (67B); throat and outside of tepals strong purplish red (67A). Spots absent; nectaries strong yellow-green (144B); pollen vivid reddish orange (34A); stigma strong red (53B). Fls 150 mm wide; tepals 80 × 35 mm, margins smooth, tips slightly recurved. Lvs 100 × 20 mm, scattered, moderate olive-green (137A). Stems 1.2 m, dark greyish reddish brown (200A), with up to 25 fls. Early to mid-July (USDA Zone 3, Alberta, Canada). Named after Tobin Banks, a close friend of the hybridizer.

‘Touching’ VIII(a-b/b-c) OT

A temporary name for ‘Lavon’ prior to registration.

‘Tramín’ I(a/b)

‘Elfi’ × ‘Tango’

H: K. Braun, pre-2006 **N:** K. Braun, pre-2006

I: K. Braun, 2006 **REG:** K. Braun, 2006

Inside of tepals violet-red; outside of tepals yellow. Spots over two-thirds of each tepal, dark violet; nectaries red; pollen and stigma brownish red. Fls 128 mm wide; tepals 69 × 36 mm, tips straight; pedicels 110 mm long, brown; with secondary buds. Lvs dull mid-green. Stems 0.7 m, infl. long. Late June.

‘Travatore’ VIII(b-a/b) OT

Seedling PG 96-43 × seedling RH 97-09

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of inner tepals brilliant yellow (8A) to light greenish yellow (8B); tips light greenish yellow (4C) to light yellow-green (2C); throat strong yellow-green (144A). Inside of outer tepals brilliant yellow (8A); shading to light greenish yellow (8B) at margins, and 8C towards tips. Outside of outer tepals pale yellow-green (4D); margins light greenish yellow (8B/C); midribs strong yellow-green (144C) at base and top. Outside of inner tepals light greenish yellow (8B/C), shading to greenish white (157D) beside midribs; midribs light greenish yellow (1C), brilliant greenish yellow (1B) at top. Spots absent; papillae vivid yellow (9A); nectaries strong yellow-green (144A); pollen moderate reddish orange (N172A); stigma dark red (187A). Fls 245 mm wide; tepals 135 × 80 mm, margins slightly ruffled, tips recurved. Lvs 236 × 37 mm, scattered, dark green. Stems 1.3 m, pale green, with up to 4 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Triora’ VII(a/c-b)

Seedling RM 96-10 × seedling RW 96-05

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2007

Inside of inner tepals strong purplish red (64B/C); throat greenish white (155C). Inside of outer tepals strong purplish red (67A), shading to moderate purplish red (64A) towards margins. Outside of tepals strong purplish red (64B) on a paler ground; midribs light purplish pink (65B). Spots deep red (60A); papillae absent; nectaries strong yellow-green (143B); stigma light grey (191C). Fls 255 mm wide, slightly scented; tepals 146 × 75 mm, margins smooth to slightly ruffled, tips slightly recurved. Lvs 200 × 50 mm. Stems 1.1–1.25 m, pale green with darker markings, with up to 3 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Tuareg’ VIII(b-a/b) LA

(*longiflorum* ‘Avita’ × ‘Nerone’) × ‘Amarone’

H: Testcentrum voor Siergewassen BV, pre-2004

N: Marklily, 2004 **REG:** Testcentrum voor Siergewassen BV, 2006

Inside of inner tepals moderate red (185B); inside of outer tepals deep red (185A), all shading to dark red (187B) towards margins; throat moderate purplish pink (186D) flushed yellow-green. Outside of outer tepals dark red to deep purplish red (187C/187D); midribs strong yellow-green (144A) at base and top. Outside of inner tepals deep purplish red (187D) shading to dark red (187C) towards margins and moderate purplish red (185C) beside midribs; midribs dark red (187B). Spots very sparse, dark purplish red (N79A); papillae absent; nectaries strong yellow-green (144A); pollen dark red (183A); stigma moderate red (181A). Fls 190 mm wide, glossy; tepals 110–115 × 35–40 mm, margins ruffled, tips recurved. Lvs 150 × 25 mm, dark green. Stems 1 m, green, with c.8 fls.

‘Una Bonita’ VII(a-b/b)

Parentage unknown

H: World Breeding BV, pre-2001 **N:** World Breeding BV, 2001 **REG:** World Breeding BV, 2007

Inside of tepals strong purplish pink (73B) to light purple (75B); midveins brilliant yellow (10A) at base. Outside of inner tepals light purple (75B) to very light purple (75C). Spots and papillae present; nectaries yellow-green; pollen brown; stigma purple. Tepals short and of narrow to medium width, margins ruffled, tips slightly recurved to recurved. Lvs of medium length and of medium width to broad. Stems of average height to tall, green with darker markings, with few to average no. of fls. Early. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007). ‘Bonita’ was used as a temporary name prior to registration.

‘Vadakste’ VI(c/a)

‘Vidzeme’ × seedling 236-S-11

H: Ā. Zorgevics, 1986 **G:** Ā. Zorgevics, 1991

N: Ā. Zorgevics, 2006 **I:** University of Latvia Botanical Garden, 2006 **REG:** University of

Latvia Botanical Garden, 2006

Inside of tepals moderate purplish pink (78D); midveins greenish in basal half; throat brilliant yellow-green (149C). Outside of tepals light violet flushed brownish; midribs brownish olive-green. Nectaries clear, filaments almost straight, pale green, pollen brown; style, longer than stamens, incurving, and stigma, greenish violet. Fls 130 mm wide, strongly scented, long-lasting; tepals 100 × 30 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 80 × 10 mm, glossy. Stems 0.9 m, green, with c.7 fls. Bulbs purple. Mid- to late July. Published in *Lilium Balticum. Latvijas liliju selekcionāru izaudzētās liliņas*, No. 1 (1995). Nomenclatural standard: colour print supplied by registrant (WSY 0100423).

'Vasari' VII(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Tepals, including throat, yellowish white (155D); midveins flushed light yellow-green (150D) above nectaries. Outside of outer tepals with two, small, deep purplish red (61A) stripes beside midribs at base; midribs strong yellow-green (144C). Spots and papillae yellowish white (155D); nectaries strong yellow-green (143C); pollen dark reddish orange (173A); stigma light grey (188D). Fls 193 mm wide, scented; tepals 120 × 70 mm, margins slightly ruffled, tips slightly recurved. Lvs 197 × 58 mm. Stems 1.2 m, green, with up to 4 fls.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Vendela' VIII(b/a) LO

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2001

N: Marklily CV, 2001

REG: Testcentrum voor Siergewassen BV, 2007

Inside of tepals strong purplish red (71B/C), paler towards tips; midveins green at base. Outside of inner tepals strong reddish purple (70B). Spots and papillae absent; nectaries green; pollen orange-brown; stigma pale grey. Tepals long and broad, margins (very) slightly ruffled, tips (very) slightly recurved. Lvs long to very long, and of medium width to broad. Stems 1.6 m, green with darker spots and stripes, with up to 3 fls. Very late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Veranda' VIII(a/b) OT

Seedling PG 96-43 × seedling RH 99-07

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2007

Inside of tepals light greenish yellow (6D), shading to brilliant greenish yellow (6B) beside midveins and light yellow (10C) towards tips; midveins brilliant greenish yellow (6B); throat pale yellow-green (4D). Outside of outer tepals pale yellow-green (4D), shading to light greenish yellow (5D) towards

margins and with a broad, strong yellowish pink (38A) stripe on one side of each midrib; midribs light yellow-green (144D) at base and strong yellow-green (144C) at top. Outside of inner tepals light greenish yellow (4C), shading to pale yellow-green (4D) beside midribs; midribs light yellow-green (144D) at base. Spots brilliant greenish yellow (6B); papillae absent; nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma deep purple (79C). Fls 220 mm wide, scented; tepals 130 × 75 mm, margins smooth to slightly ruffled, tips recurved. Lvs 225 × 38 mm. Stems 1.3–1.4 m, green, with up to 3 fls. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Vernon White' II(c/d)

('Glacier' × *martagon* var. *cattaniae* 'Album Superbum') × white-flowered *martagon* hybrid from Vernon, Canada

H: E.E. Fox, 1987

G: E.E. Fox, 1994

N: E.E. Fox, 1996

I: Fox Lily Ranch, 2006

REG: E.E. Fox, 2006

Inside of tepals pure white; centre of tepals flushed brilliant yellow-green (150C) beside nectary furrows; throat brilliant yellow-green (149A). Outside of tepals white; midribs moderate yellow-green (138B). Spots and papillae absent; nectaries brilliant yellow-green (149C); pollen strong orange-yellow (22A); stigma light yellow-green (154D). Fls 70 mm wide, slightly scented; tepals 35 × 17 mm, margins smooth, pubescent towards tips, tips strongly recurved. Lvs whorled, 165 × 42 mm. Stems 1.5 m, strong yellow-green (143A) mottled moderate yellow-green (147B), with 18–24 fls. Late June to early July.

Nomenclatural standard: colour print supplied by registrant (WSY 0100422). The pollen parent originated from the Canadian city of Vernon, British Columbia.

'Vidzeme'

Unregistered seed parent of 'Vadakste'. No further details known.

VITARA VII(a/b-c)

Trade designation of 'Zanlortara'.

'Vladimira' VI(b/b)

Green Magic Group × tetraploid Aurelian (Div. VI) hybrid

H: J. Dudek, pre-2006

G: J. Dudek, pre-2006

N: J. Dudek, pre-2006

I: J. Dudek, 2006

REG: J. Dudek, 2006

Inside of tepals lemon yellow; throat green-yellow.

Outside of tepals yellow; midribs brown-bronze.

Nectaries green; pollen orange; stigma pale green. Fls 160 mm wide, slightly scented; tepals 150 × 34 mm, tips recurved; pedicels 70 mm, stout. Lvs 130 × 12 mm, dull, slightly recurved. Stems 1.45 m, stiff, green, with long infl. Early July.

'Voltage' I(a-b/b)

Parentage unknown

H: Mak 't Zand BV, pre-2004

N: Liliande BV, 2004

REG: Mak Breeding BV, 2006

Tepals vivid yellow; inside of inner tepals 13A; inside

of outer tepals 12A/13A; outside of tepals 14B. Spots very few; nectaries green; pollen orange; stigma yellow. Fls 190 mm wide; tepals short to very short and of narrow to medium width, margins slightly ruffled, tips slightly recurved. Lvs of medium length and of narrow to medium width. Stems 0.8–1.15 m, green with darker spots and stripes, with few to average no. of fls. Early to mid-season. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘West Point’ I(a/-)

Parentage unknown

H: Fa Jaap Boon, pre-1999 **N:** Trior Lelie BV/ Koninklijke Van Zanten BV, 1999 **REG:** Fa Jaap Boon, 2007

Inside of tepals vivid yellow (14A); midveins orange-yellow at base. Outside of tepals brilliant yellow (14C); margins vivid yellow (14B). Spots few; papillae present; nectaries green; pollen brown; stigma purple-red. Tepals short to very short and narrow, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length, and of narrow to medium width. Stems of average height to tall, green with darker spots and stripes, with few to average no. of fls. Very early to early.

‘White Collier’ VII(a-b/b)

Parentage unknown

H: Mak t Zand BV, pre-2003 **N:** Liliande BV, 2003 **REG:** Mak Breeding BV, 2006

Tepals greenish white (155C). Spots absent; papillae present; nectaries green; pollen orange-brown; stigma green with small purple spots. Tepals of medium length to long and of medium width to broad, margins ruffled, tips recurved to strongly recurved. Lvs of medium length and broad. Stems 1.15 m, green, with up to 7 fls. Late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘White Continental’ V(b/a)

Parentage unknown

H: C. van Duyn, pre-1990 **REG:** C. van Duyn, 2007

Inside of tepals greenish white (155C) to yellowish white (155D); midveins and throat light yellow-green (145B and 145C/D respectively). Outside of tepals greenish white (155C); midribs yellowish white (155B). Spots and papillae absent; pollen deep orange-yellow (163A); stigma pale yellow-green (155A). Fls 170 mm wide; tepals 170 × 55 mm, margins slightly ruffled, tips strongly recurved. Lvs 190 × 20 mm. Stems 1.3 m, green, with 1 or 2 fls.

‘White Express’ VII(a-b/b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007

Tepals greenish white (155C). Spots absent; papillae present; nectaries pale yellow-green; pollen orange-brown; stigma purple, speckled. Tepals of medium length to long and of medium width, margins ruffled to strongly ruffled, tips recurved to strongly recurved. Lvs of medium length and broad. Stems 1.5 m, green,

with few to average no. of fls. Late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

WHITE IMPRESSION VIII(a/b) LA

Trade designation of ‘Zanlapres’.

‘White Pixels’ I(a/b)

‘Cha Cha Cha’ × white-flowered seedling

H: Laan Flora Facilities BV, pre-2005 **N:** Laan Flora Facilities BV, 2005 **REG:** Laan Flora Facilities BV, 2006

Tepals yellowish white (155D); throat light yellow-green (145D); base of outside of outer tepals light yellow-green (145D) flushed strong purplish red (60B); midribs of outer tepals light yellow-green (145C) at base, and strong yellow-green (144A) with strong purplish red (60B) at top; midribs of inner tepals strong yellow-green (144B) at base shading to strong purplish red (60B); upper two-thirds light yellow-green (145C). Spots numerous in basal half of tepals, a few along margins of inside of tepals and along basal half of margins of outside of outer tepals, dark red (59A); papillae mostly yellowish white (155D), some dark red (59A); nectaries strong yellow-green (143A); pollen strong orange-yellow (17A); stigma strong purplish red (60C). Fls 150 mm wide; tepals 88 × 53 mm, margins smooth, tips slightly recurved. Lvs 130 × 14 mm, dark green. Stems 0.7–1.2 m, green with darker markings, with c.5 fls.

‘White Shadow’ V(b/a)

Parentage unknown

H: J.W.A. van der Wereld, pre-2003 **N:** Liliande BV, 2003 **REG:** J.W.A. van der Wereld, 2005

Tepals greenish white (155C); midveins green at base shading to white. Spots and papillae absent; nectaries green; pollen orange-yellow; stigma whitish grey.

Tepals long and of medium width to broad, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length to long and of medium width to broad. Stems very tall, green, with few fls. Mid-season.

WHITE TRIUMPH VIII(b/a) LO

Trade designation of ‘Zanlotriumph’.

‘White Valley’ VII(a/b)

(Un-named seedling × ‘Opus One’) × (*L.* × *parkmanii* ‘Allegra’ × ‘Starfighter’) F₂

H: Bischoff Tulleken Lelies BV, pre-2003

N: Bischoff Tulleken Lelies BV, 2003

REG: Mak Breeding BV, 2006

Tepals, including throat, yellowish white (155D); midveins pale greenish yellow (1D) up to half-way above nectaries; midribs of outer tepals strong yellow-green (144C) at base and top, shading to 144A between; midribs of inner tepals light yellow-green (144D) at base and top. Spots absent, papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen deep orange-yellow (163A); stigma light purplish grey (N187D). Fls 180 mm wide, scented; tepals 110 × 56 mm, margins slightly ruffled, tips recurved. Lvs 130 × 38 mm. Stems 0.75–0.85 m, green with darker markings, particularly at base, with c.5 fls.

'Wild Romance' VII(a-b/b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007
 Tepals yellowish white (155D). Spots absent; papillae present; nectaries green; pollen orange-brown; stigma purple, speckled. Tepals of medium length and width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and broad to very broad. Stems 1.2 m, green with darker markings, with up to 6 fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Winterberg' VII(a/b-c)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2006
 Tepals, including throat, yellowish white (155D); midveins tinged pale yellowish pink (159D); outside of outer tepals flushed 159D beside midribs; midribs of outer tepals strong yellow-green at base and top (143C and 144C respectively), shading to light yellow-green (145D) between; midribs of inner tepals strong yellow-green (144C) at base shading to brilliant yellow-green (150C). Midveins and midribs of young fls light yellow-green (150D). Spots absent, papillae yellowish white (155D); nectaries light yellow-green (144D); pollen dark reddish orange (172B); stigma yellowish grey (198D). Fls 260 mm wide, slightly scented; tepals 141 × 88 mm, margins ruffled, tips recurved. Lvs 225 × 48 mm. Stems 1.05–1.1 m, green, with *c.* 3 fls.

'Yellow Cab' I(a/b)

'Connecticut King' seedling × yellow-flowered seedling

H: Bischoff Tulleken Lelies BV, pre-2003
N: Bischoff Tulleken Lelies BV, 2003
REG: Mak Breeding BV, 2006
 Inside of inner tepals vivid yellow (14B); inside of outer tepals, and margins and above throat of inner tepals, vivid yellow (14A); throat brilliant greenish yellow (1B). Outside of outer tepals vivid yellow (12A) flushed strong yellow (153D), shading to vivid yellow (14A) towards margins; midribs strong yellow-green at base and top (143B and 144A respectively), flushed moderate red (181A) at base. Outside of inner tepals vivid yellow (13A); midribs strong yellow-green (144B). Spots dark red (59A); papillae vivid yellow (14A); nectaries moderate olive-green (137A); pollen dark orange (N163A); stigma dark red (59A). Buds slightly hairy. Fls 170 mm wide; tepals 100 × 58 mm, margins smooth, tips slightly recurved. Lvs 175 × 15 mm, dark green. Stems 1.1 m, dark, with *c.* 4 fls.

'Yellowknife NWT' I(a/c)

Un-named seedling hybridized by F. Fellner × 'Joyce's Favourite'
H: D. Crook, 2001 **G:** D. Crook, 2003
N: D. Crook, 2007 **I:** Valley K Greenhouses, post-2008 **REG:** D. Crook, 2007
 Inside of tepals vivid yellow (14B) in basal half, shading

to strong orange (26A) in upper half; throat vivid reddish orange (30A). Outside of tepals vivid yellow (13A); midribs vivid yellowish orange (30C). Spots of medium size, oblong, warm brown, evenly distributed either side of midrib over basal third of each tepal; papillae absent; nectaries vivid reddish orange (30A); pollen strong orange (170A); stigma moderate yellow (160A). Fls 134 mm wide; tepals 72 × 26 mm, margins smooth, tips slightly recurved; infl. racemose. Lvs 122 × 11 mm, scattered. Stems 1 m, greyish brown (166A), with up to 12 fls. Mid-July (NW Territories, Canada). Named after the capital city of Northwest Territories, Canada.

'Yellow Pixels' I(a/b-c)

'Cha Cha Cha' seedling × yellow-flowered seedling
H: Laan Flora Facilities BV, pre-2005 **N:** Laan Flora Facilities BV, 2005 **REG:** Laan Flora Facilities BV, 2006

Inside of tepals brilliant greenish yellow (6B); throat brilliant greenish yellow (1B). Outside of tepals brilliant greenish yellow (3A) shading to 5B towards margins; midribs of outer tepals strong yellow-green at base and top (144C and 144B respectively), with a little vivid reddish orange (N30C) at top; midribs of inner tepals brilliant greenish yellow (3A) shading to strong yellow-green (144C) at top. Spots numerous, inside of tepals heavily spotted dark red (59A) at base and along margins; spots on outside of tepals few, deep purplish red (59B) along margins; papillae absent; nectaries strong yellow-green (143B); pollen vivid orange-yellow (23A); stigma dark red (59A). Fls 120–145 mm wide; tepals 84 × 38 mm, margins smooth, tips straight. Lvs 114 × 15 mm. Stems 0.85–1.2 m, green with darker markings, with *c.* 9 fls.

'Yomaco' VII(a-b/b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007
 Inside of tepals deep purplish pink (N57D); margins darker towards base; midveins pale yellow-green (4D) at base. Outside of inner tepals strong purplish pink (63C). Spots and papillae present; nectaries pale green; pollen orange-brown; stigma purple, speckled. Tepals of medium length and width, margins ruffled to strongly ruffled, tips recurved. Lvs of medium length to long, and broad to very broad. Stems 1.5 m, green with darker spots and stripes, with few to average no. of fls. Late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Záletnice' I(a/b-c)

((('Redstart' × 'Nutmegger') × Wattle Bird) × ('Red Echo' × ('Connecticut King' × 'Tabasco'))) × 'Silver Bunny')

H: B. Mičulka, pre-2006 **N:** B. Mičulka, pre-2006 **I:** B. Mičulka, 2006 **REG:** B. Mičulka, 2006
 Inside of tepals pale beige, with a brownish tinge above nectaries; tips yellowish cream; throat whitish. Outside of tepals whitish; midribs old rose-pink and brownish. Spots few, fairly large, dark carmine-brown, on a quarter of each tepal; nectaries frosted white; pollen dark brown; stigma dark carmine-

brown. Fls 135 mm wide; tepals 85 × 40 mm, tips slightly recurved; pedicels 60 mm long, profusely spotted dark brown. Lvs 100 × 15 mm, dull dark green. Stems 0.85 m, profusely spotted brown, infl. fairly long. Mid-June.

‘Zanlagota’ VIII(a/b) LA

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2004

N: Van Zanten Flowerbulbs BV, 2004

REG: Van Zanten Flowerbulbs BV, 2006

Inside of tepals brilliant greenish yellow (inner 4A, outer 6B, midveins 5A); throat light greenish yellow (1C). Outside of outer tepals brilliant greenish yellow (7C), shading to brilliant yellow (7B) towards margins. Outside of inner tepals light greenish yellow (5D); margins brilliant greenish yellow (5B). All midribs strong yellow-green (143C) at base shading to brilliant greenish yellow (of outer tepals 7C, of inner 5A).

Spots and papillae absent; nectaries strong yellow-green (143B); pollen greyish brown (166A); stigma brilliant greenish yellow (1B) with deep purplish red (59B). Fls 147 mm wide; tepals 115 × 56 mm, margins smooth, tips not recurved. Lvs 155 × 33 mm. Stems 1–1.2 m, green, with up to 4 fls.

Trade designation BOGOTA

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Zanlapres’ VIII(a/b) LA

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2004

N: Van Zanten Flowerbulbs BV, 2004

REG: Van Zanten Flowerbulbs BV, 2006

Tepals yellowish white (155D); inside of tepals flushed strong yellow-green (143B) above throat; throat 144C. Outside of each outer tepal with broad strong purplish red (59D) stripe to one side; midribs strong yellow-green at base (144C) and top (144B); top with a darker (144A) spot; shading to light yellow-green (145C) between. Midribs of inner tepals strong yellow-green (144C) at base flushed strong purplish red (59D), shading to light yellow-green (145D, and 145C at top). Spots and papillae absent; nectaries strong yellow-green (143A); pollen moderate reddish brown (166B); stigma light greenish grey (190D) with small, strong purplish red (59D) stripe. Fls 200 mm wide; tepals 125 × 60 mm, margins smooth, tips slightly recurved. Lvs 136 × 25 mm. Stems 1.15–1.35 m, green with darker markings, with up to 7 fls.

Trade designation WHITE IMPRESSION

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Zanlorano’ VIII(a/b) LO

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2004

N: Van Zanten Flowerbulbs BV, 2004

REG: Van Zanten Flowerbulbs BV, 2006

Inside of tepals deep purplish red (59B), shading to yellowish white (155D) in uppermost third; narrow margins in basal two-thirds, and throat, yellowish white (155D). Outside of outer tepals flushed purple-red, for up to two-thirds, on yellowish white (155D)

ground, with strong purplish red (59D) stripes along midribs; midribs strong yellow-green at base (144C) and top (144B), shading to pale yellow-green (155A) between. Outside of inner tepals yellowish white (155D), with small strong purplish red (60C) stripes; up to half of margin strong purplish red (60C); midribs light yellow-green (144D) to strong yellow-green (144B) at base, shading to strong purplish red (60C), and yellowish white (155D) at top. Spots and papillae absent; nectaries strong yellow-green (144C); pollen vivid orange-yellow (23A); stigma greyish yellow-green (191B). Fls 225 mm wide, slightly scented; tepals 160 × 70 mm, margins smooth, tips recurved. Lvs 185 × 40 mm, dark green. Stems 1–1.2 m, green with darker markings at base, with up to 3 fls.

Trade designation CYRANO

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Zanlortara’ VII(a/b-c)

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2004

N: Van Zanten Flowerbulbs BV, 2004

REG: Van Zanten Flowerbulbs BV, 2006

Inside of tepals deep purplish pink (64D); inner tepals with small yellowish white (155D) stripes above nectaries, outer tepals yellowish white (155D) above nectaries; midveins deep purplish pink (of inner tepals N66D, of outer N66C); throat yellowish white (155D). Outside of tepals dark purplish pink (186C), tinged yellowish white (155D) at base beside midribs; midribs of outer tepals moderate yellow-green (148A) at base, shading to yellowish white (155D) flushed dark purplish pink (186C), and dark greenish yellow (152C) with a spot of strong yellow-green (143A) at top. Midribs of inner tepals light olive (152A) at base and top, shading to dark purplish pink (186C) between and flushed 186C at base. Spots absent; papillae moderate purplish pink (65A); nectaries strong yellow-green (144C); pollen strong reddish orange (169A) to moderate reddish orange (171A); stigma light greenish grey (188C). Fls 245 mm wide, slightly scented; tepals 140 × 75 mm, margins ruffled, tips recurved. Lvs 220 × 63 mm. Stems 1.15–1.4 m, green, with up to 7 fls.

Trade designation VITARA

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Zanlorum’ VII(a-b/b)

Un-named seedling × un-named seedling

H: Van Zanten Flowerbulbs BV, pre-2004

N: Van Zanten Flowerbulbs BV, 2004

REG: Van Zanten Flowerbulbs BV, 2006

Inside of tepals moderate red (185B), shading to light yellow (10B) above nectaries; margins closely spotted, and tips flushed, yellowish white (155D); inner tepals shading to deep red (60A) towards margins; throat yellowish white (155D). Outside of outer tepals dark red (59A), yellowish white (155D) beside midribs; midribs strong yellow-green at base (144C) and top (143A), shading to yellowish white (155D) and dark red (59A) between. Outside of inner tepals deep red (60A), shading to moderate purplish red (59C) at base

and along margins, and to yellowish white (155D) beside midribs at base; midribs strong yellow-green (144C) at base, shading to yellowish white (155D) and deep red (60A), and tinged light yellow-green (145C) at top. Spots dark red (59A); papillae yellowish white (155D) and dark red (59A); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma light grey (188D) with greyish reddish purple (N77C). Fls 175 mm wide, slightly scented; tepals 100 × 60 mm, margins slightly ruffled, tips recurved. Lvs 155 × 48 mm. Stems 1.1–1.3 m, green, with up to 7 fls. Trade designation **RED RUM**
Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Zanlotriumph' VIII(b/a) LO

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2004

N: Van Zanten Flowerbulbs BV, 2004

REG: Van Zanten Flowerbulbs BV, 2006

Tepals yellowish white (155D); midveins pale greenish yellow (1D) for up to half; midribs of outer tepals strong yellow-green at base (143C) and top (144A), shading to pale yellow-green (155A) between; midribs of inner tepals strong yellow-green (143C) at base, shading to pale greenish yellow (2D); throat light yellow-green (145C). Spots and papillae absent; nectaries strong yellow-green (143C); pollen vivid orange-yellow (23A); stigma very pale blue (122D). Buds yellow-green, flushed deep red (60A) along midrib. Fls 300 mm wide, slightly scented; tepals 195 × 75 mm, margins smooth, tips strongly recurved. Lvs 225 × 35 mm. Stems 1.2–1.5 m, green, with up to 3 fls. Trade designation **WHITE TRIUMPH**
Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Zanlovita' VIII(b-a/b) LO

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2004

N: Van Zanten Flowerbulbs BV, 2004

REG: Van Zanten Flowerbulbs BV, 2006

Inside of tepals pale yellow-green (157B) to greenish white (157D) in basal third, shading to strong reddish purple (70B) to dark purplish pink (186C) on paler ground; tips greenish white (155C) (for a larger area on outer tepals); throat greenish white (155C). Outside of outer tepals greenish white (157D) to pale yellow-green (4D), shading to dark purplish pink (186C) to moderate purplish pink (186D) towards margins; margins strong reddish purple (70B) to deep purplish pink (70C); midribs strong yellow-green at base (144B) and top (144A), with small moderate purplish red (186B) stripes on the right-hand-side of each midrib. Outside of inner tepals greenish white (157D), shading to dark purplish pink (186C) towards margins and tips; midribs strong yellow-green (144B) at base, shading to pale greenish yellow (2D) towards top. Spots and papillae absent; nectaries strong yellow-green (144B); pollen dark reddish orange (175B); stigma light yellow-green (145D). Buds green with strong purplish red (60B) stripes. Fls 225 mm wide; tepals 180 × 82 mm, margins smooth to slightly ruffled, tips recurved. Lvs 235 × 39 mm,

dark green. Stems 1.12–1.45 m, pale green with darker markings, with up to 5 fls.

Trade designation **NOVITA**

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Zidane' VII(a-b/b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2007

Tepals yellowish white (155D); midveins pale yellow-green (4D) at base. Spots absent; papillae present; nectaries green; pollen red-brown; stigma purple.

Tepals of medium length and of medium width to broad, margins ruffled to strongly ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and broad. Stems 1.55 m, green, with few to average no. of fls. Mid-season to late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Zlatokop' I(a-b/b)

'Encounter' × ('Apricot Supreme' × 'Capri')

polyploid seedling

H: B. Mičulka, pre-2006 **N:** B. Mičulka, pre-2006

I: B. Mičulka, 2006 **REG:** B. Mičulka, 2006

Tepals apricot, darker above nectaries; tips and central portion of outside of tepals, paler. Spots fairly large, oblong, dark red-brown, on two-fifths of each tepal; nectaries frosted white; pollen chocolate-brown; stigma brown-red. Fls 155 mm wide; tepals 100 × 43 mm, tips slightly recurved; pedicels 60 mm long, very stout, dark brown; with secondary buds. Lvs 135 × 21 mm, dull dark green. Stems 0.85 m, very stout, dark brown, infl. short. Late June.

'Zliňanka' VI(c/d)

((Palmer 582812 × 'Gold Eagle') × 'Mimosa Star') × (('Henrietina' × Palmer 56227) × unknown)

H: B. Mičulka, pre-2005 **N:** B. Mičulka, pre-2005

I: B. Mičulka, 2006 **REG:** B. Mičulka, 2005

Inside of tepals pale yellow with greenish flame, of velvet texture; throat darker yellow, gently furrowed, dusted cinnamon. Outside of tepals pale beige; midribs pale green. Pollen orange-brown; stigma green. Fls 145 mm wide; tepals 113 × 38 mm, tips recurved; pedicels 55 mm long, slender, sparsely spotted dark brown. Lvs 110 × 15 mm, dull mid-green. Stems up to 2 m, slender. Mid-July.

'Zvonařka' VI(c/b)

'Sancho' × unknown

H: B. Mičulka, pre-2006 **N:** B. Mičulka, pre-2006

I: B. Mičulka, 2006 **REG:** B. Mičulka, 2006

Inside of tepals dark beige-orange; outside of tepals paler orange; midribs mahogany. Spots and papillae absent; nectaries green; pollen orange-brown; stigma green. Fls 140 mm wide, bell-shaped; tepals 120 × 53 mm, tips recurved; pedicels 70 mm long, slender, spotted dark brown; with secondary buds. Lvs 140 × 16 mm, dull dark green. Stems 1.6 m, slender, supple, profusely spotted red-brown, infl. short. Early July.

Hybridizers, registrants and others

This list includes selectors, namers and introducers in addition to hybridizers and registrants.

(† = deceased)

- BERG en Zn, M.A., van den
't Zand, The Netherlands
- BISCHOFF TULLEKEN LELIES BV
Wieringerwerf, The Netherlands
- BLOOTHOOFD, D.
The Netherlands
- BOON, Fa Jaap
Hem, The Netherlands
- BOTTEMA BLOEMBOLLEN BV
Berkhout, The Netherlands
- BRAUN, K.
Rousínov, Czech Republic
- BURNETT, A.J.
Vermilion, Alberta, Canada
- CROOK, D.
Hay River, Northwest Territories, Canada
- DUDEK, J.
Hlučín, Czech Republic
- DUNVILLE, N.
Saskatoon, Saskatchewan, Canada
- DUNVILLE, S.T.
Saskatoon, Saskatchewan, Canada
- DUYN, C., van
Rijnsburg, The Netherlands
- EWALD, G.
Leiferde/Gifhorn, Germany
- FELLNER, F.
Vermilion, Alberta, Canada
- FIELDING, D.
West Pine, Tasmania, Australia
- FLEVO LILIES
Espel, The Netherlands
- †FOX, E. E.
See FOX LILY RANCH
- FOX LILY RANCH
Miller, Alberta, Canada
- GARTLEMAN, F.
Courtney, British Columbia, Canada
- GROOT & Zn, Fa W., de
Enkhuizen, The Netherlands
- HALL, R.
Chapel House, nr Newcastle upon Tyne, UK
- HARTLE-GILMAN GARDENS
Owatonna, Minnesota, USA
- HOBAGO BV
Lisse, The Netherlands
- HOFF, P.
Steenbergen, The Netherlands
- HOFFGAARDE BV
Steenbergen, The Netherlands
- HONEYWOOD LILIES
Parkside, Saskatchewan, Canada
- HORNER LILIES
Thorndale, Ontario, Canada
- HORNER, E.
See HORNER LILIES
- IMANSE, A.C.
Lisserbroek, The Netherlands
- INSTITUT NATIONAL de la RECHERCHE
AGRONOMIQUE
Paris, France
- JONG Beheer BV, De
Andijk, The Netherlands
- JONG LELIES HOLLAND BV, De
Andijk, The Netherlands
- KANEKO SEEDS
Japan
- KONINKLIJKE Van ZANTEN BV
Hillegom, The Netherlands
- LAAN FLORA FACILITIES BV
Zwaagdijk, The Netherlands
- LICOMBA
Bovenkarspel, The Netherlands
- LILLIANDE BV
't Zand, The Netherlands
- LYBELMEX HILLEGOM BV
Hillegom, The Netherlands
- MAK BV, Gebr.
't Zand, The Netherlands
- MAK BREEDING BV
Wieringerwerf, The Netherlands
- MAK-LEEK Inc.
Anna Paulowna, The Netherlands
- MAK 't ZAND BV
't Zand, The Netherlands
- MARKLILY CV
Lisse, The Netherlands
- MARTSCHINKE, Miss N.
Sulingen, Germany
- MEMORIAL UNIVERSITY OF
NEWFOUNDLAND BOTANICAL GARDEN
St John's, Newfoundland, Canada
- MICULKA, B.
Velehrad, Czech Republic
- MOERMAN, J.A. & J.
De Lier, The Netherlands
- NICHOLLS, W.
Memorial University of Newfoundland
Botanical Garden, St John's, Newfoundland,
Canada
- PAAUW en Zn BV, W.
Rijnsburg, The Netherlands
- PEACOCK BV
Rijnsburg, The Netherlands
- PORTER, A.J.
Honeywood Lilies, Parkside, Saskatchewan,
Canada
- RICHARDSON, B.
Nanaimo, British Columbia, Canada
- ROOYAKKERS LELIE BV
Breezand, The Netherlands
- ROYAL Van ZANTEN FLOWERBULBS BV
Hillegom, The Netherlands

S. en S., V.o.f.
 Berkhout, The Netherlands
 SANDE BV
 't Zand, The Netherlands
 SCHAEFER, Mrs D.B.
 Waukee, Iowa, USA
 STEINBRÜCK, E.
 Berlin, Germany
 STEINBRÜCK, G.
 Berlin, Germany
 TESTCENTRUM voor SIERGEWASSEN BV
 Hillegom, The Netherlands
 "THE ORIGINALS" U.A., COÖP.
 KWEKERSVERENIGING
 Breezand, The Netherlands
 †THOMAS, R.
 British Columbia, Canada
 TRIOR LELIE BV
 Hillegom, The Netherlands
 UNIVERSITY OF LATVIA BOTANICAL
 GARDEN
 Riga, Latvia
 VALLEY K GREENHOUSES
 Edberg, Alberta, Canada
 VEEN BV, J.P., van
 Noorden, The Netherlands
 VLETTER en Den HAAN Beheer BV
 Rijnsburg, The Netherlands
 VRIES VEREDELING BV De
 Breezand, The Netherlands
 WAARD en Zn, P.J., de
 Breezand, The Netherlands
 WERELD, J.W.A., van der
 Breezand, The Netherlands
 WESTFALL, L.
 See VALLEY K GREENHOUSES
 WILLOUGHBY, S.K.
 Spruce Grove, Alberta, Canada
 WILLOUGHBY, T.D.
 Spruce Grove, Alberta, Canada
 WORLD BREEDING BV
 Breezand, The Netherlands
 WORLD BREEDING BV and EOTA/KOSTER
 See WORLD BREEDING BV
 ZANTEN FLOWERBULBS BV, Van
 Hillegom, The Netherlands
 ZORGEVICS, Ā.
 University of Latvia Botanical Garden,
 Rīga, Latvia

ISBN 978-1-902896-97-7



9 781902 896977 ▶